



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





math. Lib

QA

374

.E124





PARTIAL DIFFERENTIAL EQUATIONS.

PARTIAL DIFFERENTIAL EQUATIONS.

Cambridge:
PRINTED BY C. J. CLAY, M.A.
AT THE UNIVERSITY PRESS.

9065-

PARTIAL DIFFERENTIAL EQUATIONS.

AN ESSAY



TOWARDS AN ENTIRELY

NEW METHOD OF INTEGRATING THEM.


BY

S. EARNSHAW, M.A.

OF ST JOHN'S COLLEGE, CAMBRIDGE.

London and New York:
MACMILLAN AND CO.
1871.

[All Rights reserved.]



PREFACE.

THE Author of this Essay has been forewarned that the adoption of the subscript differential notation, which forty years ago was in some degree of favour but afterwards laid aside, will prove a serious hinderance to the acceptableness of the present Work, as it will be new to the generality of mathematical students, and therefore somewhat repellent. He regrets the circumstance, and feels that an explanation and apology are needed; and he finds both in the fact that the notation alluded to has been forced upon him by the necessities of the case; and, moreover, he ventures to think that the reader after going through the Essay will agree with him in this, and be willing on this ground to excuse such an important deviation from established mathematical usage.

The Author denominates his Work an ESSAY, because as a TREATISE it would be very incomplete, there being many portions of the general subject not alluded to in these pages. For information on those portions he must refer the reader to Boole and Gregory and Carmichael, with whose admirable and valuable Works he has been careful to interfere as little

as possible. In fact, he has consulted them only for such illustrative examples as fell within the scope of his Essay. He is conscious of having apparently intermeddled with the integration of equations of *one* independent variable; but this has arisen out of the peculiarities of the Author's system, by which every equation, whatever be the number of its independent variables, is reduced to an equivalent equation of only one independent variable. It was therefore impossible to keep the two branches of differential equations apart.

The system of integration here proposed occurred to the mind of the Author a few years ago, but his professional engagements did not then leave him leisure to follow the general idea into its details. And, as his object is merely to render his Method thoroughly intelligible, and not the exhibition of integrals, he has for brevity's sake not always proceeded to the last steps of an integration when he conceived that the method had been made intelligible.

The Author fears a cursory glance at the pages of the Work will have a prejudicial effect; for he is aware that some of them exhibit a formidable and deterrent array of novel symbols; he therefore begs to assure the reader that the various steps of the investigations are all obedient to one general principle, and though in some degree novel, are not really difficult, but on the contrary easy when the eye has become accustomed to the novelties of the notation. And, moreover, he entertains a hope that the results of integrations (many of which are far more general than they were in the shape in which they have appeared in former Treatises) will repay the reader for any extra trouble he may find in pushing through what may at first appear to

him a forest of operative symbols. Many equations will be found in this Essay integrated with ease in finite terms, which, as far as the Author is aware, were never integrated in finite terms before. In the last Chapter, for instance, the following highly important equations will be found integrated for the first time in finite terms:

$$\frac{d^2u}{dx dy} + cu = 0;$$

$$\frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} + cu = 0;$$

and
$$\frac{d^2u}{dx^2} + \frac{d^2u}{dy^2} + \frac{d^2u}{dz^2} = 0.$$

Three more important partial differential equations could hardly be instanced, and a finite integral of each one of them is obtained in this Essay without any difficulty.

The Author can scarcely hope that in a work so novel in its methods as this is, and written as this has been under considerable pressure of other engagements, some clerical errors will not have escaped his notice; but he hopes that they will not prove to be many, and that none of them will involve errors of principle. If any are found, he will be thankful if the courteous reader will send notice of them addressed to him at the Publishers.

SHEFFIELD,

October 2, 1871.

ERRATA.

Page 63, line 19, *for question read equation.*

„ 69, „ 18, *for f read F .*

„ 70, bottom line, *for f read F .*

AN ESSAY.

CHAPTER I.

GENERAL PRINCIPLES OF THE ESSAY.

It is known that the integration of differential equations is oftentimes much facilitated by a change of one or more of the independent variables, and sometimes also by changing the dependent variable. The process is however often tedious, and it is frequently far from obvious what ought to be chosen as the new variables, even when it is seen that a change should be made.

The system of integration to be proposed in this Essay is founded on a simple principle of choosing new variables; and the change of variables is effected with great ease. And the most important feature of the system is, that whatever be the number of original independent variables, the work of integration is at once reduced to the use of one independent variable only.

Quasi-constants.

1. If $t, x, y, z \dots$ be n independent variables, and u the dependent variable, it is understood that $d_t u, d_x u, d_y u, d_z u, \dots$ are *partial* differential coefficients; by which is meant that in finding the value of $d_t u$, i.e. in operating on u with the symbol d_t , we are to consider $x, y, z, \dots$ as *constants*: in operating on u with d_x , then $t, y, z, \dots$ are to be considered as

constants; and so on. It is here proposed to express this briefly by the use of the word *quasi-constant*. Thus in reference to the use of the symbol d_t , we shall denominate $x, y, z, \dots$ and any functions of them *quasi-constants*: the use of d_t will imply that $t, y, z \dots$ and their functions are quasi-constants; and so on.

2. If s_t be the symbol of integration corresponding to d_t , the symbol of differentiation, then we know that after the operation s_t we must add to complete the integral an arbitrary function of $x, y, z, \dots$ its quasi-constants: and this principle is general and may be thus stated,—after an operation of integration we are to add to the integral *an arbitrary function of the corresponding quasi-constants*; and all the variables which were considered constants in differentiation are to be considered quasi-constants in integration.

Change of the independent variables.

3. If from the (n) independent variables $t, x, y, z, \dots$ we wish to pass to a new set $\tau, \xi, \eta, \zeta \dots$, formulæ for this purpose are ready to hand in the *Differential Calculus*. They are of the following type:

$$(d_\tau u) = d_t u \cdot (d_\tau t) + d_x u \cdot (d_\tau x) + d_y u \cdot (d_\tau y) + \dots (1).$$

This is only one of the (n) requisite equations, and the other $(n-1)$ equations may be obtained from this by writing successively in this $\xi, \eta, \zeta \dots$ for τ . But it is possible, whenever the forms of $\xi, \eta, \zeta \dots$ are at our disposal, to assign them such values in terms of $t, x, y, z, \dots$ as shall render the above equation (1) sufficient by itself for our purpose; so that by a proper choice of values for $\xi, \eta, \zeta \dots$ any partial differential equation of n variables may be reduced to an equivalent equation of *one variable* and $(n-1)$ *quasi-constants*. This is the principle on which this Essay is founded. The method of

assigning proper forms for the quasi-constants $\xi, \eta, \zeta, \dots$ in terms of $t, x, y, z, \dots$ the original independent variables will be best learned from the examples which will be given.

Distinctive Symbols of operation for different sets of independent variables.

4. In the formula (1) certain terms are enclosed in brackets; this is done to indicate that before the differential operations indicated in them are performed, $t, x, y, \dots$ must be expressed in terms of the new variables; or, that the operations are to be performed on the supposition that u is so expressed. In brief, the distinction of brackets and no brackets in equation (1) refers to the fact that there are operations to be performed in that equation in reference to two sets of independent variables. Had we to write down an equation in which differential and integral operations would have to be performed in reference to three or more sets of independent variables, we should have to employ two or more kinds of distinctive brackets to indicate that fact. Now it is proposed in this Essay to indicate this fact by employing distinctive differential and integral symbols of operation, and so to abolish the distinction by brackets. Thus, when $\tau, \xi, \eta, \zeta, \dots$ is the set of independent variables referred to in a differential or integral operation, we shall use D and S as the symbols of operation; if $\tau', \xi', \eta', \zeta', \dots$ be the set of independent variables, we shall indicate that fact by using Δ and Σ as the symbols of operation; and so on for any number of sets. Under this system equation (1) of Art. 3 will be written without brackets thus,—

$$D_{\tau}u = d_{\tau}u \cdot D_{\tau}t + d_x u \cdot D_{\tau}x + d_y u \cdot D_{\tau}y + \dots \dots \dots (1).$$

5. In speaking about different sets of independent variables, it is to be noticed that it is not necessary that every individual member of one set should be different from every

individual member of another set. If *some*, even if *one*, of one set be different from all of another set this is sufficient to establish them as different sets, and to entitle them to the use of distinctive differential and integral symbols of operation.

6. In fixing the values of a new set of independent variables in terms of the original set, it is of importance to take care that each member of the new set be really an *independent* variable; i.e. that it be such as cannot be expressed in terms of the other members of its own set. Unless this condition of independence be observed, we shall be working under an erroneous hypothesis.

7. Under the authority of Art. 5 we are at liberty to assume, whenever it can be done usefully, that one of the variables of each new set shall be identical with one of the variables of the original set,—for example, we may assume $\tau = t$, in which case equation (1) of Art. 4 becomes

$$D_t u = d_x u + d_x u . D_t x + d_y u . D_t y + \dots \dots \dots (1).$$

But when we use this equation, which will always be known from the occurrence of the symbol D_t or S_t , we must remember that there remain only $(n-1)$ new independent variables (the quasi-constants) of the set to be determined, one of them (t) being already fixed upon in using this equation.

8. In determining what shall be the forms of the constituent members of a new set of variables, we are at liberty to introduce n *arbitrary* hypotheses if we use equation (1) of Art. 4; but only $(n-1)$ arbitrary hypotheses if we employ equation (1) of the last Article. The arbitrariness of these hypotheses is however overruled by the necessity that the resulting members of a set of variables determined from them must be independent (Art. 6).

CHAPTER II.

DETERMINATION OF QUASI-CONSTANTS. EQUATIONS OF THE FIRST ORDER.

Ex. 1. To integrate $d_t u + a d_x u = bu$.

Let us change the independent variables to t, ξ . Then the formula of Art. 7 becomes for this case, which is one of two independent variables,

$$D_t u = d_t u + D_x x \cdot d_x u \dots\dots\dots (1).$$

We are at liberty to make *one* hypothesis (Art. 8) for the determination of the value of ξ ; let us assume $D_x x = a$. We make this choice, because by this assumption the right-hand member of the formula (1) will be rendered identical with the left-hand member of the proposed equation; so that we shall have $D_t u = bu$;

$$\therefore D_x x = a, \text{ and } D_t u = bu,$$

and these two equations, taken simultaneously, are together exactly equivalent to the proposed equation. Integrating them we have (see Art. 2)

$$x = at + \xi, \text{ and } e^{-bu} = F(\xi).$$

The former gives the value of ξ ; and this value being written

in the latter, we have the following as a relation that must always exist between u and the independent variables t, x ,

$$u = e^{at} F(x - at).$$

This is therefore the required integral.

9. REMARK. According to Art. (2) we ought to have written $x = at + f(\xi)$ instead of $x = at + \xi$, to complete the integration of $D_x x = a$: but had we done so the final integral would not have been affected thereby: and we were moreover at liberty to take either $x - at$, or any function of $x - at$ as the value of ξ . Under this license no additional generality would be gained by the use of $f\xi$ instead of ξ . We shall always make use of this liberty by adding either ξ , or a function of ξ , after *one* of the integrations in any proposed problem.

Ex. 2. To integrate $d_t u + a d_x u + b d_y u = cu$.

In this case, which is one of three independent variables, we have to change the variables to t, ξ, η ; and the formula of Art. 7 becomes

$$D_t u = d_t u + D_x x \cdot d_x u + D_y y \cdot d_y u.$$

We are here at liberty to make two hypotheses for the determination of the quasi-constants ξ and η . We therefore assume $D_x x = a$, $D_y y = b$; which gives $D_t u = cu$. These three equations, when taken simultaneously, are equivalent to the proposed equation.

We shall employ the first and second for fixing the values of ξ and η ;

$$\therefore x = at + \xi, \quad y = bt + \eta, \quad \text{and} \quad e^{-at} u = F(\xi, \eta).$$

And by substituting in the last of these the values of ξ and η found from the other two we find, as the required integral,

$$u = e^{at} F(x - at, y - bt).$$

Ex. 3. To integrate $d_t u + \frac{t}{x} \cdot d_x u = \frac{tu}{x^2}$.

Comparing this, which is a case of two independent variables, with $D_t u = d_t u + D_x x \cdot d_x u$, we find the two following equations which when taken simultaneously are equivalent to the proposed equation;

$$D_x x = \frac{t}{x}, \text{ and } D_t u = \frac{tu}{x^2}.$$

The former by integration gives $x^2 = t^2 + \xi$, which fixes the value of ξ : and the two equations combined give

$$x D_t u - u D_x x = 0,$$

which being integrated gives $\frac{u}{x} = F(\xi)$;

$$\therefore u = x F(x^2 - t^2).$$

Ex. 4. To integrate $d_t u + \frac{u}{t} \cdot d_x u = \frac{\alpha^2 x}{t}$.

Comparing this with the formula $D_t u = d_t u + D_x x \cdot d_x u$, we have

$$D_x x = \frac{u}{t}, \text{ and } D_t u = \frac{\alpha^2 x}{t};$$

$$\therefore u D_t u - \alpha^2 x D_x x = 0;$$

$$\therefore u^2 - \alpha^2 x^2 = \xi.$$

Also
$$D_x x = \frac{u}{t} = \frac{\sqrt{\alpha^2 x^2 + \xi}}{t};$$

$$\therefore \log(ax + \sqrt{\alpha^2 x^2 + \xi}) = a \log t + \log F(\xi);$$

$$\therefore ax + u = t^a \cdot F(u^2 - \alpha^2 x^2).$$

Ex. 5. To integrate $d_t u + \frac{x}{t} \cdot d_x u + \frac{a}{t} \cdot d_y u = \frac{nu}{t}$.

Comparing this, which is a case of three independent variables, with

$$D_t u = d_t u + D_x x \cdot d_x u + D_y y \cdot d_y u,$$

we have

$$D_x x = \frac{x}{t}, \quad D_y y = \frac{a}{t}, \quad \text{and} \quad D_t u = \frac{nu}{t}.$$

These equations being integrated give

$$x = t\xi, \quad \epsilon^{\frac{y}{a}} = t\eta, \quad \text{and} \quad t^nu = F(\xi, \eta);$$

$$\therefore u = t^nu F\left(\frac{x}{t}, \frac{1}{t} \cdot \epsilon^{\frac{y}{a}}\right).$$

Ex. 6. To integrate $d_x V \cdot d_t u - d_t V \cdot d_x u = 0$.

Divide by $d_x V$, and compare with

$$D_t u = d_t u + D_x x \cdot d_x u;$$

$$\therefore D_x x = -\frac{d_t V}{d_x V}, \quad \text{and} \quad D_t u = 0.$$

The latter gives by integration $u = \xi$, and the former may be written in the form $0 = d_t V + D_x x \cdot d_x V$ which is $= D_t V$. And hence, by integration,

$$V = F(\xi) = F(u).$$

Hence, in an equation of the form proposed, the dependent variables u , V are necessarily functions of one another. We may write the integral under the form

$$f(u, V) = 0.$$

Ex. 7. To integrate $d_t u - \frac{x}{t} \cdot d_x u + \frac{y}{t} \cdot d_y u = \frac{y^2}{t^2 x}$.

Comparing this equation with

$$D_t u = d_t u + D_x x \cdot d_x u + D_y y \cdot d_y u,$$

we have

$$D_t x = -\frac{x}{t}, \quad D_t y = \frac{y}{t}, \quad \text{and} \quad D_t u = \frac{y^2}{t^2 x}.$$

The first two give $tx = \xi$ and $y = t\eta$, and these being used in the third to eliminate x and y , we have

$$D_t u = \frac{\eta^2}{\xi} \cdot t, \quad (\xi, \eta \text{ are quasi-constants});$$

$$\begin{aligned} \therefore u &= \frac{\eta^2}{\xi} \cdot \frac{t^2}{2} + F(\xi, \eta) \\ &= \frac{y^2}{2tx} + F\left(tx, \frac{y}{t}\right). \end{aligned}$$

Ex. 8. To integrate $d_t u - \frac{x}{t} \cdot d_x u + \frac{y}{x} d_y u = \frac{y^2}{tx}$.

In this example,

$$D_t x = -\frac{x}{t}, \quad D_t y = \frac{y}{x}, \quad \text{and} \quad D_t u = \frac{y^2}{tx};$$

$$\therefore \xi = tx, \quad \eta = ye^{-\frac{t}{x}}, \quad \text{and} \quad D_t u = \frac{\eta^2}{\xi} \cdot \frac{t^2}{\xi};$$

$$\begin{aligned} \therefore u &= \frac{\eta^2}{\xi} \cdot S_t \epsilon \frac{t^2}{\xi} + F(\xi, \eta) \\ &= \frac{y^2}{tx} \cdot e^{-\frac{t}{x}} S_t \epsilon \frac{t^2}{\xi} + F(tx, ye^{-\frac{t}{x}}). \end{aligned}$$

Ex. 9. To integrate $d_t u + a d_x u + \frac{bu}{ht+kx} = 0$.

Comparing this with the usual formula we have $D_t x = a$, and $(ht+kx) D_t u = -bu$. Hence

$$x = at + \xi; \quad \text{and} \quad D_t (ht+kx) = h+ka.$$

It still remains to integrate

$$(ht + kx) D_t u + bu = 0.$$

Multiply by $(ht + kx)^{m-1}$, m being at present an unknown constant;

$$\therefore (ht + kx)^m D_t u + b (ht + kx)^{m-1} u = 0.$$

We see at a glance that this will be an exact differential, if m be such that $b = m D_t (ht + kx)$, that is, if

$$b = m(h + ka);$$

$$\therefore (ht + kx)^m u = F(\xi) = F(x - at).$$

Ex. 10. To integrate $d_t u + \frac{x}{t} \cdot d_x u = \frac{au}{ht + kx}$.

In this example,

$$D_t x = \frac{x}{t}, \quad \text{and} \quad D_t u = \frac{au}{ht + kx};$$

$$\therefore x = \xi t, \quad \text{and} \quad (ht + kx) D_t u = au;$$

$$\therefore t D_t u = \frac{au}{h + k\xi} \quad (\xi \text{ is a quasi-constant});$$

$$\therefore u = t^{\frac{a}{h+k\xi}} \cdot F(\xi)$$

$$= t^{\frac{at}{ht+kx}} \cdot F\left(\frac{x}{t}\right).$$

Ex. 11. To integrate

$$d_t u + a d_x u + b d_y u = c, \quad \text{and} \quad d_t u + a' d_x u + b' d_y u = c'$$

simultaneously.

Proceeding with the former of these as before, we have

$$D_t x = a, \quad D_t y = b, \quad D_t u = c;$$

which being integrated give

$$x = at + \xi, \quad y = bt + \eta, \quad \text{and} \quad u = ct + F(\xi, \eta) \\ = ct + F(x - at, y - bt).$$

Now substitute the value of u , given in the last equation, in the second of the simultaneous equations, using F for $F'(x - at, y - bt)$ for brevity,

$$\therefore (a - a') d_{\xi} F + (b - b') d_{\eta} F = c - c'.$$

This is an ordinary differential equation of the first order with constant coefficients, and being integrated as in Examples preceding gives

$$F = f\{(b - b') \xi + (a' - a) \eta\} + \frac{c\xi}{a - a'} - \frac{c'\eta}{b - b'},$$

$$\therefore u = \left(\frac{a'c}{a' - a} + \frac{bc'}{b' - b} \right) t + f\{(b - b')x + (a' - a)y + (ab' - ba')t\}.$$

10. In equations of the first order, if the coefficients be of a symmetrical form, it will sometimes be found to simplify the process of integration if we use the formula of Art. 4, instead of that of Art. 7.

Ex. 12. To integrate $(x + t) d_t u + (x - t) d_x u = au$.

Comparing this with

$$D_t u = d_t u, \quad D_t t = 1, \quad d_x u = d_x u, \quad D_t x = 0,$$

we have the following equations,

$$D_t u = au, \quad D_t t = x + t, \quad D_t x = x - t.$$

From the last two we obtain

$$t D_t t + x D_t x = t^2 + x^2,$$

and

$$x D_t t - t D_t x = t^2 + x^2;$$

$$\therefore \log \sqrt{t^2 + x^2} = \tau + \phi_1 \xi, \text{ and } \tan^{-1} \frac{t}{x} = \tau + \phi_2 \xi.$$

Also, the integration of the first gives

$$\log u = a\tau + \phi_3 \xi.$$

Between these three equations we have to eliminate τ and ξ ;

$$\therefore \log u - a \log \sqrt{t^2 + x^2} = \phi_3 \xi - a\phi_1 \xi,$$

and $\tan^{-1} \frac{t}{x} - \log \sqrt{t^2 + x^2} = \phi_2 \xi - \phi_1 \xi.$

Now the right-hand members of these equations being arbitrary functions of ξ , the left-hand members must also be arbitrary functions of each other;

$$\therefore u = (t^2 + x^2)^{\frac{a}{2}} F \left(\tan^{-1} \frac{t}{x} - \log \sqrt{t^2 + x^2} \right).$$

Ex. 13. To integrate

$$(bu - cx) d_t u + (ct - au) d_x u = ax - bt.$$

Comparing this with the formula of Art. 4, we have

$$D_t t = bu - cx, \quad D_t x = ct - au, \quad \text{and} \quad D_t u = ax - bt;$$

$$\therefore aD_t t + bD_t x + cD_t u = 0, \quad \text{and} \quad tD_t t + xD_t x + uD_t u = 0.$$

$$\therefore at + bx + cu = \xi, \quad \text{and} \quad t^2 + x^2 + u^2 = F(\xi);$$

$$\therefore t^2 + x^2 + u^2 = F(at + bx + cu),$$

which is the integral required.

Ex. 14. To integrate

$$(x + y + u) d_t u + (y + t + u) d_x u + (t + x + u) d_y u = t + x + y.$$

$$\therefore D_t t = x + y + u,$$

$$D_t x = y + t + u,$$

$$D_t y = t + x + u,$$

$$\text{and } D_t u = t + x + y;$$

$$\therefore \frac{D_t t - D_t x}{t - x} = \frac{D_t x - D_t y}{x - y} = \frac{D_t y - D_t u}{y - u} = -1,$$

$$\text{and } D_t (t + x + y + u) = 3(t + x + y + u).$$

Integrating these equations, and writing Q for $t + x + y + u$ for brevity, we have

$$t - x = \epsilon^{-\tau} F_1(\xi, \eta), \quad x - y = \epsilon^{-\tau} F_2(\xi, \eta), \quad y - u = \epsilon^{-\tau} F_3(\xi, \eta),$$

$$\text{and } Q = \epsilon^{3\tau} F_4(\xi, \eta).$$

Eliminating τ from these integrals, we find that

$$(t - x) \sqrt[3]{Q}, \quad (x - y) \sqrt[3]{Q}, \quad (y - u) \sqrt[3]{Q}$$

are (each of them) arbitrary functions of ξ, η ; consequently each of them is an arbitrary function of the other two.

$$\therefore 0 = F\{(t - x) \sqrt[3]{Q}, (x - y) \sqrt[3]{Q}, (y - u) \sqrt[3]{Q}\}.$$

Ex. 15. To integrate $t^2 du + x^2 dx = atx$.

Comparing this with the formula of Art. 4, we have

$$D_t u = atx, \quad D_t t = t^2, \quad D_t x = x^2,$$

From the last two of these we find $x^2 D_t t = t^2 D_t x$,

$$\therefore \frac{1}{t} = \frac{1}{x} + \xi;$$

$$\therefore D_t u = ax^2 \cdot \frac{t}{x} = \frac{a D_t x}{1 + \xi x};$$

$$\begin{aligned}\therefore u &= \frac{a}{\xi} \cdot \log(1 + \xi x) + F\xi \\ &= \frac{atx}{x-t} \cdot \log \frac{x}{t} + F\left(\frac{1}{t} - \frac{1}{x}\right),\end{aligned}$$

which is the required integral.

Ex. 16. To integrate

$$(x^3t - 2t^4) d_t u + (2x^4 - t^3x) d_x u = 9(t^3 - x^3) u.$$

Comparing this with the formula of Art. 4, we have

$$D_t u = 9(t^3 - x^3) u, \quad D_t t = x^3 t - 2t^4, \quad D_t x = 2x^4 - t^3 x.$$

$$\therefore \frac{D_t u}{3u} + \frac{D_t t}{t} + \frac{D_t x}{x} = 0,$$

$$\text{and } \frac{x D_t t - 2t D_t x}{x^3} + \frac{t D_t x - 2x D_t t}{t^3} = 0;$$

$$\therefore t^3 x^3 u = F(\xi), \text{ and } \frac{t}{x^3} + \frac{x}{t^3} = \xi;$$

$$\therefore t^3 x^3 u = F\left(\frac{t}{x^3} + \frac{x}{t^3}\right)$$

which is the integral required.

CHAPTER III.

DETERMINATION OF QUASI-CONSTANTS. EQUATIONS OF THE SECOND AND HIGHER ORDERS.

11. SINCE $D_i u = d_i u + D_i x . d_x u + D_i y . d_y u + \dots$ it follows that the single operation denoted by D_i is equivalent to the compound system of operations denoted by

$$d_i + D_i x . d_x + D_i y . d_y + \dots$$

Hence if $D_i x, D_i y, \dots$ are either constants, or quasi-constants, any function of the one operation D_i will be equivalent to the same function of the compound system

$$d_i + D_i x . d_x + D_i y . d_y + \dots$$

$$\therefore F(D_i) = F(d_i + D_i x . d_x + D_i y . d_y + \dots) \dots \dots (1).$$

Ex. 1. To integrate

$$d_i^2 u + 2ad_i d_x u + a^2 d_x^2 u + b(d_i u + ad_x u) + cu = 0.$$

Taking the terms of the second order, we have

$$d_i^2 + 2ad_i d_x + a^2 d_x^2 = (d_i + ad_x)^2.$$

We therefore assume $D_i u = d_i u + ad_x u$, which compared with the general formula gives

$$D_t x = a; \text{ and } \therefore D_t^2 u + b D_t u + c u = 0;$$

$$\begin{aligned} \therefore x = at + \xi, \text{ and } u &= e^{mt} F(\xi) + e^{nt} f(\xi) \\ &= e^{mt} F(x - at) + e^{nt} f(x - at), \end{aligned}$$

m and n being the two roots of $m^2 + bm + c = 0$.

Ex. 2. To integrate

$$d_t^2 u - a^2 d_x^2 u + b(d_t u + a d_x u) = 0.$$

Since $d_t^2 - a^2 d_x^2 = (d_t + a d_x)(d_t - a d_x)$, we shall have (here for the first time) to use *two* new sets of independent variables, corresponding to the two equations,

$$D_t u = d_t u + a d_x u, \text{ and } \Delta_t u = d_t u - a d_x u.$$

These give $D_t x = a$, and $\Delta_t x = -a$; and since the symbols D_t , Δ_t are respectively equivalent to the compound symbols $d_t + a d_x$, $d_t - a d_x$, the proposed equation becomes

$$D_t \Delta_t u + b D_t u = 0.$$

Hence the following system of three simultaneous equations is equivalent to the proposed equation,

$$D_t x = a, \Delta_t x = -a, \text{ and } D_t \Delta_t u + b D_t u = 0 \dots \dots (1).$$

The first two give by integration,

$$x = at + \xi, \quad x = -at + \xi';$$

and the third being integrated with S_t gives

$$\Delta_t u + b u = F(\xi) = F(x - at) \dots \dots \dots (2).$$

But the symbols D_t and Δ_t are commutative because their equivalents $d_t + a d_x$, $d_t - a d_x$ are so; consequently the equation (1) may be written in the form

$$\Delta_t D_t u + b D_t u = 0,$$

which being integrated with Σ_t gives

$$D_t u = \epsilon^{-at} f(\xi') = \epsilon^{-at} f(x + at).$$

Adding this equation to equation (2), and writing u for the sum of the left-hand members, we obtain finally

$$u = F(x - at) + \epsilon^{-at} f(x + at),$$

for the integral of the proposed equation.

12. The following is the reason why we may write u for $D_t u + \Delta_t u + bu$.

Since the proposed equation is represented by

$$D_t \Delta_t u + b D_t u = 0 \dots\dots\dots(1),$$

differentiate this equation separately with D_t and Δ_t ; and also multiply it by b ; the results are (since D_t and Δ_t are commutative)

$$D_t \Delta_t \cdot D_t u + b D_t \cdot D_t u = 0,$$

$$D_t \Delta_t \cdot \Delta_t u + b D_t \cdot \Delta_t u = 0,$$

$$D_t \Delta_t \cdot bu + b D_t \cdot bu = 0;$$

$$\therefore D_t \Delta_t \cdot (D_t u + \Delta_t u + bu) + b D_t \cdot (D_t u + \Delta_t u + bu) = 0.$$

This equation is of the same form as (1), and we learn from it that $D_t u + \Delta_t u + bu$ will satisfy the proposed equation quite as well as u will do so. If therefore $D_t u + \Delta_t u + bu$ contain the requisite number of arbitrary functions, it is a complete integral of the proposed equation, and being so we may write u for it.

We shall find this principle of great use hereafter; but it must be noticed that it only applies to cases of linear equations with constant or quasi-constant coefficients and when the symbols of operation are commutative.

13. It is well known that the integral of any linear differential equation of the form $F(d_t, d_x, d_y, \dots)u = W$, where W is a function of $t, x, y, \dots$, consists of two independent parts; one of them independent of W is called the absolute part of the integral; the other is dependent on W . These two parts being independent of each other, may be found separately (should it be advantageous to do so); and their sum will constitute the complete integral. The absolute part corresponds to the general supposition, $W = 0$.

As an example of this principle, let it be required to integrate the equation $d_t^2 u - a^2 d_x^2 u + b(d_t u + a d_x u) = W$.

We first find, as in the preceding example, the absolute integral, i.e. that part of the complete integral which corresponds to $W = 0$. This has already been found to be

$$u = e^{-bx} f(x + at) + F(x - at).$$

To find the part that depends on W we have the equation

$$D_t \Delta_t u + b D_t u = W, \text{ or } \Delta_t D_t u + b D_t u = W.$$

Either of these (whichever may be judged most convenient) will furnish the required result. For integrating the former with S_t , or the latter with Σ_t , we have

$$\Delta_t u + bu = S_t W, \text{ or } D_t u = e^{-bx} \Sigma_t e^{bx} W.$$

Integrate, now, the former with Σ_t , or the latter with S_t ;

$$\therefore u = e^{-bx} \Sigma_t e^{bx} S_t W, \text{ or } u = S_t e^{-bx} \Sigma_t e^{bx} W.$$

Adding to either of these the absolute part of the integral, we have for the complete result

$$u = e^{-bx} \Sigma_t e^{bx} S_t W + e^{-bx} f(x + at) + F(x - at),$$

$$\text{or } u = S_t e^{-bx} \Sigma_t e^{bx} W + e^{-bx} f(x + at) + F(x - at).$$

14. From the results just obtained we perceive that the symbols $\epsilon^{-n}\Sigma_i\epsilon^n$ and S_i are commutative in the preceding example. It will be found always necessary to notice whether the symbols of operation in any proposed example are commutative, for the commutateness of the symbols of operation is a property of immense use in facilitating the integration of equations; and in some instances failure arises solely out of the want of commutateness.

15. In seeking the absolute portion of an integral, the following property will be found of great use. It is applicable in all linear equations with constant coefficients; and its proof depends on the property proved in Art. 12.

If $d_i^m D_i^n \Delta_i^p u = 0$, and the symbols d_i , D_i , Δ_i are known to be commutative, then $u = s_i^m O + S_i^n O + \Sigma_i^p O$.

For since $d_i^m D_i^n \Delta_i^p u = 0$ is a linear equation with constant coefficients, operate on it separately with s_i^m , S_i^n , Σ_i^p ; the results are

$$\therefore D_i^n \Delta_i^p u = s_i^m O, \quad d_i^m \Delta_i^p u = S_i^n O, \quad d_i^m D_i^n u = \Sigma_i^p O.$$

Add these equations together, writing u for the sum of their left-hand members;

$$\therefore u = s_i^m O + S_i^n O + \Sigma_i^p O.$$

Now this value of u contains $m+n+p$ arbitrary functions, viz. m corresponding to the symbol s_i^m , n to the symbol S_i^n , and p to the symbol Σ_i^p ; and consequently it is the complete integral of the proposed equation, which is a differential equation of the $(m+n+p)^{\text{th}}$ order.

It will have been noticed in the above proof, that the commutateness of the symbols d_i , D_i , Δ_i has been assumed; and this assumption was made on the ground that the pro-

posed equation $d_i^m D_i^n \Delta_i^p u = 0$ represents a linear equation with constant coefficients.

16. In a similar way it may be shewn that if F, G, H be any symbols whatever of integration,

$$F.G.H.O = F.O + G.O + H.O,$$

provided the proper number of arbitrary functions be furnished by the right-hand member, and F, G, H be commutative.

Ex. 3. To integrate the equation

$$d_i^2 u + (a+b)d_x d_x u + abd_x^2 u = W.$$

The left-hand member being equal to $(d_i + ad_x)(d_i + bd_x)u$, we have $D_i = d_i + ad_x$, $\Delta_i = d_i + bd_x$; and therefore $\xi = x - at$, $\xi' = x - bt$; and the proposed equation may be written

$$D_i \Delta_i u = W + O;$$

$$\therefore u = S_i \Sigma_i W + S_i O + \Sigma_i O, \quad (\text{Art. 15})$$

$$= S_i \Sigma_i W + F\xi + f\xi'$$

$$= S_i \Sigma_i W + F(x - at) + f(x - bt).$$

Ex. 4. To integrate $d_i^2 u - a^2 d_x^2 u = W$.

Since $d_i^2 u - a^2 d_x^2 u = (d_i + ad_x)(d_i - ad_x)u$, we have $D_i = d_i + ad_x$, $\Delta_i = d_i - ad_x$; and therefore $\xi = x - at$, $\xi' = x + at$;

$$\therefore D_i \Delta_i u = W + O;$$

$$\therefore u = S_i \Sigma_i W + F(x - at) + f(x + at).$$

Ex. 5. To integrate

$$d_i^2 u = a^2 d_x^2 u + 2abd_x d_y u + b^2 d_y^2 u.$$

The right-hand member is equivalent to $(ad_x + bd_y)^2 u$; hence the equation is equivalent to

$$\begin{aligned} 0 &= d_t^2 u - (ad_x + bd_y)^2 u \\ &= (d_t + ad_x + bd_y)(d_t - ad_x - bd_y)u. \end{aligned}$$

Consequently $D_t = d_t + ad_x + bd_y$, $\Delta_t = d_t - ad_x - bd_y$; whence $\xi = x - at$, $\eta = y - bt$; $\xi' = x + at$, $\eta' = y + bt$; and the proposed equation becomes

$$\begin{aligned} D_t \Delta_t u &= 0; \\ \therefore u &= S_t O + \Sigma_t O \quad (\text{Art. 15}), \\ &= F(\xi, \eta) + f(\xi', \eta') \\ &= F(x - at, y - bt) + f(x + at, y + bt). \end{aligned}$$

Ex. 6. To integrate

$$\begin{aligned} d_t^2 u + (a + b)d_t d_x u + abd_x^2 u + A(d_t u + ad_x u) \\ + B(d_t u + bd_x u) + ABu = W. \end{aligned}$$

In this example $D_t = d_t + ad_x$, $\Delta_t = d_t + bd_x$; hence $\xi = x - at$, $\xi' = x - bt$; and the proposed equation becomes

$$\begin{aligned} D_t \Delta_t u + A D_t u + B \Delta_t u + ABu &= W; \\ \therefore (D_t + B)(\Delta_t + A)u &= W; \\ \therefore \epsilon^{-Bt} D_t \epsilon^{Bt} \cdot \epsilon^{-At} \Delta_t \epsilon^{At} u &= W \dots \dots \dots (1). \end{aligned}$$

The absolute part of the required integral is (see Art. 16)

$$\begin{aligned} u &= \epsilon^{-Bt} S_t \epsilon^{Bt} O + \epsilon^{-At} \Sigma_t \epsilon^{At} O \\ &= \epsilon^{-Bt} F\xi + \epsilon^{-At} f(\xi') \\ &= \epsilon^{-Bt} F(x - at) + \epsilon^{-At} f(x - bt). \end{aligned}$$

The part dependent on W may be determined from equation (1), which gives

$$u = \epsilon^{-At} \Sigma_t \epsilon^{(A-B)t} S_t \epsilon^{Bt} W.$$

Hence the complete integral of the proposed equation is

$$u = e^{-At} \Sigma_i e^{(A-B)t} S_i e^{Bt} W + e^{-Bt} F(x-at) + e^{-At} f(x-bt).$$

17. It may be necessary to shew that $e^{-at} s_i^n e^{at}$ is the symbol of integration corresponding to the symbol of differentiation, $e^{-at} d_i^n e^{at}$; i.e. if $e^{-at} d_i^n e^{at} u = v$, then $u = e^{-at} s_i^n e^{at} v$. This may be proved as follows.

$$\begin{aligned} \text{Since} \quad e^{-at} d_i^n e^{at} u &= v, \\ \therefore d_i^n e^{at} u &= e^{at} v; \\ \therefore e^{at} u &= s_i^n e^{at} v; \\ \therefore u &= e^{-at} s_i^n e^{at} v. \end{aligned}$$

Similar formulæ will of course be true for the symbols D_i and S_i ; for Δ_i and Σ_i ; &c.

Ex. 7. To integrate $d_i^3 u + 3d_i d_x u + 2d_x^2 u = t + x$.

Since $d_i^3 + 3d_i d_x + 2d_x^2 = (d_i + d_x)(d_i + 2d_x)$, we have

$$\begin{aligned} D_i u &= d_i u + d_x u, \quad \Delta_i u = d_i u + 2d_x u; \\ \therefore D_i x &= 1, \quad \Delta_i x = 2, \quad \text{and } x = t + \xi, \quad x = 2t + \xi'; \\ \therefore D_i \Delta_i u &= t + x. \end{aligned}$$

The absolute part of the integral is $S_i \Sigma_i O$, which is equal to $S_i O + \Sigma_i O = F(\xi) + f(\xi') = F(x-t) + f(x-2t)$.

The part of the integral dependent on W or $t+x$ is found as follows: since $D_i(t+x) = 2$, and $\Delta_i(t+x) = 3$,

$$\begin{aligned} u &= \Sigma_i S_i(t+x) = \Sigma_i \cdot \frac{1}{4} (t+x)^3 \\ &= \frac{1}{36} (t+x)^3. \end{aligned}$$

Hence the complete integral, found by adding to this the absolute part, is

$$u = \frac{1}{36} \cdot (t+x)^3 + F(x-t) + f(x-2t).$$

18. The integration of $t+x$ in the preceding example suggests the necessity of the following general explanation. If in any example we know that $D_t x = a$ and $\Delta_t x = b$; and if we have to integrate such an expression as $f(ht+kx)$, we must remember that

$$\text{as } D_t(ht+kx) = h+ka, \text{ and } \Delta_t(ht+kx) = h+kb,$$

$$\therefore S_t^m f(ht+kx) = \frac{f_m(ht+kx)}{(h+ka)^m};$$

$$\text{and } \therefore \Sigma_t^n S_t^m f(ht+kx) = \frac{f_{m+n}(ht+kx)}{(h+ka)^m(h+kb)^n}.$$

Consequently, had the integral of the equation

$$d_t^3 u + (a+b) d_t d_x u + ab d_x^2 u = (ht+kx)^n$$

been required, we should have found

$$u = \frac{(ht+kx)^{n+3}}{(n+1)(n+2)(h+ka)(h+kb)} + F(x-at) + f(x-bt).$$

Ex. 8. To integrate $(d_t - 2d_x)(d_t + d_x)(d_x - 3d_y)u = 0$.

Here we have

$$D_t u = d_t u - 2d_x u, \quad \Delta_t u = d_t u + d_y u, \quad \delta_x u = d_x u - 3d_y u;$$

consequently $D_t x = -2$; $\Delta_t y = 1$, $\delta_x y = -3$;

$$\text{and } x = -2t + \xi, \quad y = t + \xi', \quad y = -3x + \xi'';$$

$$\text{and } D_t \Delta_t \delta_x u = 0;$$

$$\begin{aligned} \therefore u &= S_t O + \Sigma_t O + \sigma_x O \\ &= F(\xi, y) + f(\xi', x) + \phi(\xi'', t) \\ &= F(x+2t, y) + f(y-t, x) + \phi(y+3x, t). \end{aligned}$$

CHAPTER IV.

THE TREATMENT OF QUASI-CONSTANTS.

19. We propose in this Chapter to develop more fully the doctrine of quasi-constants, and we think that it will be seen that by a right use of them a very great extension of integration-power is effected; and that this increase of power is almost wholly the consequence of the use of symbols of operation instead of brackets (Art. 4). From what has been said in Arts. 1 and 2, about the nature of quasi-constants, it will be understood that when we meet with such an equation as $D_i u = d_i u + \xi d_x u + \eta d_y u + \zeta d_z u + \dots$ we may treat it as an equation with constant coefficients. Now, on comparing it with the general formula of Art. 7, we find

$$D_i x = \xi, \quad D_i y = \eta, \quad D_i z = \zeta, \quad \&c. \dots\dots\dots (1),$$

and these by integration give

$$x = \xi t + \phi_1 \xi, \quad y = \eta t + \phi_1 \eta, \quad z = \zeta t + \phi_1 \zeta, \quad \&c. \dots\dots\dots (2).$$

We might give to these integrals an apparently still more general form by writing $\phi_1(\xi, \eta, \zeta \dots)$, $\phi_2(\xi, \eta, \zeta \dots)$,

$$\phi_3(\xi, \eta, \zeta \dots), \dots \text{ for } \phi_1 \xi, \phi_2 \eta, \phi_3 \zeta \dots$$

The forms above given are in general better suited for our purpose; and our conclusion from them is this, that if the

independent variables of a new set are such that each one (as ξ) will satisfy an equation of the typical form $x = \xi t + \phi \xi$, then we know that

$$D_t = d_t + \xi d_x + \eta d_y + \zeta d_z + \dots,$$

and $\therefore$ also $F(D_t) = F(d_t + \xi d_x + \eta d_y + \zeta d_z + \dots) \dots\dots (3).$

20. The simplest case of quasi-constants is found by equating each of the functions $\phi_1, \phi_2, \dots$ to zero, in which case we have

$$\xi = \frac{x}{t}, \quad \eta = \frac{y}{t}, \quad \zeta = \frac{z}{t}, \quad \dots \&c. \dots\dots\dots (4),$$

as the simplest set of quasi-constants that can be found.

21. If in the equation $D_t = d_t + Ld_x + Md_y + Nd_z + \dots$ some of the coefficients $L, M, N, \dots$ are absolute constants, and some of them quasi-constants, we shall have

$$F(D_t) = F(d_t + Ld_x + Md_y + Nd_z + \dots) \dots\dots\dots (5).$$

22. We may here make one general remark which will apply to great numbers of equations. If we have an equation involving differential symbols of operation of one kind only, as d_t , or D_t , or Δ_t , ... then any *absolute* constant in that equation may be replaced by a *quasi*-constant, without in any way affecting the validity of the integral. We shall illustrate this property of quasi-constants by an example or two; and afterwards it will be assumed, even when not mentioned, that the compass of every differential equation and its integral *may* be extended by means of this substitution of quasi-constants for absolute constants.

23. We may also mention, for form's sake, what is obvious enough, that in integrating an equation in which two or more differential symbols of different kinds occur, the quasi-con-

stants corresponding to one symbol will not all (though some of them *may*) be quasi-constants under the other differential symbols.

And, moreover, when the operating symbol in a differential equation is d_t , we may use functions of $(d_x, d_y, d_z \dots)$ as quasi-constants.

Ex. 1. To integrate $td_t u + xd_x u + yd_y u = x^a y^b u^c$.

Dividing by t we see that the coefficients of $d_x u$ and $d_y u$ are quasi-constants (Art. 20), and that $\xi = \frac{x}{t}$, $\eta = \frac{y}{t}$;

$$\therefore D_t u = t^{-1} x^a y^b u^c;$$

$$\therefore u^{-c} D_t u = \xi^a \eta^b t^{a+b-1};$$

$$\begin{aligned} \therefore \frac{u^{1-c}}{1-c} &= \xi^a \eta^b \cdot \frac{t^{a+b}}{a+b} + F(\xi, \eta) \\ &= \frac{x^a y^b}{a+b} + F\left(\frac{x}{t}, \frac{y}{t}\right). \end{aligned}$$

Now this integral has been found on the supposition that a, b, c are absolute constants, but every step of the investigation will be equally valid if we suppose a, b, c to be functions of $\frac{x}{t}, \frac{y}{t}$, (Art. 22).

Ex. 2. To integrate $t^2 d_t^2 u + 2tx d_t d_x u + x^2 d_x^2 u = t^a x^b$.

Dividing by t^2 we find $D_t = d_t + \frac{x}{t} \cdot d_x$, and consequently the coefficients are quasi-constants, and $\xi = \frac{x}{t}$.

$$\therefore D_t^2 u = t^{a-2} x^b = \xi^b \cdot t^{a+b-2};$$

$$\begin{aligned}\therefore u &= \xi^b \cdot \frac{t^{a+b}}{(a+b)(a+b-1)} + t F(\xi) + f(\xi) \\ &= \frac{t^a x^b}{(a+b)(a+b-1)} + t F\left(\frac{x}{t}\right) + f\left(\frac{x}{t}\right).\end{aligned}$$

We are at liberty to write for a and b any functions of $\frac{x}{t}$, (Art. 22).

Ex. 3. To integrate

$$t^2 d_t^2 u + 2tx d_t d_x u + x^2 d_x^2 u + b(ht + kx) \left(d_t u + \frac{x}{t} d_x u \right) + cu = 0.$$

Here as in the last example $D_t = d_t + \frac{x}{t} d_x$, and $\xi = \frac{x}{t}$;

$$\therefore t^2 D_t^2 u + bt(h + k\xi) D_t u + cu = 0,$$

the integral of which is obtained as in ordinary integration for one independent variable;

$$\begin{aligned}\therefore u &= t^m F(\xi) + t^n f(\xi) \\ &= t^m F\left(\frac{x}{t}\right) + t^n f\left(\frac{x}{t}\right);\end{aligned}$$

m and n being the two roots of the equation

$$m(m-1) + \frac{b}{t}(ht + kx)m + c = 0.$$

We are at liberty to write for b and c any functions of $\frac{x}{t}$.

Ex. 4. To integrate

$$d_t^2 u + 2ad_t d_x u + a^2 d_x^2 u + \frac{b}{t}(d_t u + ad_x u) + \frac{cu}{t^2} = 0.$$

In this example $D_t = d_t + ad_x$; $\therefore D_tx = a$, and $x = at + \xi$.

$$\therefore D_t^2 u + \frac{b}{t} \cdot D_t u + \frac{c}{t^2} \cdot u = 0;$$

$$\begin{aligned}\therefore u &= t^m F(\xi) + t^n f(\xi) \\ &= t^m F(x - at) + t^n f(x - at); \end{aligned}$$

m and n being the roots of the equation

$$m(m-1) + bm + c = 0.$$

In this example we are at liberty to write for b and c any functions of $x - at$.

Ex. 5. To integrate $(ht + kx)^2 (d_t^2 u + 2ad_t d_x u + a^2 d_x^2 u)$
 $+ b(ht + kx)(d_t u + ad_x u) + cu = 0.$

Here we assume $D_t = d_t + ad_x$, whence $\xi = x - at$;
 also $D_t(ht + kx) = h + ka$;

$$\therefore (ht + kx)^2 D_t^2 u + b(ht + kx) D_t u + cu = 0.$$

This is an equation of one independent variable, because $D_tx = a$;

$$\begin{aligned}\therefore u &= (ht + kx)^m \cdot F\xi + (ht + kx)^n \cdot f\xi \\ &= (ht + kx)^m \cdot F(x - at) + (ht + kx)^n \cdot f(x - at), \end{aligned}$$

m, n being the values of m in the equation

$$(h + ak)^2 m(m-1) + b(h + ak)m + c = 0.$$

We might add here the same remark as before respecting the constants b, c .

24. The preceding examples will also establish the principle, that if any equation of one independent variable, $\phi(d_t)u = 0$, can be integrated, then the corresponding equa-

tion $\phi(D_i).u=0$ can be integrated, even when the quantities which were constant in $\phi(d_i).u=0$ are replaced in $\phi(D_i).u=0$ by quasi-constants, or in some cases by such quantities as $\psi(ht+kx)$. Thus because the integral of

$$d_i^2u + ad_iu + bu = 0 \text{ is } u = A\epsilon^{mt} + B\epsilon^{nt},$$

m and n being the roots of the equation $m^2 + am + b = 0$; therefore the integral of

$$D_i^2u + aD_iu + bu = 0$$

will be $u = \epsilon^{mt} F'(\xi, \eta, \dots) + \epsilon^{nt} f(\xi, \eta, \dots),$

in which $\xi, \eta, \dots$ are quasi-constants of any form we choose to assign to them.

The equation

$$d_i^2u + 2\frac{x}{t}.d_id_xu + \frac{x^2}{t^2}.d_x^2u + a\left(d_iu + \frac{x}{t}.d_xu\right) + bu = 0,$$

may be taken in illustration of this principle.

Ex. 6. To integrate

$$td_iu + xd_xu + yd_yu + \dots = t^n \phi\left(\frac{x}{t}, \frac{y}{t}, \dots\right).$$

Dividing by t we have

$$\xi = \frac{x}{t}, \eta = \frac{y}{t}, \dots \text{ and } D_iu = t^{n-1} \phi\left(\frac{x}{t}, \frac{y}{t}, \dots\right);$$

$$\therefore D_iu = t^{n-1} \phi(\xi, \eta, \dots);$$

$$\therefore u = \frac{t^n}{n} \cdot \phi(\xi, \eta, \dots) + F(\xi, \eta, \dots)$$

$$= \frac{t^n}{n} \cdot \phi\left(\frac{x}{t}, \frac{y}{t}, \dots\right) + F\left(\frac{x}{t}, \frac{y}{t}, \dots\right).$$

Ex. 7. To integrate

$$d_t u + \xi d_x u + \eta d_y u + \dots = t^{-1} \phi(\xi, \eta, \dots).$$

As in the last example we have here

$$u = \frac{t^n}{n} \cdot \phi(\xi, \eta, \dots) + F(\xi, \eta, \dots).$$

This differs from the preceding example in this, that the values of $\xi, \eta, \dots$ are not given as there, but are to be determined from the equations

$$x = \xi t + f_1(\xi, \eta, \dots), \quad y = \eta t + f_2(\xi, \eta, \dots), \quad \&c.$$

Ex. 8. To integrate $d_t u - a d_x u + \frac{y}{t} d_y u = \frac{y^2}{tx}$.

This is a case where the coefficients are mixed constants.

$D_x x = -a$, and therefore $x = -at + \xi$; but $\frac{y}{t} = \eta$ is a quasi-constant;

$$\therefore D_t u = \frac{y^2}{tx} = \frac{\eta^2 t}{\xi - at};$$

$$\therefore u = -\frac{\eta^2 t}{a} - \frac{\eta^2 \xi}{a^2} \cdot \log(\xi - at) + F(\xi, \eta)$$

$$= -\frac{y^2}{at} - \frac{y^2}{a^2 t^2} (x + at) \log x + F\left(x + at, \frac{y}{t}\right).$$

CHAPTER V.

THE GENERAL LINEAR EQUATION OF THE SECOND ORDER.
COEFFICIENTS CONSTANT.

25. EVERY linear differential equation of one independent variable, and of the second order with constant coefficients, is by custom taken to be integrable in finite terms; that is, the ultimate elementary forms to which that general equation can be reduced are accepted as integrable. Formulæ both algebraic and transcendental have been invented, —logarithmic, exponential and trigonometric, direct and inverse, and have been agreed upon for the purpose that an integral expressed in terms of such forms shall be accepted as a finite integral. Would it be too much to ask that whether there be only one or any number of independent variables, the elementary forms of the general equation of the second order with constant coefficients may be considered integrable forms, and that new transcendentals and appropriate symbols shall be invented for that purpose and take rank with $\log$, $\sin$, $\cos^{-1}$, ...? To agree to this would only be consenting to put all integrations of the second order on the same level, whatever the number of variables.

We shall proceed to shew how on the principles of this Essay the general partial differential equation, which is the

subject of this Chapter, may be reduced to its ultimate elementary forms. We shall take the case of two independent variables only, but we shall treat it in such a manner that the method will be equally applicable to any number of independent variables. We confine our operations to equations of two independent variables, because when there are more, the algebraic expressions become tediously long without illustrating any new principle.

26. The general equation of the second order embraces the two following forms:—

$$\text{I. } d_t^2 u + (a+b)d_t d_x u + ab d_x^2 u + A d_t u + B d_x u + C u = W.$$

$$\text{II. } d_t d_x u + A d_t u + B d_x u + C u = W.$$

We begin with the former.

Case I. When a and b are not equal.

$$\text{Since } d_t^2 + (a+b)d_t d_x + ab d_x^2 = (d_t + a d_x)(d_t + b d_x),$$

$$\text{we assume } D_t = d_t + a d_x, \quad \Delta_t = d_t + b d_x;$$

$$\therefore D_t x = a, \quad x = at + \xi; \text{ and } \Delta_t x = b, \quad x = bt + \xi':$$

and the proposed equation becomes

$$D_t \Delta_t u + \frac{B-Ab}{a-b} \cdot D_t u + \frac{B-Aa}{b-a} \cdot \Delta_t u + C u = W \dots\dots(1);$$

$$\therefore \left(D_t + \frac{B-Aa}{b-a} \right) \left(\Delta_t + \frac{B-Ab}{a-b} \right) u + \left\{ C + \frac{(B-Aa)(B-Ab)}{(a-b)^2} \right\} u = W(2).$$

Now for u write $v e^{m\xi' + n\xi}$, remembering that $D_t \xi = 0$, $\Delta_t \xi = 0$, $D_t \xi' = a - b$, and $\Delta_t \xi' = b - a$.

$$\therefore \left(D_t + \frac{B-Aa}{b-a}\right) u = \epsilon^{m\xi+n\xi} \left\{ D_t + m(a-b) + \frac{B-Aa}{b-a} \right\} \cdot v,$$

$$\text{and } \left(\Delta_t + \frac{B-Ab}{a-b}\right) u = \epsilon^{m\xi+n\xi} \left\{ \Delta_t + n(b-a) + \frac{B-Ab}{a-b} \right\} \cdot v.$$

As m, n are arbitrary constants we are at liberty to assume

$$(a-b)^2 m = B - Aa, \text{ and } (a-b)^2 n = B - Ab;$$

$$\therefore D_t \Delta_t v + \left\{ C + \frac{(B-Aa)(B-Ab)}{(a-b)^2} \right\} v = W \epsilon^{-m\xi-n\xi}.$$

Consequently when $W=0$, the form I. can be reduced to the elementary form,

$$D_t \Delta_t v = cv.$$

Case II. When a and b are equal.

In this case the form I. becomes

$$(d_t + ad_x)^2 u + A \left(d_t + \frac{B}{A} d_x\right) u + Cu = W.$$

Hence $D_t = d_t + ad_x$, $x = at + \xi$ as before; and $\Delta_t = d_t + \frac{B}{A} d_x$

whence $\Delta_t x = \frac{B}{A}$, and $x = \frac{B}{A} t + \xi'$;

$$\therefore D_t^2 u + A \Delta_t u + Cu = W.$$

Now for u write $v \epsilon^{m\xi}$, remembering that $\Delta_t \xi = \frac{B}{A} - a$;

$$\therefore D_t^2 v + A \Delta_t v + \{(B-Aa)m + C\} v = W \epsilon^{-m\xi}.$$

We are at liberty to assume

$$(Aa - B)m = C;$$

$$\therefore D_t^2 v + A \Delta_t v = W \epsilon^{-m\xi}.$$

Consequently when $W = 0$, the form I. when $a = b$ can be reduced to the elementary form,

$$D_t^2 v = c \Delta_t v.$$

There is however an exception to this result, viz. when $B = Aa$, for then the original equation becomes of the following form :

$$D_t^2 u + A D_t u + C u = W,$$

which is always integrable in finite terms (Art. 24).

Case III. When $d_t d_x u + A d_t u + B d_x u + C u = W$;

$$\therefore (d_t + B) (d_x + A) u + (C - AB) u = W.$$

For u write $v e^{mt+nx}$, and assume $m + B = 0$, $n + A = 0$, then

$$u = v^{-Ax-Bt},$$

$$\text{and } d_t d_x v + (C - AB) v = W e^{Ax+Bt};$$

consequently when $W = 0$, the form II. can be reduced to the elementary form,

$$d_t d_x v = cv,$$

which is the same as in the first case.

Thus it appears that there are two elementary forms, and no more, viz.

$$D_t \Delta_t u = cu, \text{ and } D_t^2 u = c \Delta_t u.$$

The question is—are we to accept these as integrable forms? Their symbolical integrals can be easily found, but they cannot be generally integrated in finite terms.

Ex. 1. To find the symbolical integral of $D_t \Delta_t u = cu$.

Since D_t and Δ_t are independent symbols, each of them may be treated as a constant in reference to the other ;

$$\therefore D_t u = \frac{c}{\Delta_t} \cdot u = c \Sigma_t \cdot u;$$

$$\therefore u = e^{c t \Sigma_t} F(\xi).$$

If it should be thought that there is any ambiguity in this expression, we may exhibit it in another form, free from all ambiguity, by considering that

$$\Delta_t F(\xi) = F'(\xi) \cdot \Delta_t \xi = d_\xi F(\xi) \cdot (b-a);$$

$$\therefore (b-a) \Sigma_t = s_\xi;$$

$$\therefore u = e^{\frac{c t s_\xi}{b-a}} F(\xi).$$

In a similar way we should find

$$u = e^{\frac{c t s_{\xi'}}{a-b}} f(\xi').$$

Consequently the complete symbolical integral is

$$u = e^{\frac{c t}{b-a} \cdot s_\xi} F(\xi) + e^{\frac{c t}{a-b} \cdot s_{\xi'}} f(\xi').$$

REMARK. If in these symbolic exponentials we had $d_\xi, d_{\xi'}$, instead of $s_\xi, s_{\xi'}$, the value of u would be expressible in finite terms. Why not extend this character of finiteness to the above form by the invention of an appropriate transcendental?

Ex. 2. To find the symbolical integral of $D_t^2 u = c \Delta_t u$.

Since Δ_t is constant in reference to D_t , we have for the partial integral of this equation,

$$u = (\epsilon^{t\sqrt{c\Delta_t}} + \epsilon^{-t\sqrt{c\Delta_t}}) F(\xi) + (\epsilon^{t\sqrt{c\Delta_t}} - \epsilon^{-t\sqrt{c\Delta_t}}) \cdot \frac{f(\xi)}{\sqrt{c\Delta_t}}.$$

The connexion between these two terms is this:—if the second be represented by u , the first will be represented by $D_t u$ with a change of the arbitrary function. As the pro-

posed equation is linear $D_t^2 u$, $D_t^3 u$, ... each with a change of the arbitrary function will be an integral; but as $D_t^2 u$ will coincide in form with u , and $D_t^3 u$ with $D_t u$, there are only two forms which are different, and the sum of these two given above constitute the whole integral of u as far as the differential equation depends on D_t .

But since the differential equation depends also on Δ , the integral must contain an arbitrary function of ξ . To obtain this we put the equation in the form

$$\Delta u = \frac{D_t^2}{c} \cdot u;$$

$$\therefore u = e^{\frac{t}{c} \cdot D_t^2} \phi(\xi).$$

The complete integral will be obtained by adding this to the above.

27. But now we come upon a difficulty. If we add this to the preceding portion of the integral, we appear to have an integral with *three* instead of *two* arbitrary constants. Will it be sufficient to take the sum of this with *one* of the former two? Do two of the three terms of the whole include the other term? The probability of this may perhaps be due to the circumstance that each term of the integral represents in reality an infinite series, and that in the whole integral each arbitrary function enters with all its derived forms.

Since in the symbolical integral each arbitrary function enters with all its derived forms, the finiteness of the integral can be secured only by assigning to each arbitrary function a periodic form. It is easy to shew that the integral cannot be finite on any other supposition. For if possible let the integral of $D_t^2 u = c \Delta u$ be finite, and let it involve

the arbitrary functions $F(\xi)$ and $f(\xi')$. Then $D_t^2 u$ will involve $f(\xi')$, $f'(\xi')$ and $f''(\xi')$ in addition to $F(\xi)$; it will not contain $F'(\xi)$, $F''(\xi)$, because ξ is constant in reference to D_t .

The other side of the equation $c\Delta_t u$ will involve $F(\xi)$, $F''(\xi)$ in addition to $f(\xi')$. Hence when we substitute for u its value in terms of $F(\xi)$ and $f(\xi')$, we shall have on the left-hand side of the equation $f'(\xi')$, $f''(\xi')$, which are not to be found at all on the right-hand side. On the right hand also there will be $F'(\xi)$, which does not exist on the left hand; and as these functions are arbitrary, and not periodic, the two sides of the equation never can be equivalent to each other. If then a finite integral is required, periodic forms must be found for the arbitrary functions.

28. The following integrals are easily found on this principle. They are given for the sake of reference and illustration.

(1) Given $\xi = x - at$, $\xi' = x - bt$, to find a finite integral of the equation $D_t \Delta_t u = cu$.

$$u = C_1 \epsilon^{\frac{\sqrt{c}}{a-b} \cdot (m\xi + n\xi')} + C_2 \epsilon^{\frac{\sqrt{c}}{b-a} \cdot (m\xi + n\xi')}.$$

C_1 , C_2 , m and n being arbitrary constants, the last two being subject to the condition $mn + 1 = 0$.

(2) Given $\xi = x - at$, and $\xi' = x - bt$, to find a finite integral of the equation $D_t^2 u = c\Delta_t u$.

$$u = (C_1 \epsilon^{\sqrt{c} \cdot \xi} + C_2 \epsilon^{-\sqrt{c} \cdot \xi}) \cdot \epsilon^{n^2(a-b)\xi} + (C_3 \epsilon^{\sqrt{c} \cdot \xi} + C_4 \epsilon^{-\sqrt{c} \cdot \xi}) \cdot \epsilon^{n^2(b-a)\xi}.$$

C_1 , C_2 , C_3 , C_4 , n are all arbitrary constants.

(3) Given $d_t^2 u = a^2 d_x^2 u + b^2 d_y^2 u$, to find an integral in finite terms.

This equation gives

$$(d_t + ad_x)(d_t - ad_x)u = b^2 d_y^2 u.$$

Hence $D_t = d_t + ad_m$, $\Delta_t = d_t - ad_x$, $\xi = x - at$, $\xi' = x + at$;

$$\therefore D_t \Delta_t u = d_y^2 u.$$

The symbolical integral may be at once obtained from this by integrating separately with S_t , Σ_t , s_y ; and it will involve four arbitrary functions,

$$F(\xi, y), f(\xi', y), \phi_1(\xi, \xi'), \phi_2(\xi, \xi').$$

But the following integral in finite terms, though involving only two arbitrary functions, may be useful in certain cases.

$$u = F\left(m\xi + \frac{\xi'}{m} + \frac{2a}{b} \cdot y\right) + f\left(m\xi + \frac{\xi'}{m} - \frac{2a}{b} \cdot y\right),$$

m being an arbitrary constant.

The same may be given in the following form:

$$u = F\left(m\xi - \frac{\xi'}{m} + \frac{2a\sqrt{-1}}{b} \cdot y\right) + f\left(\frac{\xi'}{m} - m\xi + \frac{2a\sqrt{-1}}{b} \cdot y\right),$$

and these two forms may be combined by addition with others derived from these by writing $\frac{1}{m}$ for m .

(4) The equation $d_t^2 u + d_x^2 u = cu$ by a change of the independent variables may be reduced to the form $d_t d_x u = cu$.

The symbolical integral of the latter form is

$$u = e^{cta} F(t) + e^{cxa} f(x).$$

And the following is an integral in finite terms,

$$u = C_1 e^{\left(mt + \frac{x}{m}\right)\sqrt{c}} + C_2 e^{-\left(mt + \frac{x}{m}\right)\sqrt{c}},$$

C_1 , C_2 , m being all arbitrary constants.

If c be negative, the finite integral of $d_t d_x u + cu = 0$ is

$$u = C_1 e^{(mt - \frac{x}{m})\sqrt{c}} + C_2 e^{(\frac{x}{m} - mt)\sqrt{c}}.$$

In this and the preceding case we are at liberty to add other terms to the finite integrals, derived from those given by changing m into $\frac{1}{m}$.

The equation $d_t^2 u - a^2 d_x^2 u = cu$, being treated in the usual manner, gives $D_t = d_t + ad_x$, $\Delta_t = d_t - ad_x$, $\xi = x - at$, $\xi' = x + at$;

$$\therefore D_t \Delta_t u = cu.$$

This result shews that it is vain to look for a finite general integral of any equation of the form $d_t^2 u - a^2 d_x^2 u = cu$.

Ex. 3. To integrate

$$d_t^2 u + (a+b) d_t d_x u + ab d_x^2 u + Ad_t u + Bd_x u + Cu = W,$$

when the coefficients satisfy the equation

$$C(a-b)^2 + (B-Aa)(B-Ab) = 0.$$

On reference to equation (2) Art. 26, Case I., it will be seen that when the above condition is satisfied by the constants, the proposed equation may be reduced to

$$\left(D_t + \frac{B-Aa}{b-a}\right) \left(\Delta_t + \frac{B-Ab}{a-b}\right) u = W.$$

For brevity write

$$(D_t + a)(\Delta_t + \beta)u = W;$$

$$\therefore e^{-at} D_t e^{at} \cdot e^{-\beta t} \Delta_t e^{\beta t} u = W;$$

$$\begin{aligned} \therefore u &= e^{-at} S_t e^{at-\beta t} \sum_t e^{\beta t} W + e^{-at} S_t e^{at} O + e^{-\beta t} \sum_t e^{\beta t} O \\ &= e^{-at} S_t e^{(a-\beta)t} \sum_t e^{\beta t} W + e^{-at} F(\xi) + e^{-\beta t} f(\xi'), \end{aligned}$$

in which $\xi = x - at$ and $\xi' = x - bt$.

Ex. 4. To integrate

$$d_t^2 u + 2a d_t d_x u + a^2 d_x^2 u + A (d_t u + a d_x u) + Cu = W.$$

This is equivalent to

$$D_t^2 u + A D_t u + Cu = W;$$

$$\therefore (D_t + \alpha) (D_t + \beta) u = W;$$

if α, β be the roots of $a^2 - Aa + C = 0$.

$$\therefore u = \epsilon^{-\alpha t} S_t \epsilon^{(\alpha-\beta)t} S_t \epsilon^{\beta t} W + \epsilon^{-\alpha t} F(\xi) + \epsilon^{-\beta t} f(\xi).$$

The double integration in the first term of this integral may be avoided.

In this example A, C may be quasi-constants (Art. 22).

29. The resolution of multiple integration into single integration may be thus effected.

$$\text{Since} \quad (d_t + \alpha) W = \epsilon^{-\alpha t} d_t \epsilon^{\alpha t} W,$$

$$\text{and} \quad (d_t + \beta) W = \epsilon^{-\beta t} d_t \epsilon^{\beta t} W;$$

$$\therefore (\beta - \alpha) W = \epsilon^{-\beta t} d_t \epsilon^{\beta t} W - \epsilon^{-\alpha t} d_t \epsilon^{\alpha t} W.$$

Integrate both sides with $\epsilon^{-\alpha t} s_t \epsilon^{\alpha t} \cdot \epsilon^{-\beta t} s_t \epsilon^{\beta t}$;

$$\therefore (\beta - \alpha) \epsilon^{-\alpha t} s_t \epsilon^{\alpha t} \epsilon^{-\beta t} s_t \epsilon^{\beta t} W = \epsilon^{-\alpha t} s_t \epsilon^{\alpha t} W - \epsilon^{-\beta t} s_t \epsilon^{\beta t} W.$$

This is the formula for the reduction of a double integral to two single ones.

If we integrate both sides with $\epsilon^{-\gamma t} s_t \epsilon^{\gamma t}$ the left hand will become a triple integral, and the right hand two double integrals. Each of the latter may be resolved into single integrals, and the final result will be found to be

$$\epsilon^{-\alpha t} s_t \epsilon^{(\alpha-\beta)t} s_t \epsilon^{(\beta-\gamma)t} s_t \epsilon^{\gamma t} W = \frac{\epsilon^{-\alpha t} s_t \epsilon^{\alpha t} W}{(\gamma-\alpha)(\beta-\alpha)} + \frac{\epsilon^{-\beta t} s_t \epsilon^{\beta t} W}{(\gamma-\beta)(\alpha-\beta)} + \frac{\epsilon^{-\gamma t} s_t \epsilon^{\gamma t} W}{(\beta-\gamma)(\alpha-\gamma)}.$$

The law is obvious for any number of integrations.

If any of the quantities $\alpha, \beta, \gamma, \dots$ be equal, the method will hold good, but the formula just found will fail. The method will furnish the right formula in every case.

Thus if $\alpha = \beta$, $\epsilon^{-\alpha t} s_t \epsilon^{\alpha t} \cdot \epsilon^{-\alpha t} s_t \epsilon^{\alpha t} W = \epsilon^{-\alpha t} s_t^2 \epsilon^{\alpha t} W$, which does not admit of further reduction.

By putting $W = 0$ we obtain the following formula, which agrees with Art. 16,

$$\epsilon^{-\alpha t} s_t \epsilon^{\alpha t} \cdot \epsilon^{-\beta t} s_t \epsilon^{\beta t} \cdot \epsilon^{-\gamma t} s_t \epsilon^{\gamma t} O = \epsilon^{-\alpha t} s_t O + \epsilon^{-\beta t} s_t O + \epsilon^{-\gamma t} s_t O.$$

Ex. 5. To integrate $d_t d_x u + A d_t u + B d_x u + A B u = W$.

This is equivalent to $(d_t + B)(d_x + A)u = W$;

$$\therefore \epsilon^{-Bt} d_t \epsilon^{Bt} \cdot \epsilon^{-Ax} d_x \epsilon^{Ax} u = W;$$

$$\begin{aligned} \therefore u &= \epsilon^{-Bt} s_t \epsilon^{Bt - Ax} s_x \epsilon^{Ax} W + \epsilon^{-Bt} s_t \epsilon^{Bt} O + \epsilon^{-Ax} s_x \epsilon^{Ax} O \\ &= \epsilon^{-Bt} s_t \epsilon^{Bt - Ax} s_x \epsilon^{Ax} W + \epsilon^{-Bt} F(x) + \epsilon^{-Ax} f(t). \end{aligned}$$

Ex. 6. To integrate

$$d_t^2 u + 2a d_t d_x u + a^2 d_x^2 u = \phi''(ht + kx, \xi);$$

where ϕ'' is the second differential coefficient of ϕ with regard to $ht + kx$, ξ being considered constant.

In this case

$$D_t = d_t + a d_x, \quad \xi = x - at, \quad \text{and} \quad D_t(ht + kx) = h + ka;$$

$$\therefore D_t^2 u = \phi''(ht + kx, \xi);$$

$$\therefore u = \frac{\phi(ht + kx, \xi)}{(h + ka)^2} + tF(\xi) + f(\xi).$$

Ex. 7. To integrate

$$d_t^2 u + \frac{2x}{t} \cdot d_t d_x u + \frac{x^2}{t^2} \cdot d_x^2 u = \phi''(ht + kx, \xi).$$

In this case

$$D_t = d_t + \frac{x}{t} \cdot d_x, \quad \xi = \frac{x}{t}, \quad \text{and} \quad D_t (ht + kx) = h + k \cdot \frac{x}{t};$$

$$\therefore D_t^2 u = \phi'' (ht + kx, \xi);$$

$$\therefore u = \frac{t^2}{(ht + kx)^2} \cdot \phi (ht + kx, \xi) + tF\left(\frac{x}{t}\right) + f\left(\frac{x}{t}\right).$$

Ex. 8. To integrate

$$d_t^2 u + 2\xi d_t d_x u + \xi^2 d_x^2 u = \phi'' (ht + kx, \xi).$$

In this case $x = \xi t + \phi_1(\xi) \dots \dots \dots (1).$

$$u = \frac{\phi (ht + kx)}{(h + k\xi)^2} + tF(\xi) + f(\xi).$$

ξ in this Example is any quantity which satisfies equation (1); and the function ϕ_1 is arbitrary.

Several of the examples contained in this Chapter will be more fully treated of in the last Chapter.

CHAPTER VI.

EQUATIONS OF THE SECOND ORDER WITH VARIABLE
COEFFICIENTS.

30. As our object is to exhibit the new method of integration, we shall, in order to avoid complexity, produce formulæ for only two independent variables, but the methods used will always be equally applicable to any number of variables.

Since $D_t u = d_t u + D_x . d_x u$;

$$\therefore \Delta_t D_t u = \Delta_t (d_t u) + \Delta_t (D_x . d_x u) + D_x . \Delta_t (d_x u).$$

By writing $d_t u$, $d_x u$ successively for u in the former equation, and substituting the results in the latter, we have

$$\Delta_t D_t u = d_t^2 u + (D_x + \Delta_x) d_t d_x u + D_x . \Delta_x . d_x^2 u \\ + \Delta_t D_x . d_x u \dots \dots \dots (1).$$

This is a very important formula, and will enable us in solving a certain class of equations with variable coefficients to eliminate at one step all terms of the second order, by substituting $\Delta_t D_t u - \Delta_t D_x . d_x u$ for them.

31. It is important to notice that if the operative symbols $D_t \Delta_t$ be commutative on x , they will be so on u , i.e. on any function of t, x . The commutativeness of the symbols of

operation is of such great utility in the work of integration, that we shall prove this proposition for any number of variables.

Let u be represented by $\phi(x)$ as far as it depends on the independent variable x . Then as far as x is concerned,

$$D_i u = \phi'(x) \cdot D_i x, \text{ and } \Delta_i D_i u = \phi'(x) \cdot \Delta_i D_i x + \phi''(x) \cdot \Delta_i x \cdot D_i x.$$

But by performing the operations in a contrary order we should find

$$D_i \Delta_i u = \phi'(x) \cdot D_i \Delta_i x + \phi''(x) \cdot D_i x \cdot \Delta_i x.$$

Consequently if $\Delta_i D_i x = D_i \Delta_i x$, i.e. if D_i, Δ_i be commutative on x , then as far as u depends on x , they are commutative on u , any function of the variable. The same may be proved for $y, z, \dots$ the remaining variables. Hence the symbols of operation D_i, Δ_i are commutative on u in reference to all the variables on which they are individually commutative.

As a general rule, also, the symbols will be commutative on any variable, as x , if $D_i x$ and $\Delta_i x$ be both constant.

32. When it is desirable to commute the symbols D_i, Δ_i , and they are not commutative *per se*, they may be commuted by the formula,

$$D_i \Delta_i u = \Delta_i D_i u + (D_i \Delta_i x - \Delta_i D_i x) \cdot d_x u.$$

For since

$$\begin{aligned} \Delta_i D_i u &= d_i^2 u + (D_i x + \Delta_i x) \cdot d_i d_x u + D_i x \cdot \Delta_i x \cdot d_x^2 u + \Delta_i D_i x \cdot d_x u; \\ \therefore D_i \Delta_i u &= d_i^2 u + (\Delta_i x + D_i x) \cdot d_i d_x u + \Delta_i x \cdot D_i x \cdot d_x^2 u + D_i \Delta_i x \cdot d_x u \\ &= \Delta_i D_i u + (D_i \Delta_i x - \Delta_i D_i x) \cdot d_x u. \end{aligned}$$

33. The following formulæ are easily proved, P, Q being any functions of the independent variables.

$$\begin{aligned} \epsilon^{-P} \Delta_i \epsilon^P D_i u &= d_i^2 u + (D_i x + \Delta_i x) d_i d_x u + D_i x \cdot \Delta_i x \cdot d_x^2 u \\ &\quad + \Delta_i P \cdot d_i u + (\Delta_i P \cdot D_i x + \Delta_i D_i x) \cdot d_x u \dots \dots \dots (2). \end{aligned}$$

$$\begin{aligned} \epsilon^{-P}\Delta_i\epsilon^{P-Q}D_i\epsilon^Qu &= d_i^2u + (D_ix + \Delta_ix).d_id_xu + D_ix.\Delta_ix.d_x^2u \\ &+ (\Delta_iP + D_iQ).d_iu + (\Delta_iP.D_ix + D_iQ.\Delta_ix + \Delta_iD_ix).d_xu \\ &+ (\Delta_iP.D_iQ + \Delta_iD_iQ)u \dots\dots\dots (3). \end{aligned}$$

34. It will be seen that the formulæ (1), (2), (3) increase step by step in comprehensiveness. Formula (1) can, by giving proper values to D_ix , Δ_ix (which are arbitrary), be made to agree with all the terms of the second order in any proposed equation. Formula (2) can be made to agree with one term more by giving a proper value to P (which is arbitrary). Formula (3) can be made to agree with one term more by assigning to Q a proper value. If therefore we have an equation of the general form

$$d_i^2u + Hd_xd_xu + Kd_x^2u + Ld_iu + Md_xu + Ru = W,$$

it can always be reduced to the form

$$\epsilon^{-P}\Delta_i\epsilon^{P-Q}D_i\epsilon^Qu + R'u = W,$$

and if R' be equal to zero, it can be integrated.

35. The formula (1) shews that we can substitute

$$\Delta_iD_iu - (\Delta_iD_ix).d_xu$$

for the whole of the terms of the second order; and since

$$D_iu = d_iu + D_ix.d_xu, \text{ and } \Delta_iu = d_iu + \Delta_ix.d_xu,$$

these two equations will enable us to eliminate d_iu and d_xu from the remaining part of any proposed equation.

Ex. 1. To integrate

$$d_i^2u + 2ad_id_xu + (a^2 - b^2t^2)d_x^2u - bd_xu = W.$$

Comparing this equation with formula (1) we find

$$D_ix + \Delta_ix = 2a, \text{ and } D_ix.\Delta_ix = a^2 - b^2t^2;$$

$$\therefore D_t x = a - bt, \text{ and } \Delta_t x = a + bt; \text{ and } \therefore \Delta_t D_t x = -b;$$

$$\therefore x = at - \frac{1}{2} bt^2 + \xi, \text{ and } x = at + \frac{1}{2} bt^2 + \xi';$$

$$\therefore \Delta_t D_t u = W;$$

$$\therefore D_t u = \Sigma_t W + f(\xi');$$

$$\therefore u = S_t \Sigma_t W + S_t f(\xi') + F(\xi)$$

$$= S_t \Sigma_t W + S_t f(\xi - bt^2) + F\left(x - at + \frac{1}{2} bt^2\right).$$

REMARK. In this example $\Delta_t D_t x = -b$, and $D_t \Delta_t x = b$; consequently $\Delta_t D_t x$ is not equal to $D_t \Delta_t x$, and D_t , Δ_t are not commutative. As an instance of the important influence of commutateness of the symbols D_t , Δ_t , we may mention that had D_t , Δ_t been commutative the absolute part of the above integral would have been

$$S_t O + \Sigma_t O = F(\xi) + f(\xi') = F\left(x - at + \frac{1}{2} bt^2\right) + f\left(x - at - \frac{1}{2} bt^2\right).$$

Ex. 2. To integrate $\frac{d_t^3 u}{t^3} - \frac{d_x^3 u}{x^3} = \frac{d_t u}{t^3} - \frac{d_x u}{x^3} + W$.

Multiply this equation by t^3 ,

$$\therefore d_t^3 u - \frac{t^3}{x^3} \cdot d_x^3 u - \frac{1}{t} \cdot d_t u + \frac{t^2}{x^3} \cdot d_x u = W t^2.$$

Now by formula (1)

$$\Delta_t D_t u - \Delta_t D_t x \cdot d_x u = d_t^3 u + (D_t x + \Delta_t x) d_t d_x u + D_t x \cdot \Delta_t x \cdot d_x^2 u;$$

and this will agree with the terms of the second order of the proposed equation if we assume $D_t x = -\Delta_t x = \frac{t}{x}$;

$$\therefore x^2 = t^2 + \xi, \quad x^2 = -t^2 + \xi', \quad \text{and} \quad \Delta_t D_x x = \frac{1}{x} + \frac{t^2}{x^3};$$

$$\therefore \Delta_t D_t u - \left(\frac{1}{x} + \frac{t^2}{x^3} \right) \cdot d_x u - \frac{1}{t} \cdot d_t u + \frac{t^2}{x^3} \cdot d_x u = W t^2;$$

$$\therefore \Delta_t D_t u - \frac{1}{t} \left(d_t u + \frac{t}{x} \cdot d_x u \right) = W t^2;$$

$$\therefore \Delta_t D_t u - \frac{1}{t} D_t u = W t^2.$$

Divide by t and integrate with Σ_t ;

$$\therefore D_t u = t \Sigma_t (W t) + t f(\xi);$$

$$\therefore u = S_t \Sigma_t (W t) + S_t f(\xi) + F(\xi).$$

But $S_t f(\xi) = S_t f(\xi + 2t^2) = \phi(\xi + 2t^2) = \phi(x^2 + t^2)$;

$$\therefore u = S_t \Sigma_t W t + F(x^2 - t^2) + \phi(x^2 + t^2).$$

Ex. 3. To integrate $(1 - t^2)^2 d_t^2 u + 2(1 - t^2)(1 - tx) d_t d_x u + (1 - tx)^2 d_x^2 u - 2t(1 - t^2) d_t u - (t + x - 2t^2 x) d_x u + c^2 u = 0$.

Divide by $(1 - t^2)^2$;

$$\begin{aligned} \therefore d_t^2 u + \frac{2(1 - tx)}{1 - t^2} \cdot d_t d_x u + \left(\frac{1 - tx}{1 - t^2} \right)^2 d_x^2 u - \frac{2t}{1 - t^2} \cdot d_t u \\ - \frac{t + x - 2t^2 x}{(1 - t^2)^2} d_x u + \frac{c^2 u}{(1 - t^2)^2} = 0. \end{aligned}$$

Comparing this with formula (1) we see immediately that the symbols D_t and Δ_t must be identical, and that

$$D_x x = \frac{1 - tx}{1 - t^2}, \quad \text{and} \quad \therefore D_t^2 x = \frac{t - x}{(1 - t^2)^2};$$

$$\therefore D_t^2 u - \frac{t - x}{(1 - t^2)^2} \cdot d_x u - \frac{2t}{1 - t^2} \cdot d_t u - \frac{t + x - 2t^2 x}{(1 - t^2)^2} \cdot d_x u + \frac{c^2 u}{(1 - t^2)^2} = 0;$$

$$\therefore D_t^2 u - \frac{2t}{1-t^2} \cdot \left(d_t u + \frac{1-tx}{1-t^2} \cdot d_x u \right) + \frac{c^2 u}{(1-t^2)^2} = 0;$$

$$\therefore D_t^2 u - \frac{2t}{1-t^2} \cdot D_t u + \frac{c^2 u}{(1-t^2)^2} = 0;$$

$$\therefore D_t \{ (1-t^2) D_t u \} + \frac{c^2 u}{1-t^2} = 0;$$

$$\therefore \{ (1-t^2) D_t \}^2 u + c^2 u = 0.$$

Now change the independent to τ , such that $D_\tau t = 1 - t^2$.

$$\therefore D_\tau^2 u + c^2 u = 0;$$

$$\therefore u = F(\xi) \cos c\tau + f(\xi) \sin c\tau.$$

ξ is known from $(1-t^2) D_t x = 1 - tx$, and τ from $D_\tau t = 1 - t^2$.

$$\therefore \xi = \frac{x-t}{\sqrt{1-t^2}}, \text{ and } \tau = \log \sqrt{\frac{1+t}{1-t}}.$$

CHAPTER VII.

INTEGRATION BY FACTORIAL SYMBOLS OF OPERATION.

36. If T be any function of t , and $f(d_i)$ or f any function of the symbol of operation d_i , then the symbol $f(d_i)$ applied to TU is equivalent to the symbol $Tf + T'f' + \frac{1}{1.2} T''f'' + \frac{1}{1.2.3} T'''f''' + \dots$, applied to U .

Since

$$\begin{aligned} d_i^n . TU &= T d_i^n U + \frac{n}{1} d_i T . d_i^{n-1} U + \frac{n(n-1)}{1.2} d_i^2 T . d_i^{n-2} U + \dots \\ &= (T d_i^n + d_i T . n d_i^{n-1} + d_i^2 T . \frac{n(n-1)}{2} d_i^{n-2} + \dots) U, \\ &= (T . \kappa^n + T' . d_i . \kappa^n + \frac{1}{1.2} T'' . d_i^2 . \kappa^n + \dots) . U, \end{aligned}$$

where the Greek letter κ is used to represent d_i as a symbol of *quantity*.

This being true for all integer values of n , it follows for all interpretable forms of $f(\kappa)$, that

$$\begin{aligned}
 f(d_i) \cdot TU &= \{Tf(\kappa) + T'd_\kappa f(\kappa) + \frac{1}{1 \cdot 2} \cdot T''d_\kappa^2 f(\kappa) + \dots\} \cdot U, \\
 &= (Tf + T'f' + \frac{1}{1 \cdot 2} \cdot T''f'' + \frac{1}{1 \cdot 2 \cdot 3} T'''f''' + \dots) \cdot U.
 \end{aligned}$$

37. This theorem may be symbolically represented by the exponential expression,

$$f(d_i) \cdot TU = e^{d_i d_\kappa} T f(\kappa) \cdot U,$$

in the use of which we are to notice that d_i in the exponential acts only on T , and d_κ only on $f(\kappa)$: and before the series of operations is to be applied to U the value of κ is to be restored, i.e. d_i is to be written for κ .

38. If P be a function of several independent variables $t, x, y, \dots$ and $f(d_i, d_x, \dots)$ any function of the operative symbols, then the symbol $f(d_i, d_x, \dots)$ applied to PU is equivalent to the symbol $e^{d_i d_\kappa + d_x d_\lambda + \dots} P f(\kappa, \lambda, \dots)$ applied to U , the Greek letters $\kappa, \lambda, \dots$ representing $d_i, d_x, \dots$ when these are to be treated as symbols of quantity.

$$\begin{aligned}
 \text{For} \quad d_i^n \cdot PU &= e^{d_i d_\kappa} P \kappa^n \cdot U, \\
 \therefore d_x^m d_i^n \cdot PU &= d_x^m e^{d_i d_\kappa} P \kappa^n \cdot U, \\
 &= e^{d_i d_\kappa} d_x^m P \kappa^n \cdot U, \\
 &= e^{d_i d_\kappa} e^{d_x d_\lambda} P \kappa^n \lambda^m \cdot U, \\
 &= e^{d_i d_\kappa + d_x d_\lambda} P \kappa^n \lambda^m \cdot U.
 \end{aligned}$$

Hence for all interpretable forms of $f(d_i, d_x)$ we have

$$f(d_i, d_x) \cdot PU = e^{d_i d_\kappa + d_x d_\lambda} P f(\kappa, \lambda) \cdot U.$$

The same process may be continued to any number of independent variables; and consequently the theorem is true for any number.

39. If P be any function of the independent variables $t, x, y, z, \dots$ and if $f(D_i, \Delta_i, \delta_i, \dots)$ be any interpretable function of the operative symbols when these are commutative among themselves, then the symbol $f(D_i, \Delta_i, \delta_i, \dots)$ applied to PU is equivalent to the symbol $e^{D_i d_\kappa + \Delta_i d_\lambda + \delta_i d_\mu + \dots} P f(\kappa, \lambda, \mu, \dots)$ applied to U ; $\kappa, \lambda, \mu, \dots$ representing $D_i, \Delta_i, \delta_i, \dots$ when these are to be treated as symbols of quantity.

The proof of this theorem is similar in principle to that given in the preceding article, and it is unnecessary to repeat it.

The cases of the theorems of this and the preceding articles of this chapter which most frequently occur are the following

$$(1) \dots f(d_i) \cdot tU = (tf + f') U.$$

$$(2) \dots f(d_i, d_x) \cdot (ht + kx) U = \{(ht + kx)f + (kd_i + hd_x)f'\} \cdot U.$$

$$(3) \dots f(D_i \Delta_i) \cdot tU = \{tf + (D_i + \Delta_i)f'\} \cdot U.$$

40. Examples of the use of these theorems will now be given.

Ex. 1. To integrate

$$d_i^2 u + (a + b) d_i d_x u + a b d_x^2 u + A d_i u + B d_x u + C u = \frac{C'}{t^2} u.$$

By proceeding as in Art. 26, Case I, we reduce this equation to the form

$$D_i \Delta_i v + cv = \frac{C'}{t^2} \cdot v, \dots\dots\dots (1)$$

in which equation $v = u e^{-m\xi + n\xi'}$, $m(a - b)^2 = B - Aa$, $n(a - b)^2 = B - Ab$, and $\xi = x - at$, $\xi' = x - bt$.

We may now transform equation (1) by writing w^k for v , and assuming $k(k-1) = C$.

$$\therefore D_t \Delta_t w + \frac{k}{t} (D_t + \Delta_t) w + cw = 0.$$

$$\therefore t (D_t \Delta_t + c) w + k (D_t + \Delta_t) w = 0 \dots\dots\dots(2).$$

Operate on this with $f(D_t \Delta_t)$ by the formula (3) of last Article.

$$\begin{aligned} \therefore \{t f(D_t \Delta_t) + (D_t + \Delta_t) f'(D_t \Delta_t)\} (D_t \Delta_t + c) w \\ + k (D_t + \Delta_t) f(D_t \Delta_t) w = f(D_t \Delta_t) 0. \end{aligned}$$

Now as the form of the function f is arbitrary we are at liberty to assume the sum of the terms after the first equal to zero.

$$\therefore f'(D_t \Delta_t) (D_t \Delta_t + c) + k f(D_t \Delta_t) = 0.$$

This is a linear equation of the first order,

$$\therefore f(D_t \Delta_t) = (D_t \Delta_t + c)^{-k};$$

$$\therefore t (D_t \Delta_t + c)^{1-k} w = (D_t \Delta_t + c)^{-k} 0 \dots\dots\dots(3);$$

$$\therefore w = (D_t \Delta_t + c)^{k-1} t^{-1} (D_t \Delta_t + c)^{-k} 0.$$

$$\therefore u = t^k \epsilon^{m\xi + n\xi} \cdot (D_t \Delta_t + c)^{k-1} t^{-1} (D_t \Delta_t + c)^{-k} 0.$$

This is the integral of the proposed equation; but it is obvious that it is only symbolical, except when k is an integer and $c=0$, i.e. when

$$C(a-b)^2 + (B-Aa)(B-Ab) = 0.$$

Ex. 2. To integrate the equation given in the last example, when the conditions just stated are satisfied.

$$\therefore u = t^k \epsilon^{m\xi + n\xi} \cdot (D_t \Delta_t)^{k-1} \frac{1}{t} (S_t \Sigma_t)^k 0.$$

Now $S_i^* \Sigma_i^* 0 = S_i^* 0 + \Sigma_i^* 0,$

consequently u consists of two independent parts, one depending on $S_i^* 0$, and the other on $\Sigma_i^* 0$. We may seek them separately and similarly.

$$\begin{aligned} \frac{1}{t} S_i^* 0 &= \frac{1}{t} (A_0 + A_1 t + A_2 t^2 + \dots + A_{k-1} t^{k-1}) \\ &= A_0 t^{-1} + A_1 + A_2 t + \dots + A_{k-1} t^{k-2}, \end{aligned}$$

$A_0, A_1, A_2 \dots$ being functions of ξ .

$$\therefore D_i^{k-1} \cdot \frac{1}{t} S_i^* 0 = F(\xi) \cdot t^k;$$

$$\therefore (D_i \Delta_i)^{k-1} \frac{1}{t} \Sigma_i^* 0 = \Delta_i^{k-1} \cdot t^k F(\xi).$$

Similarly $(D_i \Delta_i)^{k-1} \frac{1}{t} \Sigma_i^* 0 = D_i^{k-1} t^k f(\xi).$

$$\therefore u = \epsilon^{m\xi+n\xi} t^k \{D_i^{k-1} t^{-k} f(\xi) + \Delta_i^{k-1} t^{-k} F(\xi)\}.$$

This is the integral required; but it fails in the case of $k = \text{a negative integer}$.

To find the integral in this case write $-k$ for k ; then since $c = 0$, equations (2), (3) of example (1) become

$$t D_i \Delta_i w = k (D_i + \Delta_i) w,$$

and $t (D_i \Delta_i)^{k+1} w = (D_i \Delta_i)^k 0 = 0;$

$$\therefore w = S_i^{k+1} 0 + \Sigma_i^{k+1} 0.$$

The two parts of this integral are independent. We take them separately.

$$\therefore w = S_i^{k+1} 0 = F_0 \xi + t F_1 \xi + t^2 F_2 \xi + \dots + t^{k-1} F_{k-1} \xi + t^k F_k \xi.$$

Here we seem to have $k+1$ arbitrary functions of ξ , and there ought to be only one. The forms of the others must be determined by substituting this result in the equation $tD_t \Delta_t w = k(D_t + \Delta_t)w$. If this be done the following will be found to be the law of the functions,

$$F_{n+1}(\xi) = \frac{k-n}{2k-n} \cdot \frac{b-a}{n+1} \cdot d_\xi F_n(\xi).$$

In the same way we shall find

$$f_{n+1}(\xi) = \frac{k-n}{2k-n} \cdot \frac{a-b}{n+1} \cdot d_\xi f_n(\xi);$$

$$\therefore u = e^{m\xi + n\xi} t^{-\frac{1}{2}} \cdot \left\{ F_0 \xi + t F_1 \xi + t^2 F_2 \xi + \dots + t^k F_k \xi \right. \\ \left. + f_0 \xi' + t f_1 \xi' + t^2 f_2 \xi' + \dots + t^k f_k \xi' \right\}.$$

Ex. 3. To integrate the equation

$$d_t^2 u + 2a d_t d_x u + a^2 d_x^2 u + A(d_t u + a d_x u) + Bu = \frac{Cu}{t^2}.$$

In this example $D_t = d_t + a d_x$, and $\xi = x - at$;

$$\therefore D_t^2 u + A D_t u + Bu = \frac{Cu}{t^2}.$$

For u write vt^m , and assume $m(m-1) = C$;

$$\therefore t(D_t^2 + A D_t + B)v + m D_t v = 0.$$

It will be convenient to assume

$$D_t^2 + A D_t + B = (D_t + \alpha)(D_t + \beta),$$

$$\text{now } m D_t = \frac{m}{\alpha - \beta} \cdot \{\alpha(D_t + \beta) - \beta(D_t + \alpha)\};$$

$$\therefore t(D_t + \alpha)(D_t + \beta)v + \frac{m\alpha}{\alpha - \beta}(D_t + \beta)v - \frac{m\beta}{\alpha - \beta}(D_t + \alpha)v = 0.$$

Differentiate this equation with $f(D_i)$;

$$\therefore (tfD_i + f'D_i) (D_i + \alpha) (D_i + \beta) v + \frac{m\alpha}{\alpha - \beta} (D_i + \beta) fD_i \cdot v \\ - \frac{m\beta}{\alpha - \beta} (D_i + \alpha) fD_i \cdot v = fD_i \cdot 0.$$

But because the form of f is arbitrary we are at liberty to assume, either

$$(D_i + \beta) fD_i - \frac{m\beta}{\alpha - \beta} \cdot fD_i = 0, \text{ or, } (D_i + \alpha) fD_i + \frac{m\alpha}{\alpha - \beta} \cdot fD_i = 0.$$

The former gives

$$fD_i = (D_i + \beta)^{\frac{m\beta}{\alpha - \beta}} \dots \dots \dots (1)$$

and the latter gives

$$fD_i = (D_i + \alpha)^{\frac{m\alpha}{\beta - \alpha}} \dots \dots \dots (2)$$

If we take the former, and for brevity write n for $\frac{m\beta}{\alpha - \beta}$, we have

$$tfD_i \cdot (D_i + \alpha) (D_i + \beta) v + \frac{m\alpha}{\alpha - \beta} (D_i + \beta) fD_i \cdot v = fD_i \cdot 0,$$

$$\text{or, } t(D_i + \alpha) \cdot (D_i + \beta)^{n+1} v + \frac{m\alpha}{\alpha - \beta} \cdot (D_i + \beta)^{n+1} v = (D_i + \beta)^n \cdot 0.$$

This being an equation of the first order with $(D_i + \beta)^{n+1} v$ as the dependent variable can be integrated as in the former part of this Essay.

REMARKS. It is to be remarked respecting this Example that the final integral will be only symbolical, unless n be an integer positive or negative. But as we might have used equation (2) instead of (1), we should succeed in the integra-

tion with that equation if $\frac{m\alpha}{\beta-\alpha}$ be an integer: and thus the only condition of success is that one of the two quantities $m\alpha, m\beta$ shall be a multiple of $\alpha-\beta$, where α, β are the two roots of the equation $\alpha^2 - A\alpha + B = 0$.

The method of integration will be similar if the right-hand member of the proposed equation be $C'(ht+kx)^{-2}u$ instead of $C't^{-2}u$.

Ex. 4. We come now to an example which has excited a good deal of attention, viz.

$$0 = d_t^2 u + (a+b) d_t d_x u + ab d_x^2 u + \frac{Ad_t u + Bd_x u}{ht+kx} + \frac{Cu}{(ht+kx)^2}.$$

Here $D_t = d_t + ad_x$, $\Delta_t = d_t + bd_x$, $\xi = x - at$, $\xi' = x - bt$, $D_t(ht+kx) = h + ka$, $\Delta_t(ht+kx) = h + kb$.

In the proposed equation write $(ht+kx)^m v$ for u , and equate to zero the coefficient of v .

$$\therefore 0 = (h+ka)(h+kb)m(m-1) + (Ah+Bk)m + C,$$

and

$$0 = (ht+kx)(d_t^2 u + \overline{a+b} \cdot d_t d_x u + ab d_x^2 u) + \{A + m(2h + \overline{a+b} \cdot k)\} d_t v \\ + \{B + m(2abh + \overline{a+b} \cdot h)\} d_x v;$$

Eliminate d_t and d_x

$$\therefore 0 = (ht+kx) D_t \Delta_t v + \left\{ \frac{B-Ab}{a-b} + m(h+bk) \right\} D_t v \\ + \left\{ \frac{B-Aa}{b-a} + m(h+ak) \right\} \Delta_t v.$$

For brevity represent the coefficients of $D_t v$ and $\Delta_t v$ by $\alpha(h+bk)$, and $\beta(h+ak)$;

$$\therefore 0 = (ht + kx) D_t \Delta_i v + \alpha (h + bk) D_t v + \beta (h + ak) \Delta_i v.$$

Operate on this with $f(\Delta_i)$,

$$\begin{aligned} \therefore f(\Delta_i) 0 &= \{(ht + kx) f \Delta_i + (h + bk) f' \Delta_i\} D_t \Delta_i v \\ &\quad + \alpha (h + bk) D_t f \Delta_i v + \beta (h + ak) \Delta_i f \Delta_i v. \end{aligned}$$

Now as the form of f is arbitrary we may assume

$$\Delta_i f' \Delta_i + \alpha f \Delta_i = 0;$$

$$\therefore f \Delta_i = \Delta_i^{-\alpha} = \Sigma_i^{\alpha}.$$

Hence

$$f \Delta_i \cdot 0 = (ht + kx) f \Delta_i \cdot D_t \Delta_i v + \beta (h + ak) \Delta_i f \Delta_i v,$$

$$\text{or} \quad \Sigma_i^{\alpha} 0 = (ht + kx) D_t \cdot \Delta_i^{1-\alpha} v + \beta (h + ak) \Delta_i^{1-\alpha} v.$$

This is an equation of the first order with respect to $\Delta_i^{1-\alpha} v$ as the dependent variable, and may therefore be integrated as in Chap. II. Ex. 9.

REMARK. The final integral will be only symbolical unless α be an integer; but as we might have operated with $f D_t$ instead of $f \Delta_i$, we should succeed in this case if β be an integer; and consequently the proposed equation will be integrable in finite terms if either α or β , i.e. if either

$$m(a-b) + \frac{B-Ab}{h+kb} \quad \text{or} \quad m(b-a) + \frac{B-Aa}{h+ka}$$

be a multiple of $a-b$. And as there are two values of m either of which we are at liberty to use, we shall be able to integrate the proposed equation in finite terms if one of the values of m will render one of the above quantities a multiple of $a-b$.

Ex. 5. To integrate

$$(ht + kx) (d_t^2 u + 2ad_t d_x u + a^2 d_x^2 u) + b(d_t u + ad_x u) + cu = 0.$$

In this example

$$D_t = d_t + ad_x, \quad \xi = x - at, \quad \text{and} \quad D_t(ht + kx) = h + ka;$$

$$\therefore (ht + kx) D_t^2 u + b D_t u + cu = 0.$$

Operate on this equation with $f(D_t)$;

$$\therefore \{(ht + kx) f(D_t) + (h + ka) f'(D_t)\} \cdot D_t^2 u + b D_t \cdot f(D_t) u + cf(D_t) u = f(D_t) \cdot 0.$$

We are at liberty to assume

$$(h + ka) f'(D_t) \cdot D_t^2 + (b D_t + c) f(D_t) = 0;$$

$$\begin{aligned} \therefore f(D_t) &= D_t^{\frac{-b}{h+ka}} \cdot \frac{e D_t^{-1}}{e^{h+ka}} \\ &= S_t^{\frac{b}{h+ka}} \cdot \frac{e S_t}{e^{h+ka}} \dots \dots \dots (1); \end{aligned}$$

$$\therefore (ht + kx) D_t^2 \cdot f(D_t) u = f(D_t) \cdot 0.$$

The occurrence of the exponential operative symbol in equation (1) shews that the proposed equation cannot be generally integrated in finite terms. Success would also require that b should be a multiple of $h + ka$.

Ex. 6. To integrate

$$d_t d_x u + A d_t u + B d_x u + Cu = \frac{C'}{t^2} \cdot u.$$

Reduce the equation as in Art. 26, Case III., by assuming $u = v e^{-Ax - Bt}$;

$$\therefore d_t d_x v + (C - AB) v = \frac{C'}{t^2} \cdot v \dots \dots \dots (1);$$

$$\therefore \frac{t^2 d_t}{(C - AB) t^2 - C'} \cdot d_x v + v = 0.$$

Now assume a new variable τ such that

$$t^2 d_t \tau = (C - AB) t^2 - C';$$

$$\therefore \tau = (C - AB) t + \frac{C'}{t},$$

$$\text{and } d_\tau d_x v + v = 0.$$

A form which we have seen cannot be integrated generally in finite terms. The proposed equation is therefore of a form which in general can only be integrated symbolically.

Ex. 7. To integrate

$$d_t d_x u + \frac{A d_t u + B d_x u}{ht + kx} + \frac{Cu}{(ht + kx)^2} = 0.$$

Write $(ht + kx)^m v$ for u , and equate to zero the coefficient of v ;

$$\therefore 0 = hkm(m-1) + (Ah + Bk)m + C,$$

$$\text{and } 0 = (ht + kx) d_t d_x v + (A + mk) d_t v + (B + mh) d_x v.$$

Operate on this equation with $f(d_t)$;

$$\begin{aligned} \therefore \{ (ht + kx) f d_t + h f' d_t \} \cdot d_t d_x v + (A + mk) d_t f(d_t) v \\ + (B + mh) d_x f(d_t) v = f(d_t) \cdot 0. \end{aligned}$$

Now as the form of f is arbitrary we are at liberty to assume

$$h d_t f'(d_t) + (B + mh) f(d_t) = 0;$$

$$\therefore f(d_t) = (d_t)^{-m - \frac{B}{h}} = s_t^{m + \frac{B}{h}},$$

$$\text{and } (ht + kx) d_t f(d_t) d_x v + (A + mk) d_t f(d_t) v = f(d_t) \cdot 0,$$

$$\therefore (ht + kx) d_x \cdot d_t f(d_t) v + (A + mk) d_t f(d_t) v = f(d_t) \cdot 0.$$

This equation is of the first order with respect to $d_t f(d_t) v$ as the dependent variable, and it can be integrated by the general method given in the former part of this Essay. It is however essential to the finiteness of this integral that $\frac{B}{h} + m$ be an integer: but as we might have operated with $f(d_x)$ instead of $f(d_t)$, in which case the condition of finiteness would have depended on $\frac{A}{h} + m$ being an integer, it is evident that the proposed equation admits of integration in finite terms if either

$$\frac{A}{h} + m, \text{ or } \frac{B}{h} + m$$

be an integer. And as there are two values of m , either of which we are at liberty to use, the integral will be finite if either of the values of m will satisfy this condition.

Ex. 8. To integrate

$$d_t d_x u + A d_t u + B d_x u + A B u = \frac{C u}{(h t + k x)^2}.$$

For u write $v e^{-Bt - Ax}$ as in Art. 24, Case III.;

$$\therefore d_t d_x v = \frac{C v}{(h t + k x)^2}.$$

Now for v write $w (h t + k x)^m$, and assume

$$m(m-1)hk = C;$$

$$\therefore (h t + k x) d_t d_x w + m k d_t w + m h d_x w = 0.$$

Differentiate this equation with $f(d_t d_x)$;

$$\begin{aligned} \therefore \{ (h t + k x) f(d_t d_x) + (h d_x + k d_t) f'(d_t d_x) \} d_t d_x w \\ + m (k d_t + h d_x) f(d_t d_x) w = f(d_t d_x) 0. \end{aligned}$$

We are at liberty to assume

$$f'(d_i d_x) d_i d_x + m f(d_i d_x) = 0;$$

$$\therefore f(d_i d_x) = (d_i d_x)^{-m} = s_i^m s_x^m,$$

and

$$(ht + kx) f(d_i d_x) d_i d_x w = f(d_i d_x) 0;$$

$$\therefore (ht + kx) s_i^{m-1} s_x^{m-1} w = s_i^m s_x^m 0 = s_i^m 0 + s_x^m 0.$$

$$\text{Now } s_i^m 0 = A_0 + A_1 t + A_2 t^2 + \dots + A_{m-1} t^{m-1},$$

where $A_0, A_1, A_2, \dots$ are arbitrary functions of x ; but it will be more convenient to assume

$$s_i^m 0 = A_0 + A_1 (ht + kx) + A_2 (ht + kx)^2 + \dots \\ \dots + A_{m-1} (ht + kx)^{m-1},$$

which we are at liberty to do because x is here a quasi-constant;

$$\therefore s_i^{m-1} s_x^{m-1} w = A_0 (ht + kx)^{-1} + A_1 + \dots + A_{m-1} (ht + kx)^{m-2};$$

$$\therefore w = d_x^{m-1} (ht + kx)^{-m} F(x).$$

Similarly we have

$$w = d_i^{m-1} (ht + kx)^{-m} f(t).$$

Hence the complete integral is

$$w = d_i^{m-1} (ht + kx)^{-m} f(t) + d_x^{m-1} (ht + kx)^{-m} F(x).$$

If m be a negative integer, write $-m$ for it; then

$$f(d_i d_x) = d_i^m d_x^m;$$

$$\therefore (ht + kx) d_i^{m+1} d_x^{m+1} w = d_i^m d_x^m 0 = 0;$$

$$\therefore w = s_i^{m+1} s_x^{m+1} 0 = s_i^{m+1} 0 + s_x^{m+1} 0$$

$$= F_0 x + (ht + kx) F_1 x + \dots + (ht + kx)^{m-1} F_{m-1} x$$

$$+ f_0 t + (ht + kx) f_1 t + \dots + (ht + kx)^{m-1} f_{m-1} t.$$

The laws of the arbitrary functions must be determined as in Art. 40, Ex. 2, by substituting this result in the equation from which they were derived, viz.

$$(ht + kx) d_x d_x w = m k d_x w + m h d_x w.$$

If this be done we find

$$F_{n+1} x = \frac{n-m}{(2m-n)(n+1)} \cdot \frac{1}{k} \cdot F_n' x,$$

$$f_{n+1} t = \frac{n-m}{(2m-n)(n+1)} \cdot \frac{1}{h} \cdot f_n' t.$$

Ex. 9. The equation

$$t \phi(d_i) u + A \phi'(d_i) u = 0$$

can be deprived of its last term, and integrated if A be an integer.

Differentiate it with $f(d_i)$;

$$\therefore (t f d_i + f' d_i) \phi(d_i) u + A f(d_i) \phi'(d_i) u = f(d_i) 0.$$

We are at liberty to assume

$$\phi(d_i) f' d_i + A \phi'(d_i) f d_i = 0;$$

$$\therefore f d_i = (\phi d_i)^{-A},$$

and
$$t (\phi d_i)^{1-A} u = (\phi d_i)^{-A} 0;$$

$$\therefore u = (\phi d_i)^{A-1} t^{-1} (\phi d_i)^{-A} 0.$$

Ex. 10. The equation

$$(ht + kx) \phi(D_i) u + A (h + ka) \phi'(D_i) u = 0,$$

where $D_x a = a$, if treated in the same manner, will give

$$u = (\phi D_i)^{A-1} (ht + kx)^{-1} \phi(D_i)^{-A} 0.$$

41. If we have a linear differential equation, the coefficients of which are of the form $A_n t^n + A_{n-1} t^{n-1} + \dots$ we may arrange the terms of the equation according to the powers of t . It will then take the form

$$\phi_0 (d_t) u + t \phi_1 (d_t) u + t^2 \phi_2 (d_t) u + \dots = 0.$$

We shall shew how every equation of the kind here proposed may be solved, at least symbolically. And this we shall do by shewing how the equation can be reduced to another of one dimension lower in t : and as this process may be repeated on the reduced equation, we shall at length arrive at an equation of the form

$$t \psi_1 (d_t) u = \psi_0 0,$$

from which u will be known. We shall not attempt to prove this in a general way, in order to avoid complexity, but a few examples will be given which will render the general proof unnecessary.

Ex. 11. To reduce to

$$t \phi_1 (d_t) u + \phi_0 (d_t) u = 0.$$

Operate on this ~~equation~~ ^{equation} with $f(d_t)$;

$$\therefore (t f d_t + f' d_t) \phi_1 (d_t) u + f d_t \cdot \phi_0 (d_t) u = f d_t \cdot 0.$$

As the form of f is arbitrary we are at liberty to assume

$$f' (d_t) \phi_1 (d_t) + f (d_t) \phi_0 (d_t) = 0 \dots \dots \dots (1);$$

$$\therefore t f (d_t) \phi_1 (d_t) u = f (d_t) 0;$$

$$\therefore u = (f d_t \cdot \phi_0 d_t)^{-1} t^{-1} f (d_t) 0.$$

Equation (1) will give the form of $f(d_t)$, and that of $\phi_1(d_t)$ is given in the proposed equation: and consequently u is symbolically known from the last equation.

Ex. 12. To reduce

$$t^2 \phi_2 (d_t) u + t \phi_1 (d_t) u + \phi_0 (d_t) u = 0,$$

one degree lower in the powers of t .

Differentiate with $f(d_t)$;

$$\therefore (t^2 f d_t + 2t f' d_t + f'' d_t) \phi_2 (d_t) u + (t f d_t + f' d_t) \phi_1 (d_t) u + f(d_t) \cdot \phi_0 (d_t) u = f(d_t) 0.$$

Equate to zero the terms on the left-hand side which do not contain t , and divide the remaining portion by t ;

$$\therefore \phi_2 (d_t) \cdot f''(d_t) + \phi_1 (d_t) \cdot f'(d_t) + \phi_0 (d_t) \cdot f(d_t) = 0,$$

and

$$t f(d_t) \phi_2 (d_t) u + \{2f'(d_t) \cdot \phi_2 (d_t) + f(d_t) \phi_1 (d_t)\} u = t^{-1} f(d_t) 0.$$

The former of these being of the second order will give two forms of $f(d_t)$, either of which may be used; and the latter equation falls under the preceding example.

These two examples are sufficient to exhibit the principle of the method, and to shew that any equation arranged according to powers of t can be integrated, symbolically at least, if the successive equations for finding the operative factorial $f(d_t)$ can be solved.

It is hardly necessary to say that we may write $ht + kx$ for t , and D_t for d_t ; and a similar proposition will hold good respecting an equation of the form

$$\phi_0 (D_t) u + (ht + kx) \phi_1 (D_t) u + (ht + kx)^2 \phi_2 (D_t) u + \dots = 0,$$

if $D_t x = \text{constant}$, or quasi-constant.

Ex. 13. To integrate

$$t (d_t + \alpha) (d_t + \beta) u = A (d_t + \alpha) u + B (d_t + \beta) u.$$

Differentiate with $f(d_i)$,

$$\begin{aligned}\therefore (t f d_i + f' d_i) (d_i + \alpha) (d_i + \beta) u &= A f(d_i) (d_i + \alpha) u \\ &+ B f(d_i) (d_i + \beta) u + f(d_i) 0.\end{aligned}$$

We are at liberty to assume $f'(d_i) (d_i + \alpha) (d_i + \beta) u$ equal to either of the terms of the right-hand member which contain u ;

$$\therefore \frac{f' d_i}{f d_i} = \frac{A}{d_i + \beta}, \text{ or } = \frac{B}{d_i + \alpha};$$

$$\therefore f(d_i) = (d_i + \beta)^A, \text{ or } (d_i + \alpha)^B.$$

If now either A or B be an integer, we may take the corresponding form for $f(d_i)$. Suppose A an integer;

$$\therefore f(d_i) = (d_i + \beta)^A,$$

$$\text{and } t(d_i + \alpha) (d_i + \beta)^{A+1} u = B (d_i + \beta)^{A+1} u + (d_i + \beta)^A 0.$$

If we write v for $(d_i + \beta)^{A+1} u$, we have

$$t(d_i + \alpha) v = B v + (d_i + \beta)^A 0,$$

which being an equation of the first order can be integrated as in the early part of this Essay. The proposed equation can therefore be integrated in finite terms if either A or B be an integer.

CHAPTER VIII.

 θ OR $\log t$ AS LEADING INDEPENDENT VARIABLE.

42. THERE is an important class of equations which seem to be most easily integrated by the use of θ ($= \log t$) as leading independent variable. A few of the requisite formulæ are added.

$$\text{Since} \quad t = e^\theta; \therefore d_\theta t = t \dots\dots\dots (1).$$

$$\text{Again, since} \quad d_\theta u = d_i u \cdot d_\theta t = t d_i u; \\ \therefore d_\theta = t d_i \dots\dots\dots (2).$$

$$\text{Again;} \quad d_\theta \cdot t^{-n} u = t^{-n} (d_\theta - n) u; \\ \therefore t^n d_\theta t^{-n} = d_\theta - n \dots\dots\dots (3); \\ \therefore t^{n+1} d_i t^{-n} = d_\theta - n \dots\dots\dots (4) \text{ from } (2).$$

In (4) write 0, 1, 2, 3, ... $(n-1)$ for n ; multiply the results together, and reverse them;

$$\therefore d_\theta (d_\theta - 1) (d_\theta - 2) \dots (d_\theta - n + 1) \\ = (d_\theta - n + 1) (d_\theta - n + 2) \dots (d_\theta - 1) d_\theta \\ = t^n d_i t^{-n} \cdot t^{n-1} d_i t^{-n} \dots t^2 d_i t^{-1} \cdot t d_i \\ = t^n \cdot d_i^n \dots\dots\dots (5).$$

43. The following formulæ of operations are useful abbreviations.

Since $(d_\theta + k) = t^k d_\theta t^k;$

$$\therefore (d_\theta + k)^n = t^k d_\theta t^k \cdot t^k d_\theta t^k \cdot t^k d_\theta t^k \dots \text{to } n \text{ terms,}$$

$$= t^k d_\theta^n t^k \dots \dots \dots (1);$$

$$\therefore f(d_\theta + k) = t^k f(d_\theta) t^k \dots \dots \dots (2);$$

$$\therefore f(d_\theta + k) \cdot f(d_\theta + 2k) \dots f(d_\theta + nk)$$

$$= t^k f(d_\theta) t^k \cdot t^{2k} f(d_\theta) t^{2k} \dots t^{nk} f(d_\theta) t^{nk}$$

$$= (t^k f d_\theta)^n \cdot t^{nk} \dots \dots \dots (3).$$

Since $f(d_\theta + h + k) = t^h f(d_\theta + k) t^h,$
and $f(d_\theta + h + 2k) = t^h f(d_\theta + 2k) t^h,$
..... = ...

$$\therefore f(d_\theta + h + k) \cdot f(d_\theta + h + 2k) \dots f(d_\theta + h + nk)$$

$$= t^h f(d_\theta + k) f(d_\theta + 2k) \dots f(d_\theta + nk) t^h$$

$$= t^h (t^k f d_\theta)^n \cdot t^{nk+h} \dots \dots \dots (4).$$

It is very important to notice that the formulæ numbered (1), (2), (3), (4) in this, and (1), (2), (3), (4), (5) in the preceding article are commutative with d_θ and with each other, because their equals in terms of d_θ are so.

44. By formula (2) of the last article

$$f(d_\theta + h) = t^h f(d_\theta) t^h;$$

$$\therefore (d_\theta + h) (d_\theta + h - 1) (d_\theta + h - 2) \dots (d_\theta + h - n + 1)$$

$$= t^h d_\theta \cdot (d_\theta - 1) \cdot (d_\theta - 2) \dots (d_\theta - n + 1) t^h$$

$$= t^h \cdot t^n d_\theta^n \cdot t^h \dots \dots \dots (\text{Art. 42, 5}).$$

By this formula we are able to express the product of a series of factorials of the form $(d_\theta + \alpha) (d_\theta + \beta) (d_\theta + \gamma) \dots (d_\theta + \mu)$ in terms of t and d_θ , on the one condition that $\alpha, \beta,$

$\gamma \dots \mu$ can be so arranged as to form an Arithmetical progression whose common difference is unity.

And it is in consequence of this ability to pass from d_θ to d_t with such facility when such a series can be formed out of $\alpha, \beta, \gamma \dots \mu$, that in the following articles our chief object will be to change the values of $\alpha, \beta, \gamma \dots \mu$ or some of them, if necessary, so as to produce such a series when possible.

45. THEOREM. If we have an equation of the form $f(d_\theta) \cdot \phi(d_\theta) u + Ct^k u = 0$, then by writing vt^{-a} for u , the subjects of f and ϕ will be both reduced by α .

$$\text{For } f(d_\theta) \phi(d_\theta) t^{-a} v + Ct^k \cdot t^{-a} v = 0;$$

$$\therefore t^a f(d_\theta) \phi(d_\theta) t^{-a} \cdot v + Ct^k v = 0;$$

$$\therefore f(d_\theta - \alpha) \phi(d_\theta - \alpha) v + Ct^k v = 0 \text{ (Art. 43, 2).}$$

Hence the following Rule:—To pass from the proposed equation to this, or from this to the proposed equation, nothing more is necessary than to use the equation

$$u = vt^{-a}, \text{ or } v = ut^a.$$

46. By the last Article we are able to reduce the subjects of the operative factorials when they are to be affected alike; but if we wish to affect one of them only we must proceed differently.

Given the equation $f(d_\theta) \phi(d_\theta) u + Ct^k u = 0$, to increase or diminish the subject of f , without altering that of ϕ .

Commutate f and ϕ , and differentiate the equation with $t^{-h} (t^k f d_\theta)^n t^{nk+h}$ (see Art. 43, 4); and assume $nk + h = -k$;

$$\therefore \phi(d_\theta) t^{-h} (t^k f d_\theta)^n t^{-k} f(d_\theta) u + Ct^{-h} (t^k f d_\theta)^n u = 0.$$

Now for $(t^k f d_\theta)^n u$ write $(t^{-k})^n v$.

$$\therefore \phi(d_\theta) t^{nk} f(d_\theta) t^{-nk} v + Ct^k v = 0;$$

$$\therefore \phi(d_\theta) f(d_\theta - nk) v + Ct^k v = 0 \dots \text{Art. 43, 2.}$$

THEOREM. Hence if we desire to pass from the equation,

$$f(d_\theta) \phi(d_\theta) u + Ct^k u = 0, \text{ to } f(d_\theta - nk) \phi(d_\theta) v + Ct^k v = 0,$$

or, from the latter to the former, we must employ the equation

$$(t^{-k} f d_\theta)^n u = (t^{-k})^n v.$$

47. Though we have thus succeeded in diminishing or increasing the subject of one operative function without affecting the other, it is to be noted that we can only increase or diminish a subject by integer multiples of k .

48. If we have three factorials $f(d_\theta)$, $F(d_\theta)$, $\phi(d_\theta)$, we can change the subject of f by nk ; and then that of F by $n'k$; and then that of ϕ by $n''k$; and so on for any number of operative factorials; and to accomplish this we have merely to employ in succession the equations

$$(t^{-k} f d_\theta)^n u = (t^{-k})^n v,$$

$$(t^{-k} f d_\theta)^n v = (t^{-k})^{n'} w, \quad (\text{for } f, v \text{ and } F)$$

$$\&c. = \&c.$$

Thus the result of the preceding articles is, that if we have an equation of the form

$$f(d_\theta) \cdot F(d_\theta) \cdot \phi(d_\theta) w + Ct^k w = 0,$$

we can affect the subjects of f , F , ϕ in two ways:—1st, we can alter them all alike by any proposed quantity; and 2nd, we can alter any one or more of them by multiples of k .

49. In practice a somewhat more convenient form of the preceding equation is the following.

70 θ OR $\log t$ AS LEADING INDEPENDENT VARIABLE.

THEOREM. We can pass from one of the two following equations, viz.

$f(d_\theta + \alpha) \phi(d_\theta) u + Ct^k u = 0$, and $f(d_\theta + \alpha - nk) \phi(d_\theta) v + Ct^k v = 0$,
to the other by means of the equation,

$$(t^{-k} f d_\theta)^n \cdot ut^\alpha = (t^{-k})^n \cdot vt^\alpha.$$

For since $f(d_\theta + \alpha) = t^{-\alpha} f(d_\theta) t^\alpha \dots (\text{Art. 43, 2})$;

$$\therefore f(d_\theta) t^\alpha \phi(d_\theta) u + Ct^k \cdot ut^\alpha = 0;$$

$$\therefore f(d_\theta) t^\alpha \phi(d_\theta) t^{-\alpha} \cdot ut^\alpha + Ct^k \cdot ut^\alpha = 0;$$

$$\therefore f(d_\theta) \phi(d_\theta - \alpha) \cdot ut^\alpha + Ct^k \cdot ut^\alpha = 0.$$

Let us now assume (in order to reduce the subject of f by nk) in this equation,

$$(t^{-k} f d_\theta)^n \cdot ut^\alpha = (t^{-k})^n \cdot vt^\alpha \quad (\text{Article 46}).$$

$$\therefore f(d_\theta - nk) \cdot \phi(d_\theta - \alpha) vt^\alpha + Ct^k \cdot vt^\alpha = 0.$$

Divide by t^α ;

$$\therefore t^{-\alpha} f(d_\theta - nk) \phi(d_\theta - \alpha) t^\alpha v + Ct^k v = 0;$$

$$\therefore f(d_\theta + \alpha - nk) \phi(d_\theta) \cdot v + Ct^k v = 0. \quad (\text{Art. 43, 2.})$$

50. From this it is evident that we may pass from one of the following equations to the other,

$$f(d_\theta + \alpha) F(d_\theta + \beta) \phi(d_\theta) u + Ct^k u = 0,$$

$$\text{and } f(d_\theta + \alpha - nk) F(d_\theta + \beta - n'k) \phi(d_\theta) w + Ct^k w = 0,$$

by assuming, in succession, the equations

$$(t^{-k} f d_\theta)^n \cdot ut^\alpha = (t^{-k})^n \cdot vt^\alpha,$$

$$(t^{-k} f d_\theta)^{n'} \cdot vt^\beta = (t^{-k})^{n'} \cdot wt^\beta. \quad (\text{for } f \text{ and } F).$$

Ex. 1. To reduce $(d_\theta + \alpha)(d_\theta + 5)u + Ct^k u = 0$ to a simpler form in d_t .

Reduce the first factor by nk ,

$$\therefore (d_\theta + \alpha - nk)(d_\theta + 5)v + Ct^k v = 0 \dots \dots \dots (1),$$

and $(t^{-k}d_\theta)^s \cdot ut^s = (t^{-k})^s \cdot vt^s \dots \dots \dots (2).$

Now we have to form, if possible, of the two operative factors of equation (1) two adjacent terms of an Arithmetic Series whose common difference is unity. (Art. 44.)

$$\therefore \alpha - nk - 5 = \pm 1.$$

We have therefore to consider the two cases, $\alpha = nk + 6$ and $\alpha = nk + 4$.

Case 1. When $\alpha = nk + 6$, equation (1) becomes

$$(d_\theta + 6)(d_\theta + 5)v + Ct^k v = 0;$$

$$\therefore d_\theta(d_\theta - 1) \cdot vt^s + Ct^k \cdot vt^s = 0 \text{ (Art. 43, 2);}$$

$$\therefore d_t^2 \cdot vt^s + Ct^{k-2} \cdot vt^s = 0 \dots \dots \dots (3).$$

Case 2. When $\alpha = nk + 4$, equation (1) becomes

$$(d_\theta + 4)(d_\theta + 5)v + Ct^k v = 0,$$

$$\therefore (d_\theta - 1)d_\theta \cdot vt^s + Ct^k \cdot vt^s = 0, \text{ (Art. 43, 2),}$$

$$\therefore d_t^2 \cdot vt^s + Ct^{k-2} \cdot vt^s = 0 \dots \dots \dots (4).$$

The reduced equations (3) and (4) are of the same form, and if that form can be integrated then the value of u may be found from (2).

In the *first* case, if V be the integral of the equation

$$d_t^2 V + Ct^{k-2} V = 0,$$

72 θ OR $\log t$ AS LEADING INDEPENDENT VARIABLE

we find from equation (2)

$$u = t^{-nk-1} (s_i t^{k-1})^n V.$$

In the *second* case,

$$u = t^{-nk-1} (s_i t^{k-1})^n V t.$$

Ex. 2. To integrate the equation

$$d_i^2 u + \frac{A}{t} \cdot d_i u + C t^{k-2} u = 0.$$

Multiply by t^2 and introduce d_θ ;

$$\therefore (d_\theta + A - 1) d_\theta u + C t^k u = 0.$$

Diminish the first factor by nk (Art. 49);

$$\therefore (t^{-k} d_\theta)^n \cdot u t^{A-1} = (t^{-k})^n \cdot v t^{A-1} \dots\dots\dots(1)$$

and

$$(d_\theta + A - nk - 1) d_\theta v + C t^k v = 0.$$

Hence we must have $A - nk - 1 = \pm 1$; there are therefore two cases to consider.

1st, when $A = nk$;

$$(d_\theta - 1) d_\theta v + C t^k v = 0;$$

$$\therefore d_i^2 v + C t^{k-2} v = 0 \text{ (Art. 42, 5) } \dots\dots\dots(2),$$

2nd, when $A = nk + 2$;

$$\therefore (d_\theta + 1) d_\theta v + C t^k v = 0;$$

$$\therefore d_\theta (d_\theta - 1) \cdot v t + C t^k \cdot v t = 0 \text{ (Art. 45);}$$

$$\therefore d_i^2 \cdot v t + C t^{k-2} \cdot v t = 0 \dots\dots\dots(3).$$

The value of u will be known from equation (1), if the equation

$$d_i^2 V + C t^{k-2} V = 0$$

can be integrated.

In the *first* case

$$u = t^{1-nk}(s_t t^{k-1})^n \cdot \frac{V}{t}.$$

In the *second* case

$$u = t^{1-nk}(s_t t^{k-1})^n \cdot Vt.$$

Ex. 3. To find for what values of k the equation $d_t^2 u + Ct^{k-2}u = 0$ is integral in finite terms.

Change the independent variable from t to τ , where $\tau = t^n$.

$$\therefore \tau d_\tau^2 u + \frac{n-1}{n} \cdot d_\tau u + \frac{C}{n^2} \cdot t^{k-n} u = 0.$$

As n is at our disposal we may assume $t^{k-n} = \tau$, which gives $2n = k$.

$$\therefore \tau \left(d_\tau^2 + \frac{C}{n^2} \right) u + \frac{k-2}{2k} (2d_\tau) u = 0.$$

Now on referring to Article 40, Example 9, it will be seen that this exactly corresponds with the equation there integrated, $\phi(d_\tau)$ there representing $d_\tau^2 + \frac{C}{n^2}$ here. The only condition of integrability in finite terms is that A in that example, which represents $\frac{k-2}{2k}$ in this example, shall be an integer;

$$\therefore \frac{k-2}{2k} = \pm \lambda, \text{ an integer;}$$

$$\therefore k = \frac{2}{1 \mp 2\lambda}.$$

Hence if k be a fraction, the numerator of which is 2, and the denominator of which is any positive or negative odd

integer, the proposed equation will be integrable; and its integral will be

$$u = \left(d_r^2 + \frac{C}{n^2}\right)^{\pm \lambda - 1} t^{-1} \left(d_r^2 + \frac{C}{n^2}\right)^{\mp \lambda} \cdot 0.$$

REMARK. C may be a function of d_x .

EX. 4. To take away the second term of the equation $d_i^m u + \frac{A}{t} \cdot d_i^{m-1} u + Ct^k u = 0$.

Multiply by t^m and introduce d_θ ;

$$\therefore t^{m-1} (d_\theta + A) d_i^{m-1} u + Ct^k u = 0;$$

$$\therefore t^{m-1} (d_\theta + A) t^{1-m} \cdot t^{m-1} d_i^{m-1} u + Ct^k u = 0;$$

$$\therefore (d_\theta + A - m + 1) \phi(d_\theta) u + Ct^k u = 0 \text{ (Art. 42, 3)...(1)}$$

where $\phi(d_\theta) = t^{m-1} d_i^{m-1} = d_\theta(d_\theta - 1) \dots (d_\theta - m + 2)$. (Art. 42, 5).

Diminish the first factor of (1) by nk ;

$$\therefore (d_\theta + A - nk - m + 1) \phi(d_\theta) v + Ct^k v = 0 \dots\dots(2),$$

and $(t^{-k} d_\theta)^n \cdot u t^{A-nk+1} = (t^{-k})^n \cdot v t^{A-m+1} \dots\dots\dots(3).$

On looking at the factorial form of $\phi(d_\theta)$ it will be seen at once (Art. 44) that we may only place the first factorial of (2) either before $\phi(d_\theta)$, or after it. There are therefore two cases for consideration.

1st. Let $A = nk$; then $d_\theta + A - nk - m + 1$ comes after $\phi(d_\theta)$.

$$\therefore \phi(d_\theta) (d_\theta - m + 1) v + Ct^k v = 0;$$

$$\therefore d_i^m v + Ct^k v = 0 \dots\dots\dots(4).$$

2nd. Let $A = nk + m$; then $d_\theta + A - nk - m + 1$ comes before $\phi(d_\theta)$.

$$\therefore (d_{\theta} + 1) \phi(d_{\theta}) v + Ct^k v = 0;$$

$$\therefore d_{\theta} \cdot \phi(d_{\theta} - 1) \cdot vt + Ct^k \cdot vt = 0 \text{ (Art. 45);}$$

$$\therefore d_i^m \cdot vt + Ct^{k-m} \cdot vt = 0 \dots \dots \dots (5).$$

These equations (4) and (5) are of the same form as the proposed equation deprived of its second term; and if this form can be integrated, v will be known, and then u from (3).

Ex. 5. To deprive

$$d_i^m u + \frac{A+B+1}{t} \cdot d_i^{m-1} u + \frac{AB}{t^2} d_i^{m-2} u + Ct^{k-m} u = 0$$

of its second and third terms.

Multiply by t^m and introduce d_{θ} ;

$$\therefore (d_{\theta} + A - m + 2) (d_{\theta} + B - m + 2) \phi(d_{\theta}) u + Ct^k u = 0 \dots (1),$$

where $\phi(d_{\theta}) = d_{\theta} (d_{\theta} - 1) \dots (d_{\theta} - m + 3)$.

Diminish the first factor of (1) by nk , and the second by $n'k$ (Art. 50),

$$\therefore (d_{\theta} + A - nk - m + 2) (d_{\theta} + B - n'k - m + 2) \phi(d_{\theta}) w + Ct^k w = 0 \dots \dots \dots (2),$$

and
$$\left. \begin{aligned} (t^{-k} d_{\theta})^n \cdot ut^{A-m+2} &= (t^{-k})^n \cdot vt^{A-m+2} \\ (t^{-k} d_{\theta})^{n'} \cdot vt^{B-m+2} &= (t^{-k})^{n'} \cdot wt^{B-m+2} \end{aligned} \right\} \dots \dots \dots (3).$$

Now there are only three ways in which we can dispose, in accordance with Art. 44, the first two factorials of (2): viz. we can place them (1st) *after* $\phi(d_{\theta})$; (2nd) one *before* and the other *after* $\phi(d_{\theta})$; (3rd) both *before* $\phi(d_{\theta})$. There are therefore these three cases to be taken into consideration.

(1). Let $A = nk$, $B = n'k - 1$. In this case both are after $\phi(d_{\theta})$.

$$\therefore A + B + 1 = (n + n') k, \text{ and } AB = nk (n'k - 1),$$

and $\phi(d_\theta) (d_\theta - m + 2) (d_\theta - m + 1) w + Ct^k w = 0$;

$$\therefore d_i^m . w + Ct^{k-m} . w = 0 \dots \dots \dots (4).$$

(2). Let $A = nk$, $B = n'k + m - 1$. In this case one is before and one after $\phi(d_\theta)$.

$$\therefore A + B + 1 = (n + n')k + m, \text{ and } AB = nk(n'k + m - 1);$$

and $(d_\theta + 1) \phi(d_\theta) (d_\theta - m + 2) w + Ct^k w = 0$;

$$\therefore d_\theta \phi(d_\theta - 1) (d_\theta - m + 1) . wt + Ct^k . wt = 0;$$

$$\therefore d_i^m . wt + Ct^{k-m} . wt = 0 \dots \dots \dots (5).$$

(3). Let $A = nk + m$, $B = n'k + m - 1$. In this case both are before $\phi(d_\theta)$.

$$\therefore A + B + 1 = (n + n')k + 2m, \text{ and } AB = (nk + m)(n'k + m - 1);$$

and $(d_\theta + 2) (d_\theta + 1) \phi(d_\theta) w + Ct^k w = 0$;

$$\therefore d_\theta (d_\theta - 1) \phi(d_\theta - 2) . wt^2 + Ct^k . wt^2 = 0;$$

$$\therefore d_i^m . wt^2 + Ct^{k-m} . wt^2 = 0 \dots \dots \dots (6).$$

The equations (4), (5), (6) are of the same form as the proposed equation deprived of its second and third terms. If this form $(d_i^s V + Ct^{k-m} V = 0)$ can be integrated, the value of u will be given by equations (3).

Ex. 6. To deprive the equation

$$d_i^m u + \frac{2A + 1}{t} . d_i^{m-1} u + \frac{A^2}{t^2} . d_i^{m-2} u + Ct^{k-m} u = 0$$

of its second and third terms.

This is a particular case of the last example, and treated in the same way it gives

$$(d_\theta + A - m + 2)^2 \phi(d_\theta) u + Ct^k u = 0 \dots \dots \dots (1).$$

The factors before $\phi(d_\theta)$ being *equal*, we cannot place them *together* so as to constitute two adjacent terms of an arithmetic series whose common difference is unity. The only course open to us is, to diminish one by nk , and the other by $n'k$, and then place one *before* $\phi(d_\theta)$ and the other *after* $\phi(d_\theta)$, as in the second case of last example.

$$\therefore A = nk + m - 1 = n'k \dots\dots\dots(2),$$

and $(d_\theta + 1) \phi(d_\theta) (d_\theta - m + 2) w + Ct^k w = 0;$

$$\therefore d_i^m . wt + Ct^{k-m} . wt = 0 \dots\dots\dots(3).$$

From (2) we learn that A and $m - 1$ must be multiples of k .

Ex. 7. To deprive

$$d_i^m u - \frac{A(A+1)}{t^2} . d_i^{m-2} u + Ct^{k-m} u = 0$$

of its second term.

This, like the last example, is a particular case of Ex. 5, and treated in the same way it gives (since $B = -A - 1$);

$$(d_\theta + A - m + 2) (d_\theta - A - m + 1) \phi(d_\theta) u + Ct^k u = 0.$$

Corresponding to the three cases of Ex. 5, we have

(1). Let $A = nk$, which gives $n' = -n$.

$$\therefore A(A+1) = nk(nk+1);$$

and $d_i^m w + Ct^{k-m} w = 0 \dots\dots\dots(1).$

It is a condition in this case that A shall be a multiple of k .

(2). Let $A = nk$, which gives $(n + n')k + m = 0$;

$$\therefore A(A + 1) = nk(nk + 1),$$

and $d_i^m \cdot wt + Ct^{k-m} \cdot wt = 0 \dots\dots\dots(2).$

It is a condition in this case that m and A shall be multiples of k .

(3). Let $A = nk + m$, which gives $(n + n')k + 2m = 0$;

$$\therefore A(A + 1) = (nk + m)(nk + m + 1);$$

and $d_i^m \cdot wt^2 = Ct^{k-m} \cdot wt^2 \dots\dots\dots(3).$

It is a condition in this case that $A - m$, and $2m$ shall be multiples of k .

In these three cases if the proposed equation when deprived of its middle term can be integrated, then the integral of the equation with its middle term can be deduced from that integral by means of the equations (3) of Example (5).

51. The method of reduction employed in the preceding examples being general enables us to generalise the results. Thus if we have an equation of the form

$$d_i^m u + \frac{A'}{t} \cdot d_i^{m-1} u + \frac{B'}{t^2} \cdot d_i^{m-2} u + \dots + \frac{L'}{t^\lambda} \cdot d_i^{m-\lambda} u + Gt^{k-m} u = 0,$$

we must multiply this by t^m , introduce d_θ , and cast the first part of the resulting equation into the form of operative factorials. It will then have the form

$$(d_\theta + A - m + \lambda)(d_\theta + B - m + \lambda) \dots (d_\theta + L - m + \lambda) \phi(d_\theta) u + Gt^k u = 0,$$

where $\phi(d_\theta) = d_\theta(d_\theta - 1) \dots (d_\theta - m + \lambda + 1).$

All the symbolic factors preceding $\phi(d_\theta)$ in this equation are to be increased or diminished by multiples of k ; and they are then to be arranged in every possible way with $\phi(d_\theta)$ in which they can form with $\phi(d_\theta)$ a series of m factorials in arithmetic progression whose common difference is unity. There are, speaking generally, and subject to conditions as in the preceding examples, $\lambda + 1$ ways of doing this: and in any of the cases if μ be the number of factorials which stand *before* $\phi(d_\theta)$, then $\lambda - \mu$ will be the number that come *after* $\phi(d_\theta)$, and the reduced equation will be

$$d_t^m \cdot Vt^\mu + Gt^{k-m} \cdot Vt^\mu = 0.$$

If this equation can be integrated, then the integral of the proposed equation can be at once deduced from it.

52. The principal object of this Chapter has been the reduction of equations to integrable forms by depriving them of some of their terms. The symbol d_t has been used throughout and $d_x, d_y, d_z \dots$ have nowhere appeared, so that it would seem as though this Chapter has no connexion with the subject of partial differential equations. It is therefore necessary to point out that the articles of this Chapter are all true when D_t is used for d_t , and S_t for s_t . And not only so, but that even when d_t is used, the coefficient which we have denoted by C may be a function of $d_x, d_y, d_z \dots$, without affecting the truth of the reductions of the proposed equations. Also when D_t is substituted for d_t , then C may be a quasi-constant.

Also since $D_t(ht + kx) = h + ka$ when $D_tx = a$, the propositions admit of our writing $ht + kx$ instead of t when D_t is used. The methods of reduction will be just the same, and the results similar in form. And the quantity a may be a quasi-constant.

80 θ OR $\log t$ AS LEADING INDEPENDENT VARIABLE.

$$\begin{aligned} \text{Thus } (ht + kx)^n D_\theta (ht + kx)^{-n} &= D_\theta - n(h + ka) \\ &= t^{n(h+ka)} D_\theta t^{-n(h+ka)}. \end{aligned}$$

In this case the only effect of using $ht + kx$ for t is to substitute $n(h + ka)$ for n . It will be found that only constants and not forms will be affected by the change of writing $ht + kx$ for t .

CHAPTER IX.

MISCELLANEOUS EXAMPLES.

53. THE design of this Chapter is the illustration of various modes of applying the principles and methods of former Chapters.

Ex. 1. To integrate

$$tD_i\Delta_i u + AD_i u = 0,$$

on the supposition that D_i and Δ_i are commutative.

Commutate the symbols D_i , Δ_i , and multiply by t^{i-1} ;

$$\therefore t^i \Delta_i \cdot D_i u + A t^{i-1} \cdot D_i u = 0.$$

Integrate with Σ_i ;

$$\therefore t^i D_i u = \Sigma_i 0.$$

Divide by t^i and integrate with S_i ;

$$\therefore u = S_i t^{-i} \Sigma_i 0 + S_i 0.$$

Ex. 2. To integrate

$$tD_i\Delta_i u + AD_i u + B\Delta_i u = 0,$$

D_i and Δ_i being commutative.

We know that this can be integrated if we can deprive it of one of its last two terms.

Operate on the equation with $f(\Delta_i)$;

$$\therefore (tf\Delta_i + f'\Delta_i) D_i\Delta_i u + AD_i f(\Delta_i) u + B\Delta_i f(\Delta_i) u = f(\Delta_i) \cdot 0.$$

Since f is arbitrary we are at liberty to assume

$$f'(\Delta_i) D_i\Delta_i + AD_i f(\Delta_i) = 0;$$

$$\therefore \Delta_i f'(\Delta_i) + Af(\Delta_i) = 0;$$

$$\therefore f\Delta_i = \Delta_i^{-A} = \Sigma_i^A;$$

$$\text{and } \therefore tD_i\Delta_i \cdot f(\Delta_i) u + B\Delta_i \cdot f(\Delta_i) u = f(\Delta_i) \cdot 0.$$

$$\therefore tD_i \cdot \Sigma_i^{A-1} u + B \cdot \Sigma_i^{A-1} u = \Sigma_i^A \cdot 0.$$

Multiply by t^{B-1} and integrate with S_i ;

$$\therefore t^B \Sigma_i^{A-1} u = S_i t^{B-1} \Sigma_i^A \cdot 0 + S_i \cdot 0;$$

$$\therefore u = \Delta_i^{A-1} t^{-B} S_i t^{B-1} \Sigma_i^A \cdot 0 + \Delta_i^{A-1} t^{-B} S_i \cdot 0.$$

The form of this integral indicates that unless A be an integer the value of u will be merely symbolical. But no such restriction attaches to B .

If however A be not an integer, and B be integral, we must operate on the proposed equation with $f(D_i)$ instead of $f(\Delta_i)$. The result may be obtained from the above by interchanging A and B , D_i and Δ_i , S_i and Σ_i .

Ex. 3. To integrate

$$(ht + kx) D_i \Delta_i u + AD_i u = 0,$$

on the supposition that $D_i x = a$, $\Delta_i x = b$.

The symbols D_i , Δ_i are commutative (Art. 31), and

$$\xi = x - at, \quad \xi' = x - bt.$$

Commutate the symbols and multiply by $(ht + kx)^{m-1}$;

$$\therefore (ht + kx)^m \Delta_t . D_t u + A (ht + kx)^{m-1} . D_t u = 0.$$

Integrate with Σ_t ;

$$\therefore (ht + kx)^m D_t u = F(\xi'), \text{ if } A = m(h + kb).$$

Divide by $(ht + kx)^m$ and integrate with S_t ;

$$\therefore u = S_t (ht + kx)^{-m} F(\xi') + f(\xi).$$

Ex. 4. To integrate

$$d_t^2 u - c^2 d_x^2 u = \frac{2}{t} . d_t u,$$

In this example $D_t = d_t + c d_x$, $\Delta_t = d_t - c d_x$, $\xi = x - ct$, $\xi' = x + ct$, and D_t , Δ_t are commutative. Also $2d_t = D_t + \Delta_t$.

$$\therefore t D_t \Delta_t u = (D_t + \Delta_t) u \dots \dots \dots (1).$$

Operate on this equation with $D_t \Delta_t$ (Art. 39, 3);

$$\therefore t D_t^2 \Delta_t^2 u + (D_t + \Delta_t) D_t \Delta_t u = (D_t + \Delta_t) D_t \Delta_t u;$$

$$\therefore D_t^2 \Delta_t^2 u = 0;$$

$$\therefore u = S_t^2 0 + \Sigma_t^2 0$$

$$= t F_1(\xi) + F(\xi) + t f_1(\xi') + f(\xi').$$

The forms of the supernumerary arbitrary functions must be determined by substituting this value of u in equation (1); which being done we find

$$0 = F_1(\xi) - c F''(\xi) + f_1(\xi') + c f'(\xi') :$$

and as ξ and ξ' are independent this resolves itself into the two equations

$$F_1(\xi) = c F''(\xi), \text{ and } f_1(\xi') = -c f'(\xi') ;$$

$$\therefore u = ct (F'' \xi - f'' \xi') + F \xi + f \xi'.$$

Ex. 5. To integrate

$$d_t^2 u - c^2 d_x^2 u = \frac{2u}{t^3}.$$

In this case $D_t = d_t + cd_x$, $\Delta_t = d_t - cd_x$, as in the previous example;

$$\therefore D_t \Delta_t u = \frac{2u}{t^3}.$$

Now for u write vt^m , and assume $m(m-1) = 2$, which gives $m = -1$ or 2 .

$$\therefore t D_t \Delta_t v + m(D_t + \Delta_t)v = 0.$$

If we take the value -1 for m , this equation will coincide with the equation (1) of the last example;

$$\therefore ut = v = ct(F'\xi - f'\xi') + F\xi + f\xi';$$

$$\therefore u = c(F'\xi - f'\xi') + t^{-1}(F\xi + f\xi').$$

Ex. 6. To integrate

$$d_t^2 u - c^2 d_x^2 u = \frac{Au}{(ht + kx)^2}.$$

In this example D_t , Δ_t retain the same values as in the last example. Write $v(ht + kx)^m$ for u , and assume

$$m(m-1)(h^2 - k^2 c^2) = A.$$

$$\therefore (ht + kx) D_t \Delta_t v + m(h - kc) D_t v + m(h + kc) \Delta_t v = 0.$$

Operate on this with $f(D_t \Delta_t)$; (for brevity we may write f for this function)

$$\begin{aligned} \therefore \{(ht + kx)f + (h - kc) D_t \cdot f' + (h + kc) \Delta_t \cdot f'\} D_t \Delta_t v \\ + m(h - kc) D_t f \cdot v + m(h + kc) \Delta_t f \cdot v = f(D_t \Delta_t) 0. \end{aligned}$$

Now as f is arbitrary we are at liberty to assume

$$D_i \cdot f'(D_i \Delta_i) \cdot D_i \Delta_i + m D_i f(D_i \Delta_i) = 0;$$

$$\therefore D_i \Delta_i \cdot f'(D_i \Delta_i) + m f(D_i \Delta_i) = 0;$$

$$\therefore f(D_i \Delta_i) = (D_i \Delta_i)^{-m} = (S_i \Sigma_i)^m.$$

$$\therefore (ht + kx) (S_i \Sigma_i)^{m-1} v = (S_i \Sigma_i)^m 0 = S_i^m 0 + \Sigma_i^m 0;$$

$$\therefore v = D_i^{m-1} \Delta_i^{m-1} (ht + kx)^{-1} (S_i^m 0 + \Sigma_i^m 0).$$

This formula indicates that m must be an integer; and that the value of v consists of two independent parts, one depending on $S_i^m 0$, and the other on $\Sigma_i^m 0$. We may find them separately and similarly.

Now

$$\begin{aligned} S_i^m 0 &= F_0(\xi) + F_1(\xi) \cdot (ht + kx) + \dots + F_{m-1}(\xi) \cdot (ht + kx)^{m-1}, \\ (ht + kx)^{-1} S_i^m 0 &= (ht + kx)^{-1} F_0(\xi) + F_1(\xi) + \dots \\ &\quad \dots + (ht + kx)^{m-2} F_{m-1}(\xi); \\ \therefore D_i^{m-1} (ht + kx)^{-1} S_i^m 0 &= (ht + kx)^{-m} F(\xi). \end{aligned}$$

Similarly

$$\begin{aligned} \Delta_i^{m-1} (ht + kx)^{-1} \Sigma_i^m 0 &= (ht + kx)^{-m} f(\xi'); \\ \therefore v &= \Delta_i^{m-1} (ht + kx)^{-m} F(\xi) + D_i^{m-1} (ht + kx)^{-m} f(\xi'); \end{aligned}$$

u will be known from the equation

$$u = v (ht + kx)^m.$$

Ex. 7. To integrate

$$d_i^2 u - c^2 d_u^2 u = \frac{2A}{t} \cdot d_i u.$$

This differs from Example (4) in the last term only, having $2A$ for 2 ;

$$\therefore t D_i \Delta_i u = A (D_i + \Delta_i) u \dots \dots \dots (1).$$

Operate on this equation with $f(D_i \Delta_i)$,

$$\therefore \{tf(D_i \Delta_i) + (D_i + \Delta_i)f'(D_i \Delta_i)\} D_i \Delta_i u \\ = A(D_i + \Delta_i)f(D_i \Delta_i)u + f(D_i \Delta_i)0.$$

We are at liberty to assume

$$D_i \Delta_i f'(D_i \Delta_i) = A f(D_i \Delta_i);$$

$$\therefore f(D_i \Delta_i) = (D_i \Delta_i)^A,$$

and

$$t(D_i \Delta_i)^{A+1}u = (D_i \Delta_i)^A 0 = 0;$$

$$\therefore u = (S_i \Delta_i)^{A+1} 0 = S_i^{A+1} 0 + \Sigma_i^{A+1} 0.$$

We may find the two parts of u separately and similarly.

Now

$$S_i^{A+1} 0 = F_1(\xi) + tF_2(\xi) + t^2F_3(\xi) + \dots + t^AF_{A+1}(\xi).$$

The forms of the supernumerary arbitrary functions must be determined by substituting this expression for u in equation (1). We thus find that

$$F_{n+2}(\xi) = \frac{2c}{n+1} \cdot \frac{A-n}{2A-n} \cdot F'_{n+1}(\xi).$$

In a similar way we find

$$f_{n+2}(\xi) = \frac{-2c}{n+1} \cdot \frac{A-n}{2A-n} \cdot f'_{n+1}(\xi);$$

$$\therefore u = (F\xi + f\xi') + ct(d_\xi F\xi - d_\xi f\xi')$$

$$+ \frac{A-1}{2A-1} \cdot \frac{(2ct)^2}{1 \cdot 2} \cdot (d_\xi^2 F\xi + d_\xi^2 f\xi')$$

$$+ \frac{A-1}{2A-1} \cdot \frac{A-2}{2A-2} \cdot \frac{(2ct)^3}{1 \cdot 2 \cdot 3} \cdot (d_\xi^3 F\xi - d_\xi^3 f\xi') + \dots$$

REMARK. If A be negative, write $-A$ for A , and the integral will then be

$$u = \Delta_i^{-1} \cdot t^{-A} F(\xi) + D_i^{-1} \cdot t^{-A} f(\xi).$$

Ex. 8. To integrate

$$d_i^2 u - d_x^2 u + \frac{4d_i u}{t+x} = 0.$$

Here $D_i = d_i + d_x$, $\Delta_i = d_i - d_x$, $\xi = x - t$, $\xi' = x + t$, $2d_i = D_i + \Delta_i$, and D_i , Δ_i are commutative. The equation becomes

$$(t+x) D_i \Delta_i u + 2\Delta_i u + 2D_i u = 0.$$

Now since $t+x = \xi'$, therefore $D_i \xi' = 2$; and we may integrate at once with S_i ;

$$\therefore \xi' \Delta_i u + 2u = F(\xi).$$

As ξ' is in this equation a quasi-constant, multiply by $\frac{2}{\epsilon^{\xi'}}$ and integrate with Σ_i ;

$$\therefore u \epsilon^{\frac{2}{\xi'}} = \Sigma_i \epsilon^{\frac{2}{\xi'}} F(\xi) + f(\xi').$$

Ex. 9. To integrate

$$d_i (d_i^2 u - d_x^2 u) + d_x (d_i^2 u - d_x^2 u) = \frac{2}{t+x} \cdot (d_i^2 u - d_x^2 u).$$

Here D_i and Δ_i have the same values as in the last Example.

$$\therefore (d_i + d_x) D_i \Delta_i u - \frac{2}{t+x} \cdot D_i \Delta_i u = 0;$$

$$\therefore D_i \cdot D_i \Delta_i u - \frac{D_i (t+x)}{t+x} \cdot D_i \Delta_i u = 0.$$

Divide by $t+x$, and integrate with S_t ,

$$\therefore D_t \Delta_t u = (t+x) F'(\xi) = \xi F'(\xi).$$

Now as ξ' is a quasi-constant in reference to Σ_t , we have

$$\begin{aligned} u &= S_t \xi' \cdot \Sigma_t F'(\xi) + S_t 0 + \Sigma_t 0 \\ &= \frac{1}{4} \xi'^2 \cdot F(\xi) + \phi(\xi) + f(\xi'), \\ &= \frac{1}{4} (t+x)^2 F(t-x) + \phi(t-x) + f(t+x). \end{aligned}$$

REMARK. $\Sigma_t F'(\xi)$ is not exactly equal to $F(\xi)$, but to $-\frac{1}{2} F(\xi)$, but the constant $\left(-\frac{1}{2}\right)$ may be considered as absorbed into the arbitrary function.

The form of the above integral may be simplified by writing $(t-x)^2 + 4tx$ instead of $(t+x)^2$ in the first term. It may then be written in the form

$$u = tx F(t-x) + \phi(t-x) + f(t+x).$$

Ex. 10. To integrate

$$d_t^2 u + (a+b) d_t d_x u + a b d_x^2 u = \frac{Cu}{(ht+kx)^2}.$$

Here

$$D_t = d_t + a d_x, \quad \Delta_t = d_t + b d_x, \quad \xi = x - at, \quad \xi' = x - bt;$$

$$\therefore (ht+kx)^2 D_t \Delta_t u = Cu.$$

If we write $(ht+kx)^m v$ for u we obtain, assuming

$$m(m-1)(h+ka)(h+kb) = C,$$

$$(ht+kx) D_t \Delta_t v + m(h+kb) D_t v + m(h+ka) \Delta_t v = 0.$$

This equation has already been treated in Example (6).

Ex. 11. To integrate

$$d_t^2 u + 2ad_x d_x u + a^2 d_x^2 u = \frac{Cu}{(ht + kx)^2}.$$

Here $D_t = d_t + ad_x$, $\xi = x - at$.

$$\therefore (ht + kx)^2 \cdot D_t^2 u = Cu.$$

The integral of which is

$$u = (ht + kx)^m F(\xi) + (ht + kx)^n f(\xi),$$

m and n being the roots of

$$m(m-1)(h+ka)^2 = C.$$

C may be a quasi-constant.

Ex. 12. To integrate $d_t^2 u - \frac{c^2}{t^4} \cdot d_x^2 u = 0$.

This falls under the general form considered in Art. 50,

Ex. 3. We here assume $t^{-1} = \tau$, and change the independent from t to τ .

$$\therefore \tau(d_\tau^2 u - c^2 d_x^2 u) + 2d_\tau u = 0.$$

[The rule for determining in equations of this form what power of t is to be assumed for τ is this. The power of t in the question is t^{-4} , take the square root of this ($=t^{-2}$) and integrate this ($=t^{-1}$). This is the value of τ . (If the form of t in the question had been t^{-2} , its square root would be t^{-1} , and $\therefore \tau = \log t$.)]

$$\therefore D_\tau = d_\tau + cd_x, \quad \Delta_\tau = d_\tau - cd_x, \quad \xi = x - c\tau, \quad \xi' = x + c\tau;$$

$$\therefore \tau D_\tau \Delta_\tau u + (D_\tau + \Delta_\tau) u = 0.$$

Operate with $(D_\tau \Delta_\tau)^{-1}$,

$$\therefore \tau D_\tau \Delta_\tau (D_\tau \Delta_\tau)^{-1} u = (D_\tau \Delta_\tau)^{-1} 0 = S_i \Sigma_i 0;$$

$$\therefore \tau u = S_i 0 + \Sigma_i 0;$$

$$\begin{aligned}\therefore \frac{u}{t} &= F(\xi) + f(\xi') \\ &= F(x - c\tau) + f(x + c\tau); \\ \therefore u &= tF\left(x - \frac{c}{t}\right) + tf\left(x + \frac{c}{t}\right).\end{aligned}$$

Ex. 13. To integrate $d_i^2 u - c^2 t^{-\frac{1}{2}} d_x^2 u = 0$.

In this case we must assume $t^{\frac{1}{2}} = \tau$.

$$\begin{aligned}\therefore \tau (d_\tau^2 u - c^2 d_x^2 u) &= 2d_\tau u; \\ \therefore \tau D_\tau \Delta_\tau u &= (D_\tau + \Delta_\tau) u \dots \dots \dots (1).\end{aligned}$$

Operate on this equation with $(D_\tau \Delta_\tau)$,

$$\begin{aligned}\therefore \tau (D_\tau \Delta_\tau)^2 u &= 0; \\ \therefore u &= S_\tau^2 0 + \Sigma_\tau^2 0 \\ &= F(\xi) + \tau F_1(\xi) + f(\xi') + \tau f_1(\xi').\end{aligned}$$

Substituting $F(\xi) + \tau F_1(\xi)$ in equation (1) for u we find $F_1(\xi) = cF'(\xi)$. Similarly $f_1(\xi') = -cf'(\xi')$.

$$\begin{aligned}\therefore u &= c\tau (F'\xi - f'\xi') + F\xi + f\xi' \\ &= c\tau^{\frac{1}{2}} \{F'(x - ct^{\frac{1}{2}}) - f'(x + ct^{\frac{1}{2}})\} \\ &\quad + F(x - ct^{\frac{1}{2}}) + f(x + ct^{\frac{1}{2}}).\end{aligned}$$

Ex. 14. To integrate $D_i^2 u - c^2 t^{-\frac{1}{2}} u = 0$.

The integral of $d_i^2 u - c^2 t^{-\frac{1}{2}} u = 0$ is

$$\frac{u}{t} = C_1 \epsilon^{\frac{c}{t}} + C_2 \epsilon^{-\frac{c}{t}};$$

consequently the integral of the proposed equation is

$$\frac{u}{t} = \epsilon^{\frac{c}{t}} F(\xi, \eta, \zeta \dots) + \epsilon^{-\frac{c}{t}} f(\xi, \eta, \zeta \dots),$$

and c may be a quasi-constant, or a function of $d_x, d_y, \dots$

Ex. 15. To integrate

$$d_t^2 u + \frac{2x}{t} \cdot d_t d_x u + \frac{x^2}{t^2} \cdot d_x^2 u - c^2 x^{-4} u = 0.$$

Here $D_t = d_t + \frac{x}{t} \cdot d_x$, $\xi = \frac{x}{t}$, and $x^{-4} = \frac{t^4}{x^4}$, $t^{-4} = \xi^{-4} \cdot t^{-4}$;

$$\therefore D_t^2 u - c^2 \xi^{-4} \cdot t^{-4} u = 0.$$

This agrees with the form of the preceding Example, with $c\xi^{-2}$ for c .

$$\begin{aligned} \therefore \frac{u}{t} &= \epsilon^{\frac{c}{t\xi^2}} F(\xi) + \epsilon^{-\frac{c}{t\xi^2}} f(\xi) \\ &= \epsilon^{\frac{c}{x^2}} F\left(\frac{x}{t}\right) + \epsilon^{-\frac{c}{x^2}} f\left(\frac{x}{t}\right). \end{aligned}$$

Ex. 16. To integrate $d_t^2 u + cu = \frac{A(A-1)}{t^2} \cdot u$; A being an integer.

Multiply by t^2 ;

$$\therefore (t^2 d_t^2 + ct^2) u = A(A-1) u.$$

For $t^2 d_t^2$ write $d_\theta (d_\theta - 1)$, (Art. 42, 5);

$$\therefore \{d_\theta (d_\theta - 1) - A(A-1)\} \cdot u + ct^2 u = 0;$$

$$\therefore (d_\theta - A)(d_\theta + A - 1) u + ct^2 u = 0.$$

Diminish the symbolic factorials by A (Art. 45);

$$\therefore (d_\theta - 2A)(d_\theta - 1) \cdot ut^A + ct^2 \cdot ut^A = 0,$$

or, writing v for ut^A ,

$$(d_\theta - 2A)(d_\theta - 1)v + ct^2v = 0.$$

We have now to increase the first of the factorials by nk (that is, by $2A$, for $k=2$, and $n=-A$) to reduce them to the requisite form (Art. 44). We must therefore assume (Art. 49)

$$(t^2 d_\theta)^{-A} \cdot vt^{2A} = (t^2)^{-A} \cdot wt^{2A} = w;$$

$$\therefore d_\theta(d_\theta - 1)w + ct^2w = 0;$$

$$\therefore t^2 d_i^2 w + ct^2 w = 0;$$

$$\therefore d_i^2 w + cw = 0.$$

The integral of this equation is known: and u will be known from the equations

$$u = vt^{-A}, \text{ and } v = t^{2A} (t^2 d_\theta)^A w = t^{2A} (t^{-1} d_i)^A w.$$

$$\therefore u = t^A (t^{-1} d_i)^A w.$$

For d_i in this example we may write D_i , and c may be a quasi-constant: if d_i be used, then c may be a function of $d_x, d_y, \dots$

Ex. 17. To integrate

$$d_i^2 u + A d_i u = \frac{2u}{t^2}.$$

This may be written,

$$d_i(d_i + A)u = \frac{2u}{t^2}.$$

For u write $vt^{-\frac{1}{2}A}$;

$$\therefore \left(d_i - \frac{1}{2}A\right)\left(d_i + \frac{1}{2}A\right)v = \frac{2v}{t^2};$$

$$\therefore d_t^2 v - \frac{1}{4} A^2 v = \frac{2v}{t^2},$$

which falls under the last example. We may write D_t for d_t , and a quasi-constant for A .

Ex. 18. To integrate

$$d_t^2 u + a d_t d_x u = \frac{2u}{t^2}.$$

This is equivalent to,

$$d_t (d_t + a d_x) u = \frac{2u}{t^2};$$

$$\therefore D_t = d_t + a d_x, \text{ and } \xi = x - at;$$

$$\therefore d_t D_t u = \frac{2u}{t^2}.$$

Now assume $u = vt^m$, and put $m(m-1) = 2$, which gives $m = 2$, or -1 .

Taking the latter, we have

$$t d_t D_t v = (D_t + d_t) v \dots\dots\dots (1).$$

Differentiate with $d_t D_t$,

$$\therefore d_t^2 D_t^2 v = 0;$$

$$\therefore v = s_t^2 0 + S_t^2 0$$

$$= t F_1(x) + F(x) + t f_1(\xi) + f(\xi).$$

The supernumerary constants are to be determined by substituting this result in equation (1), (see Ex. 4);

$$\therefore v = \frac{at}{2} (f' \xi - F' x) + f \xi + F x;$$

$$\therefore u = \frac{a}{2} (f' \xi - F' x) + \frac{1}{t} (f \xi + F x).$$

Ex. 19. To integrate

$$t^2 d_t^3 u + 2tx d_t d_x u + x^2 d_x^2 u = t^m x^n.$$

Dividing by t^2 ,

$$d_t^3 u + \frac{2x}{t} d_t d_x u + \frac{x^2}{t^2} d_x^2 u = t^{m-2} x^n;$$

$$\therefore D_t = d_t + \frac{x}{t} d_x, \text{ and } \xi = \frac{x}{t};$$

$$\therefore D_t^3 u = t^{m-2} \cdot (\xi t)^n = \xi^n \cdot t^{m+n-2};$$

$$\begin{aligned} \therefore u &= \xi^n \frac{t^{m+n}}{(m+n-1)(m+n)} + tF(\xi) + f(\xi) \\ &= \frac{t^m x^n}{(m+n-1)(m+n)} + tF\left(\frac{x}{t}\right) + f\left(\frac{x}{t}\right). \end{aligned}$$

Ex. 20. To integrate

$$\begin{aligned} t^n d_t^n u + nt^{n-1} x d_t^{n-1} d_x u + n \frac{n-1}{2} t^{n-2} x^2 d_t^{n-2} d_x^2 u + \dots \\ + x^n d_x^n u = t^m \phi\left(\frac{x}{t}\right). \end{aligned}$$

Divide by t^n , then

$$D_t = d_t + \frac{x}{t} d_x, \text{ and } \xi = \frac{x}{t};$$

$$\therefore D_t^n u = t^{m-n} \phi(\xi);$$

$$\begin{aligned} \therefore u &= \frac{t^m \phi(\xi)}{m(m-1)(m-2)\dots(m-n+1)} \\ &\quad + F_1(\xi) + tF_2(\xi) + t^2F_3(\xi) + \dots + t^{n-1}F_n(\xi). \end{aligned}$$

Ex. 21. To integrate

$$t^2 d_t^2 u - x^2 d_x^2 u + t d_t u - x d_x u = 0.$$

$$\therefore d_t^2 u - \frac{x^2}{t^2} \cdot d_x^2 u + \frac{1}{t} \left(d_t u - \frac{x}{t} \cdot d_x u \right) = 0.$$

Comparing this with the formula (1) Art. 30,

$$\Delta_t D_t u = d_t^2 u + (D_t x + \Delta_t x) d_t d_x u + D_t x \cdot \Delta_t x \cdot d_x^2 u + \Delta_t D_t x \cdot d_x u,$$

we have $D_t x = \frac{x}{t}, \Delta_t x = -\frac{x}{t}; \xi = \frac{x}{t}, \xi' = tx,$

$$\therefore \Delta_t D_t x = -\frac{2x}{t^2}, \text{ and } D_t \Delta_t x = 0,$$

hence D_t and Δ_t are not commutative;

$$\therefore t \Delta_t D_t u + D_t u = 0,$$

Integrate with Σ_t ,

$$\therefore t D_t u = F_1(\xi') = F_1(\xi t') = 2\xi t^2 F'(\xi t^2);$$

$$\therefore D_t u = 2\xi t F'(\xi t^2);$$

$$\therefore u = F(\xi t^2) + f(\xi)$$

$$= F(tx) + f\left(\frac{x}{t}\right).$$

Ex. 22. To integrate

$$d_t^2 u + \frac{2}{t} \cdot d_t u - a^2 d_x^2 u = \frac{2}{t^2} \cdot u.$$

$$\therefore t^2 d_t^2 u + 2t d_t u - a^2 t^2 d_x^2 u = 2u;$$

$$\therefore \{d_\theta(d_\theta - 1) + 2d_\theta - 2\} u - a^2 t^2 d_x^2 u = 0;$$

$$\therefore (d_\theta + 2)(d_\theta - 1)u - a^2 t^2 d_x^2 u = 0.$$

For convenience write C for $-a^2 d_x^2$;

$$\therefore (d_\theta + 2)(d_\theta - 1)u + C t^2 u = 0.$$

We have now to diminish the factor $d_\theta + 2$ by 2; this we do by assuming (see Art. 49)

$$(t^{-2} d_\theta) \cdot u t^2 = (t^{-2}) \cdot v t^2 = v \dots\dots\dots (1);$$

$$\therefore d_t(d_t - 1)v + Ct^2v = 0;$$

$$\therefore d_t^2v + Cv = 0;$$

$$\therefore d_t^2v - a^2d_x^2v = 0;$$

$$\therefore v = F(x - at) + f(x + at).$$

From equation (1), we have

$$u = t^{-2}s_t(tv).$$

Ex. 23. To integrate

$$d_t d_x u + \frac{d_t u + d_x u}{t + x} = \frac{2u}{(t + x)^2}.$$

For u write $v(t + x)^m$, and put $m(m + 1) = 2$ which gives $m = 1$ or -2 . We take the latter value of m ;

$$\therefore (t + x) d_t d_x v = d_t v + d_x v \dots\dots\dots (1).$$

Differentiate this with $d_t d_x$, (Art. 39, 2);

$$\therefore (t + x) d_t^2 d_x^2 v = 0;$$

$$\therefore v = s_t^2 s_x^2 0 = s_t^2 0 + s_x^2 0$$

$$= (t + x) F_1(x) + F(x) + (t + x) f_1(t) + f(t).$$

The forms of the supernumerary arbitrary functions are to be determined by substituting this value of v in equation (1). This gives

$$F_1(x) = -\frac{1}{2} F'(x), \text{ and } f_1(t) = -\frac{1}{2} f'(t).$$

$$\therefore u = v(t + x)^{-2} = (t + x)^{-2} (Fx + ft) - \frac{1}{2} (t + x)^{-1} (F'x + f't).$$

Ex. 24. To integrate

$$(t - x) (d_t^2 u - d_x^2 u) = 4.$$

In this example

$$D_t = d_t + d_u, \Delta_t = d_t - d_u, \xi = x - t, \xi' = x + t;$$

$$\therefore (t-x) D_t \Delta_t u = 4;$$

$$\therefore D_t \Delta_t u = -\frac{4}{\xi} = \frac{2\Delta_t \xi}{\xi}, (\Delta_t \xi = -2).$$

Integrate with Σ_t ,

$$\therefore D_t u = 2 \log \xi + F'(\xi);$$

$$\therefore u = 2t \log \xi + F(\xi') + f(\xi).$$

$S_t F'(\xi')$ is not equal to $F(\xi')$; but it is equivalent to it in this case, because $D_t \xi' (= 2)$ is constant, and the constant may be considered as being absorbed in the arbitrary function.

CHAPTER X.

EQUATIONS OF THE SECOND ORDER. COEFFICIENTS
FUNCTIONS OF p AND q .

54. IN this Chapter it will be a convenience to represent $d_i u$ and $d_x u$ by p and q respectively. The following are the formulæ suitable for this class of equations. They are a mere adaptation of the general formula of Art. 7 to these equations, and no new principle is involved in them.

$$D_i p = d_i p + D_x . d_x p = d_i^2 u + D_x . d_i d_x u \dots\dots (1),$$

$$D_x q = d_x q + D_i . d_i q = d_x^2 u + D_i . d_x^2 u \dots\dots (2).$$

From these we immediately deduce the following forms, which it will be observed are such as can be at once adapted to those terms of an equation which are of the second order. They are the forms under which such terms generally present themselves, and will serve as guides in choosing the starting-point in commencing the integration of any proposed equation.

$$D_i p - D_x . D_x q = d_i^2 u - (D_x)^2 . d_i^2 u \dots\dots\dots (3),$$

$$D_i p + D_x . D_i q = d_i^2 u + 2 D_x . d_i d_x u + (D_x)^2 . d_i^2 u \dots\dots\dots (4),$$

$$D_i p + M . D_x q = d_i^2 u + (D_x + M) d_i d_x u + M D_x . d_x^2 u \dots\dots (5).$$



In the last of these forms M is arbitrary. In using the form (4) the value of $D_x x$ will be definitely known by comparison of the right-hand side of the formula with the proposed equation; and it will have only one value; in consequence of which there will be only *one* first integral of the equations that fall under that form. The general integral corresponding thereto may contain two arbitrary functions, but they will be functions of the *same* subject.

In using forms (3) and (5) the arbitrary functions, which will appear in the general integral of equations which fall under those forms, will have *different* subjects. For in form (3) the value of $(D_x x)^2$ will be given by comparison of the right-hand side with the proposed equation; and consequently $D_x x$ may be either (+) or (-) a certain quantity. Hence there may be *two* first integrals, one of them to be found by using the (+) value, and the other by using the (-) value of $D_x x$. Also in using form (5) the quantities that will be known by comparison of the right-hand side with a proposed equation are $D_x x + M$ and $M \cdot D_x x$, in which M and $D_x x$ are evidently interchangeable; consequently by using first one value for $D_x x$ and the other for M , and then interchanging them, we have two different values for $D_x x$, and consequently there may be *two* first integrals, and two *different* subjects of the arbitrary functions that enter into the final integral.

Examples suitable for this Chapter are not very abundant in Treatises on Partial Differential Equations, but of those that are usually given there are some which are interesting from their connection with Geometry or Physics. These introduce us to a new class of integration difficulties,—and in consequence of these, explanations are required which it is by no means easy to give; and of those

explanations that are here given conjecture forms a large feature, but it has been judged better to offer conjectural explanations than to pass over the difficulties in silence. Some good will have been effected if nothing more is done here than to direct the attention of mathematicians to them. The difficulties here referred to are brought to light by the method of integration peculiar to this Essay.

Ex. 1. To integrate

$$q^2 d_t^2 u - 2pq d_t d_x u + p^2 d_x^2 u = 0.$$

Divide by q^2 ,

$$\therefore d_t^2 u - \frac{2p}{q} d_t d_x u + \frac{p^2}{q^2} d_x^2 u = 0.$$

On comparing this with formula (4), we find that

$$D_x x = -\frac{p}{q}, \text{ and } D_t p + D_x x \cdot D_x q = 0;$$

$$\therefore p + D_x x \cdot q = 0, \text{ and } \Gamma_t p - \frac{p}{q} \cdot D_x q = 0;$$

$$\therefore D_t u = 0, \text{ and } \frac{q D_t p - p D_x q}{q^2} = 0;$$

$$\therefore u = \xi, \text{ and } \frac{p}{q} = F(\xi);$$

$$\therefore -D_x x = F(\xi);$$

$$\therefore -x = tF(\xi) + f(\xi),$$

$$\therefore 0 = x + tF(u) + f(u).$$

This is the general equation of a certain class of *Ruled Surfaces*.

Ex. 2. To integrate

$$d_t^2 u \cdot d_x^2 u - (d_t d_x u)^2 = 0.$$

NOTES

This is the characteristic equation of developable surfaces. Its terms are not linear, but from its peculiar form it can be integrated by means of the above formulæ.

Divide by $d_x^2 u$,

$$\therefore d_t^2 u - \frac{d_t d_x u}{d_x^2 u} d_t d_x u = 0.$$

Comparing this with form (1) we have

$$D_t p = 0, \text{ and } D_t x = -\frac{d_t d_x u}{d_x^2 u};$$

$$\therefore p = \xi, \text{ and } d_t d_x u + D_t x \cdot d_x^2 u = 0.$$

And by form (2) we have

$$D_t q = d_t d_x u + D_t x \cdot d_x^2 u = 0;$$

$$\therefore q = F(\xi) = F(p).$$

This is the first integral. The final integral may be thus found:

$$D_t u = p + D_t x \cdot q = \xi + D_t x \cdot F(\xi);$$

$$\therefore u = \xi t + x F(\xi) + f(\xi).$$

But this is subject to the condition that $\xi = p = d_t u$;

$$\therefore \xi = \xi + \{t + x F'(\xi) + f'(\xi)\} \cdot d_t \xi;$$

$$\therefore \text{either } 0 = t + x F'(\xi) + f'(\xi), \text{ or } 0 = d_t \xi = d_t^2 u.$$

Now in *general* $d_t^2 u$ cannot be $= 0$, therefore the general integral of the proposed equation is that which results from the elimination of ξ between the two equations

$$\begin{cases} u = \xi t + x F \xi + f \xi, \\ 0 = t + x F' \xi + f' \xi. \end{cases}$$

55. Let us inquire to what kind of developable surfaces the condition $d_t \xi = 0$ belongs.

Since $d_t \xi = d_t u$, this being substituted in the original equation gives $d_t d_x u = 0$. We have therefore to find that form of u which will satisfy simultaneously the two equations

$$d_t^2 u = 0, \text{ and } d_x d_t u = 0;$$

$$\therefore u = Ct + \phi(x),$$

C being an arbitrary constant and ϕ an arbitrary function. This is the equation of a cylindrical surface. And therefore when the developable surface degenerates into a cylinder, the integral just found must take the place of the general integral.

We may remark that the proposed equation when written in the form $\frac{d_t p}{d_x p} = \frac{d_t q}{d_x q}$ falls under Ex. 6, Chap. 2; and consequently we have at once, as its first integral, $q = F(p)$.

Ex. 3. To integrate

$$q d_t^2 u - p d_t d_x u = 0.$$

This equation, like the preceding one, may be put under the form of Chap. 2, Ex. 6, for it may be written in the form

$$\frac{d_t p}{d_x p} = \frac{d_t u}{d_x u};$$

$$\therefore p = F(u) = \frac{1}{f'(u)} \text{ (suppose) }; \quad$$

$$\therefore f'(u) \cdot d_t u = 1;$$

$$\therefore f u = t + \phi(x).$$

Ex. 4. To integrate

$$q(1+q)d_t^2u - (p+q+2pq)d_t d_x u + p(1+p)d_x^2u = 0.$$

Divide by $q(1+q)$,

$$\therefore d_t^2u - \left(\frac{p}{q} + \frac{1+p}{1+q}\right)d_t d_x u + \frac{p}{q} \cdot \frac{1+p}{1+q} d_x^2u = 0.$$

On comparing this with form (5) of this Chapter, we have the option of assuming

$$D_t x = -\frac{p}{q}, \text{ or } -\frac{1+p}{1+q};$$

the former on trial is found to be the simpler. We therefore assume

$$M = -\frac{1+p}{1+q}, \quad D_t x = -\frac{p}{q}, \quad \text{and} \quad D_t p - \frac{1+p}{1+q} \cdot D_t q = 0;$$

$$\therefore D_t u = p + D_t x \cdot q = 0, \quad \text{and} \quad \frac{D_t p}{1+p} - \frac{D_t q}{1+q} = 0;$$

$$\therefore u = \xi, \text{ and } 1+p = (1+q)F(\xi);$$

$$\therefore p - F(\xi)q = F(\xi) - 1,$$

$$\text{or } p - F(u)q = F(u) - 1.$$

This is an equation of the first order, and may be integrated in the usual manner.

Compare it with the fundamental equation

$$\Delta_t u = p + \Delta_t x \cdot q;$$

$$\therefore \Delta_t u = F(u) - 1, \text{ and } \Delta_t x = -F(u).$$

$$\therefore \Delta_t u + \Delta_x x + 1 = 0, \text{ and } \frac{\Delta_t u}{F(u) - 1} = 1;$$

$$\therefore u + x + t = \xi', \text{ and } \phi u = t + f(\xi');$$

$$\therefore \phi(u) = t + f(u + x + t)$$

is the integral required.

Ex. 5. To integrate

$$(1 + pq + q^2) d_t^2 u + (q^2 - p^2) d_x d_x u - (1 + pq + p^2) d_x^2 u = 0.$$

We notice that

$$q^2 - p^2 = (1 + pq + q^2) - (1 + pq + p^2);$$

let this be substituted.

$$\therefore d_t(p + q) - \frac{1 + pq + p^2}{1 + pq + q^2} d_x(p + q) = 0.$$

Compare this with the general formula

$$D_t(p + q) = d_t(p + q) + D_x x \cdot d_x(p + q);$$

$$\therefore D_t(p + q) = 0, \text{ and } D_x x = -\frac{1 + pq + p^2}{1 + pq + q^2};$$

$$\therefore p + q = \xi, \text{ and } D_x x = -\frac{1 + p\xi}{1 + q\xi};$$

$$\therefore 1 + D_x x + \xi(p + D_x x \cdot q) = 0.$$

Integrate with S_t , remembering that $p + D_x x \cdot q = D_t u$,

$$\therefore t + x + \xi u = F(\xi) \dots \dots \dots (1);$$

$$\therefore t + x + (p + q) u = F(p + q).$$

This is a first integral of the proposed equation.

This equation may be integrated in the following manner.

By inspection of equation (1), we perceive that

$$\xi = p + q = \phi(u, t + x).$$

Comparing this with the general formula,

$$\Delta_t u = p + \Delta_t x \cdot q,$$

we find

$$\Delta_t x = 1, \text{ and } \Delta_t u = \phi(u, t + x) = \xi;$$

$$\therefore x = t + \xi', \text{ and } t + x + u\xi = F(\xi) \text{ from (1).}$$

Divide the last equation by ξ , and differentiate with Δ_t ,

$$\therefore \frac{2 + \xi^2}{\xi} - (t + x) \cdot \frac{\Delta_t \xi}{\xi^2} = \Delta_t \cdot \frac{F(\xi)}{\xi}.$$

Multiply by 2, which is $= \Delta_t(t + x)$, and divide by $\sqrt{2 + \xi^2}$;

$$\therefore \frac{\sqrt{2 + \xi^2}}{\xi} \cdot \Delta_t(t + x) - (t + x) \cdot \frac{2\Delta_t \xi}{\xi^2 \sqrt{2 + \xi^2}} = \frac{2}{\sqrt{2 + \xi^2}} \cdot \Delta_t \frac{F\xi}{\xi}.$$

Integrate with Σ_t ;

$$\therefore \frac{\sqrt{2 + \xi^2}}{\xi} \cdot (t + x) = \Sigma_t \frac{2\Delta_t(\xi^{-1} F\xi)}{\sqrt{2 + \xi^2}} + f(\xi') \dots\dots(2).$$

If now we eliminate ξ between this equation and equation (1), and write $x - t$ for ξ' , we shall have the required integral.

But equation (2) contains a term affected with the sign of integration. We are at liberty to write $F_1(\xi)$ for that term, but this would introduce the sign of integration into equation (1).

The integral of the proposed equation may be found in a more convenient shape from the second first integral, as follows :

The proposed equation may be thrown into the form,

$$(1 + pq + q^2) (d_t^2 u + d_t d_x u) = (1 + pq + p^2) (d_t d_x u + d_x^2 u).$$

On reference to formulæ (1) and (2), we see that if we assume $\Delta_t x = 1$, which gives $\xi' = x - t$, then

$$(1 + pq + q^2) \Delta_t p = (1 + pq + p^2) \Delta_t q.$$

For brevity write ξ for $p + q$, and express this equation in terms of $p + q$ and $p - q$;

$$\therefore (p - q) \xi \Delta_t \xi = (2 + \xi^2) \Delta_t (p - q).$$

Integrate with Σ_t ,

$$\therefore \sqrt{2 + \xi^2} = (p - q) \cdot f(\xi') \dots\dots\dots (3).$$

This is the second first integral. It may be integrated in the following way :

$$2du = 2pdt + 2qdx$$

$$= (p + q) (dt + dx) + (p - q) (dt - dx)$$

$$= \xi (dt + dx) - \sqrt{2 + \xi^2} \cdot \frac{d\xi'}{f(\xi')} \text{ from (3) ;}$$

$$\therefore \left. \begin{aligned} 2u &= \xi(t + x) - \sqrt{2 + \xi^2} \cdot f_1(\xi') + F_1(\xi) \\ 0 &= t + x - \frac{\xi f_1(\xi')}{\sqrt{2 + \xi^2}} + F_1'(\xi) \end{aligned} \right\} \dots\dots\dots (4).$$

If we eliminate ξ between these equations, and write $x - t$ for ξ' , we shall have the required integral.

The advantage of this integral over the one before found in (1) and (2) is, that it contains no quantities under the sign of integration.

The above Example may be put under a somewhat more general form which can be integrated in the same manner. It is here added as an exercise for the student.

To integrate

$$(k + pq + aq^2) (d_t^2 u + a d_t d_x u) = (h + apq + p^2) (d_t d_x u + a d_x^2 u).$$

Ex. 6. To integrate

$$q^2 d_t^2 u - p^2 d_x^2 u = 0.$$

Divide by q^2 and compare with formula (3);

$$\therefore (D_t x)^2 = \frac{p^2}{q^2}, \text{ and } D_t p - D_x q = 0.$$

There are two values of D_x , we take the negative form as being the simpler:

$$\therefore D_x = -\frac{p}{q}, \text{ and } D_t p + \frac{p}{q} \cdot D_x q = 0;$$

$$\therefore D_t u = p + q D_x u = 0, \text{ and } pq = F(\xi);$$

$$\therefore u = \xi; \therefore pq = F(u).$$

This is a first integral. It may be integrated as follows. Let θ be a new dependent variable, such that

$$p = \theta \sqrt{Fu}, \text{ and } \therefore q = \frac{1}{\theta} \cdot \sqrt{Fu}.$$

Then, because

$$\begin{aligned} du &= p dt + q dx, \\ \therefore \frac{du}{\sqrt{Fu}} &= \theta dt + \frac{dx}{\theta}; \end{aligned}$$

$$\left. \begin{aligned} \therefore \phi(u) &= \theta t + \frac{x}{\theta} + f(\theta) \\ 0 &= t - \frac{x}{\theta^2} + f'(\theta) \end{aligned} \right\} \dots\dots\dots(1).$$

The elimination of θ between these two equations will give the required integral.

This integral has been obtained from one of the values of $D_x x$; and this value has led us to the equation

$$F(u) = pq$$

as the corresponding first integral. It is easy to obtain the second first integral, if required, by employing the second value of $D_x x$. In this case we must introduce the symbol Δ_x , and the equations we have to deal with are

$$\Delta_x x = \frac{p}{q}, \text{ and } \Delta_x p - \Delta_x x \cdot \Delta_x q = 0.$$

From these we can obtain two integrable equations. For eliminating $\Delta_x x$ we have

$$\Delta_x p - \frac{p}{q} \Delta_x q = 0;$$

$$\therefore \frac{p}{q} = \xi;$$

$$\therefore \Delta_x x = \xi';$$

$$\therefore x = \xi' t + \phi(\xi')$$

$$= \frac{p}{q} \cdot t + \phi\left(\frac{p}{q}\right).$$

This is the second first integral, and if we substitute θ^2 for its equal $\frac{p}{q}$, we shall see that this first integral constitutes the second of the two equations (1). The final integral de-

rived from this equation will be found to be the same as that already found and expressed by the equations (1).

56. Hitherto we have been able to proceed to the final integrals of the Examples proposed without meeting with any impediment. But we shall not be able to do so in the remaining Examples of this Chapter. These will bring us face to face with difficulties which require careful consideration. But when they are resolved they seem to have constituted an essential part of the process of integration, because they have forced us out of the track of ordinary routine when ordinary routine was inadequate to the task of integration. They teach us that differential equations, which fall under this class, sometimes represent Geometrical properties or Physical phenomena which an ordinary integral (an integral which can be obtained in the ordinary way) cannot express. In fact the difficulties which force us out of the beaten track are necessary, because the integral which is to be found has to express results which cannot be expressed by any single integral. Two integrals are found to be necessary, each of which is independent of the other; and yet though capable of being solitary, they are to be also capable of being joined together by a single link. This language will seem mysterious at present, but it will be explained by the examples which follow.

An important question will also be suggested and in one instance answered,—must there necessarily be an integral of every partial differential equation that can be proposed? We may write down a partial differential equation at random, does it necessarily express possible geometrical or physical properties? and if it does not, ought we to expect to be able to find an integral of it? If the conditions and requirements

of a problem be of such a nature that the variables cannot satisfy them without assuming the attributes of infinity, as in Example 10 of this Chapter, is it reasonable to expect that the differential equation which embodies them can admit of an integral?

There may be degrees of impossibility of this kind. As in ordinary Algebra there are formulæ which are true for one set of numbers (say, *odd* numbers) and untrue for any other set of numbers (say, *even* numbers); and other formulæ which are true for *even* numbers and untrue for *odd* numbers; so there may be two forms of integrals of the same equation, one of which may be true for one set of values of the variables, and the other for another set. How is it to be indicated in an *integral*, when there is no indication of it in the partial differential equation from which the integral was derived, that one of the forms of the integral is to be used in one set of circumstances, and in other circumstances the other?

The next equation we shall give for integration is derived from Ex. 1 by *arbitrarily* changing the algebraic sign of the middle term. The equation in Ex. 1 represents a family of Ruled Surfaces, what does the equation in Ex. 7 represent? Does it represent a possible property either in Geometry or Physics?

Ex. 7. To integrate

$$q^2 d_x^2 u + 2pq d_x d_y u + p^2 d_y^2 u = 0.$$

Compare this, after dividing it by q^2 , with formula (4);

$$\therefore D_y x = \frac{p}{q}, \text{ and } D_y p + D_x x \cdot D_y q = 0 \dots\dots\dots(1).$$

These two combined give

$$qD_t p + pD_x q = 0;$$

$$\therefore pq = \xi.$$

No other integrable equation can be obtained from equations (1). Here then we meet with only partial success. What use can we make of this degree of success? It must be of some value. The following has suggested itself.

Let us now abandon the equations (1), and use our success as a hint to change the *dependent* variable from u to pq (or ξ). This change of the dependent variable being made, the proposed equation takes the form

$$d_t(pq) + \frac{p}{q} \cdot d_x(pq) = 0.$$

Now, on comparing this with the general formula,

$$D_t(pq) = d_t(pq) + D_x \cdot d_x(pq)$$

we perceive that we must have

$$D_t(pq) = 0,$$

an equation which can be satisfied in two ways only, viz. by supposing

either $pq = \text{constant}$; or $pq = \text{quasi-constant}$.

The latter involves the unintegrable equation $D_x = \frac{p}{q}$; but the former evades this equation. The only way of arriving at a final integral is therefore the supposition,

$$pq = C, \text{ a constant.}$$

We consider the impossibility of forming *two* integrable equations out of equations (1) to be a necessary incident,—necessary to force us upon this alternative equation.

$$\begin{aligned} \therefore du &= p dt + q dx \\ &= p dt + \frac{C}{p} dx; \\ \therefore u &= pt + \frac{C}{p} x + F(p) \\ 0 &= t - \frac{C}{p^2} x + F'(p) \end{aligned} \left. \vphantom{\begin{aligned} \therefore du &= p dt + q dx \\ &= p dt + \frac{C}{p} dx; \\ \therefore u &= pt + \frac{C}{p} x + F(p) \\ 0 &= t - \frac{C}{p^2} x + F'(p) \end{aligned}} \right\}.$$

The elimination of p between these equations will give the only possible integral; which therefore contains *one arbitrary function, and one independent arbitrary constant*. According to the usual rule, there might be in a general integral *two* arbitrary functions, but in this Example the place of one of the arbitrary functions is supplied by an independent arbitrary constant.

Ex. 8. To integrate

$$d_t^2 u + 2p d_t d_x u + (p^2 - a^2) d_x^2 u = 0.$$

Since $2p = (p+a) + (p-a)$, we may compare this equation with formula (5),

$$D_t p + M D_t q = d_t^2 u + (D_t x + M) d_t d_x u + M D_t x \cdot d_x^2 u;$$

$$\therefore \text{either } D_t x = p + a, \text{ and } M = p - a;$$

$$\text{or } D_t x = p - a, \text{ and } M = p + a.$$

We begin with the former of these;

$$\therefore D_t x = p + a, \text{ and } D_t p + (p - a) D_t q = 0;$$

$$\therefore (p - a) \epsilon^2 = \xi.$$

The equation $D_x x = p + a$ cannot be integrated; we therefore proceed as in the last Example. We take $(p - a) \epsilon^q$ for a new dependent variable, and this conducts us to the equation

$$(p - a) \epsilon^q = C, \text{ a constant.}$$

And this being integrated in the same manner as before, gives

$$\left. \begin{aligned} u &= (C\epsilon^{-q} + a) t + qx + F(q) \\ 0 &= -C\epsilon^{-q} t + x + F'(q) \end{aligned} \right\} \dots\dots\dots (1),$$

and if we eliminate q between these equations we shall have a general integral of the proposed equation.

We say a general because we have another set of equations to deal with, viz.

$$D_x x = p - a, \text{ and } M = p + a;$$

which are also included in the proposed equation. The resulting integral in this case is

$$\left. \begin{aligned} u &= (C'\epsilon^{-q} - a) t + qx + f(q) \\ 0 &= -C'\epsilon^{-q} t + x + f'(q) \end{aligned} \right\} \dots\dots\dots (2).$$

Thus we have in an independent manner obtained two integrals of equal authority. We have now to examine the significance of this circumstance, which it is obvious could not have occurred had an integral been obtainable in the ordinary manner. Let us examine the two integrals.

1. The two arbitrary constants C and C' are not necessarily equal. Nothing in the processes of integration necessitates their equality. So likewise the two functions F and f are not necessarily equal, being absolutely arbitrary.

2. Nevertheless, if we assume $t=0$, the two integrals become

$$\left. \begin{aligned} u &= qx + F(q) \\ 0 &= x + F'(q) \end{aligned} \right\} \dots\dots\dots (3),$$

and

$$\left. \begin{aligned} u &= qx + f(q) \\ 0 &= x + f'(q) \end{aligned} \right\} \dots\dots\dots (4),$$

which place before us the following alternative:

(α). Either F and f must be equal;

or (β). The two integrals cannot both be satisfied with the same values of x ; that is, they cannot cover the same ground.

3. If F and f are equal, write in equations (2) F for f ; and the same in equations (4).

Two circumstances will now be evident.

(γ). The equations (3) and (4) exactly agree; but equations (1) and (2) differ, and cannot be made to agree even by assuming $C' = C$.

(δ). The phenomena represented by the first and second integrals (1) and (2) exactly agree when $t=0$; but they disagree for all other values of t .

If for easiness of expression we call t time, and x place, we may express these results (γ), (δ), as follows:—

When F and f are equal, the two integrals (1), (2) express phenomena which had a common origin at the time $t=0$; but afterwards they become essentially distinct,—*solitary*, not existing in the same place. If t have the same value in both integrals, x cannot have the same value in both. The

phænomena therefore which coexisted both in time and place at their origin afterwards become solitary,—each existing in its own place.

In their most general form the two integrals of equal authority which we have obtained represent phænomena which do not coexist in space, and are every way independent of each other. But in the restricted forms obtained by assuming F equal to f they represent phænomena which have no other connexion than this,—that they are due to a common originating cause.

The general inference therefore is,—that the proposed differential equation embodies properties or phænomena which may be either,—1st, absolutely independent, and capable of existing separately and alone; or, 2nd, dependent in their origin, but afterwards independent, and incapable of coexisting with the same values of co-ordinates t and x .

The Wave of Sound.

The two preceding Examples will in some measure have prepared us for entering on the integration of the dynamical equation for the transmission of the sound-wave through a thin straight horizontal tube of elastic medium.

Ex. 9. To integrate

$$d_t^2 u = \frac{a^2}{q^2} \cdot d_x^2 u.$$

This is the dynamical equation for the motion of the sound-wave. On comparing it with formula (1), we find

$$(D_x)^2 = \frac{a^2}{q^2}; \text{ and } D_t p - D_x \cdot D_t q = 0.$$

We have therefore two cases to consider arising from the two equations,

$$D_x x = \frac{a}{q}, \text{ and } \Delta_x x = -\frac{a}{q}.$$

We begin with the former of these. Both of them are unintegrable; but the former enables us to obtain the following integrable equation,

$$\therefore D_t p - \frac{a}{q} \cdot D q = 0,$$

$$\therefore q \epsilon^{-\frac{p}{a}} = \xi.$$

On account of the impossibility of obtaining another integrable equation with the help of $D_x x = \frac{a}{q}$ on which the integral just obtained depends, we are obliged to take $q \epsilon^{-\frac{p}{a}}$ as the new dependent variable instead of u ; and this leads to the equation,

$$q \epsilon^{-\frac{p}{a}} = C, \text{ a constant;}$$

whence we find, as in the preceding examples,

$$\left. \begin{aligned} u &= pt + C \epsilon^{\frac{p}{a}} x + F(p) \\ 0 &= t + \frac{C}{a} \epsilon^{\frac{p}{a}} x + F'(p) \end{aligned} \right\} \dots\dots\dots(1).$$

The elimination of p between these equations will give us one of the general integrals. The other will be found by writing $-a$ for a .

$$\left. \begin{aligned} \therefore u &= pt + C' \epsilon^{-\frac{p}{a}} x + f(p) \\ 0 &= t - \frac{C'}{a} \epsilon^{-\frac{p}{a}} x + f'(p) \end{aligned} \right\} \dots\dots\dots(2).$$

All the remarks that were made on the two general integrals of the last Example might be repeated here; but as in this example we know the meaning of the symbols t, x ; and also know the origin of the proposed differential equation, we can use more definite language here than there.

The origin of the proposed differential equation is this:— Given a uniform elastic medium (as air) at rest. A sudden short disturbance occurs at a particular part of the medium. The details of the original disturbance are given. Find the state of the medium at any subsequent time. This is expressed in differential symbols in the proposed equation.

We know that a denotes the value of the fraction $\frac{\text{pressure}}{\text{density}}$; x the distance of a particle of air from a fixed section of the tube before it was disturbed; and u the corresponding distance of the same particle when in motion at the time t . Also $p = d_t u = v$ the velocity of the same particle at the same time, and

$$q = d_x u = \frac{p}{\rho_0},$$

where ρ_0 is the equilibrium density of every part of the medium, and ρ the density at the distance u and time t .

Hence, in reference to the first general integral (1), the equation,

$$q e^{-\frac{p}{a}} = C, \text{ becomes } \frac{p}{\rho_0} e^{-\frac{v}{a}} = C.$$

But in the forefront of a wave ρ becomes ρ_0 and $v = 0$, which values being written in the equation for ρ and v give

$$C = 1;$$

$$\therefore \text{ always } \rho = \rho_0 e^{\frac{v}{a}} \dots\dots\dots (3).$$

In a similar way we arrive in the case of the second integral at the equations

$$C' = 1, \text{ and } \rho = \rho_0 e^{-\frac{v}{a}} \dots\dots\dots(4).$$

It is evident then that (3) and (4) cannot coexist in the same part of the tube, since (3) exists where ρ is greater, and (4) where it is less, than ρ_0 .

Hence the two integrals (1) and (2) cannot both hold good at the same time in the same part of the tube.

Taken separately (supposing v positive) (1) represents a wave of condensation (a positive wave); and (2) a wave of rarefaction (a negative wave); and consequently such waves if they exist at the same time must be in different parts of the tube.

We shall now shew how two such waves may have a common origin, in which case they must both be embraced in a connected system of general integrals.

Let us suppose a rigid diaphragm in the middle of the tube, and let this diaphragm suddenly move forward through a short space. The result of this *one* movement will be the generation of *two* waves,—a positive wave on the front of the diaphragm, and a negative wave on the back of the diaphragm. In the strictest sense the two waves have a common origin, and they are alike included in the same differential equation of motion. It is the work of integration to bring to light the fact that one of them ever remains on one side of the diaphragm, and the other on the other side; and that the motion of one wave is in one direction, and that of the other in the opposite direction, while yet v has the same algebraic sign in both.

Now the motion of the diaphragm, or (which is the same thing) of the particles of air in contact with the diaphragm on both its sides, is the datum from which both waves are to be determined. If we take the original position of the diaphragm to be where $x=0$, we see that (since $C=C'=1$) for the particles in contact with the diaphragm, we have from equations (1), on the front side, where $x=0$,

$$\left. \begin{aligned} u &= pt + F(p) \\ 0 &= t + F'(p) \end{aligned} \right\},$$

and from equations (2), on the back side, where $x=0$,

$$\left. \begin{aligned} u &= pt + f(p) \\ 0 &= t + f'(p) \end{aligned} \right\}.$$

The comparison of these sets with one another shews us that $F=f$, since t and p (or v) are the same in both sets of equations.

Hence when a positive and a negative wave have a common genesis, the former will be represented by equations (1) and the latter by equations (2) when F is written for f . And when this is done, the two sets of equations (1) (2) taken together embrace the *whole* effect due to the motion of the diaphragm, and consequently for that case represent the complete integral of the proposed equation.

Here then we have a device by which Analytical Symbols meet all possible cases. We may have to consider the properties of a solitary wave, it may be a wave of condensation; or it may be a wave of rarefaction; we are furnished with appropriate integrals, which in themselves shew that they cannot be used together for the same part of the medium, corresponding to the physical fact that two such waves cannot exist together in the same place. Or we may have

given, the fact that an agitation has been produced in one place in the cylinder, and we know that two waves moving in opposite directions will result from it. The originating cause is one, and the arbitrary function which has to represent that cause is one. The two integrals become connected by writing F for f ; but still the integrals are *two*, and though connected in the way just mentioned cannot be combined together so as to be coexistent in the same part of the cylinder, all which corresponds accurately with physical fact.

Thus what at first seems to be a failure of integration is a necessary part of the process for obtaining the two integrals which may be taken separately; or when taken together constitute the complete integral of the proposed equation.

These ideas are the necessary results of the method of integration proposed in this Essay. They are put forward with diffidence for the consideration of mathematicians who have more leisure to devote to such difficulties than the author.

One thing is evident, that the existence of an accumulation of medium in one part of the cylinder necessitates the existence at the same time of an equal deficiency of medium in another part of the cylinder, since the motion of the diaphragm in the genesis of the two waves cannot alter the whole quantity of medium in the tube: and as these two effects of one cause are coexistent in time but not in place, two integrals seem to be inevitably necessary. If necessary, by what means is the process of integration to force them upon us? In the last two Examples we have seen how this may be done. No doubt Analytical Symbols may in other cases have other methods of accomplishing the same object.

Equal Radii of Curvature of Contrary Signs.

An example will now be given which will bring before us a difficulty of another kind. Mathematicians have investigated the partial differential equation which embodies the condition, that at each point of a surface the two principal radii of curvature are equal and of opposite signs. And this equation has been proposed for integration with the object of finding the general equation of the family of surfaces which all possess this property. This will be our next example.

Ex. 10. To integrate

$$(1+p^2) d_i^2 u - 2pq d_i d_x u + (1+q^2) d_x^2 u = 0.$$

Divide by $1+p^2$ and compare the equation with formula (5).

$$\therefore D_i x + M = -\frac{2pq}{1+p^2}, \text{ and } M \cdot D_i x = \frac{1+q^2}{1+p^2};$$

$$\text{and } D_i p + M D_i q = 0.$$

The first two of these equations give us two values for $D_i x$, and two corresponding values for M . But neither of the values of $D_i x$ will enable us to obtain an integrable equation. It is not, as in the preceding Examples, that we are furnished with one integrable and one unintegrable equation. In this example nothing integrable can be found.

Let us endeavour to find out the reason of this difficulty.

If at any point of a surface C_1 , C_2 represent the principal curvatures, and C any other curvature at the same point, we have the following equation, connecting these quantities together, in Treatises on the Geometry of Surfaces,

$$C = C_1 \cos^2 \theta + C_2 \sin^2 \theta.$$

The meaning of θ is well known. Now if $C_1 = -C_2$, which is the case in this example, we have

$$C = C_1 \cos 2\theta.$$

And this equation shews that

as θ passes from -45° to $+45^\circ$, C is +;
 $+45^\circ$ to 135° , C is -;
 135° to 225° , C is +;
 225° to 315° , C is -.

Thus at a point of a surface where the principal radii of curvature are equal and of contrary signs, the surface will be corrugated, the corrugations tending *from* the point as a centre, and being alternately above and below the tangent plane at the point. Briefly we may say, such a point is a *centre of corrugation*. But from the very nature of a continuous surface *every* point of it cannot be a centre of corrugation; and consequently there is no surface that possesses the property embodied in the proposed differential equation. Are we not correct in expecting that such an equation cannot have an integral?

Yet there is one way of finding an integral,—to assume that simultaneously we must have

$$d_t^2 u = 0, \quad d_t d_x u = 0, \quad d_x^2 u = 0.$$

These satisfy the proposed differential equation and give us the following integral,

$$u = At + Bx + C,$$

which is the equation of a plane surface, and is in fact the only answer the proposed problem admits of.

Strictly speaking, however, this integral is neither an answer to the proposed problem, nor an integral of the proposed equation. For a plane surface has no curvature, and consequently also no radii of curvature: and the three equations $d_t^2 u = 0$, $d_t d_x u = 0$, $d_x^2 u = 0$ are an evasion of the proposed equation, not an equivalent for it.

CHAPTER XI.

SUPPLEMENTAL ARTICLES.

IN the preceding Chapters the forms of quasi-constants have been determined by comparison of a proposed equation with general formulæ; and this has been found sufficient for the majority of equations generally proposed for integration in the ordinary Works upon this subject. But the principle on which the *Method* of this Essay is founded is embodied in the right use of the equation

$$D_t u = d_t u + D_x x \cdot d_x u + D_y y \cdot d_y u + \dots$$

or, more generally,

$$D_t u = D_t t \cdot D_t u + D_x x \cdot d_x u + D_y y \cdot d_y u + \dots$$

The use of other formulæ is a convenience in most cases, but not a necessity: and there are equations and classes of equations, the successful integration of which will depend solely on a skilful use of the above fundamental form whereby the change of the system of independent variables is effected.

To the skilful use of this form we must look for an extension of success into fresh fields of difficulties. An example or two will illustrate what is here meant.

I. Ex. 1. To integrate

$$x = \frac{d_t u}{d_x u} t + \phi \left(\frac{d_t u}{d_x u} \right).$$

None of our formulæ are of this form, we therefore arbitrarily assume that the new system of variables is such as to allow us to write

$$D_t x = - \frac{d_t u}{d_x u}.$$

The reason of the minus sign being assumed in this equation will be seen in the next step,—it gives us an integrable equation.

$$\therefore d_t u + D_t x \cdot d_x u = 0 = D_t u;$$

$$\therefore u = \xi.$$

For brevity write v for $\frac{d_t u}{d_x u}$ in the proposed equation,

$$\therefore x = vt + \phi(v) \dots\dots\dots (1).$$

Differentiate this equation with D_t , remembering that $D_t x = -v$;

$$\therefore 2v + t D_t v + \phi'(v) \cdot D_t v = 0.$$

Divide by $\sqrt{v}$ and integrate with S_t ;

$$\therefore 2t\sqrt{v} + s_v \left(\frac{\phi'v}{\sqrt{v}} \right) = F(\xi) \dots\dots\dots (2).$$

Substituting u for ξ and eliminating v between equations (1) and (2), we have the integral required.

The above is applicable to two independent variables only, but the following example is so stated as to be applicable

to any number of independent variables, $D_i u$ being equal to $d_i u + D_i x \cdot d_x u + D_i y \cdot d_y u + \dots$; and the quantity represented by Q is under no restriction but this, that $D_i Q$ is to be either a constant or a quasi-constant.

Ex. 2. To integrate

$$Q + u D_i u = \phi(D_i u).$$

For brevity write v for $D_i u$,

$$\therefore Q + uv = \phi(v) \dots \dots \dots (1).$$

Divide this by v and differentiate with D_i , remembering that $D_i u = v$;

$$\therefore \frac{D_i Q + v^2}{v} - \frac{Q}{v^2} \cdot D_i v = D_i \left(\frac{\phi v}{v} \right).$$

Divide by $\sqrt{D_i Q + v^2}$ and integrate with S_i ;

$$\therefore \left(\frac{D_i Q}{v^2} + 1 \right)^{\frac{1}{2}} \cdot \frac{Q}{D_i Q} + S_i (D_i Q + v^2)^{-\frac{1}{2}} D_i \left(\frac{\phi v}{v} \right) = F(\xi, \eta, \zeta, \dots);$$

$$\text{or, } \left(\frac{D_i Q}{v^2} + 1 \right)^{\frac{1}{2}} \cdot \frac{Q}{D_i Q} + S_i (D_i Q + v^2)^{-\frac{1}{2}} d_v \left(\frac{\phi v}{v} \right) = F(\xi, \eta, \zeta, \dots).$$

We must substitute for $\xi, \eta, \zeta, \dots$ their values and then eliminate v between this and equation (1); and the result will be the required integral.

The following are particular examples of this class.

(3) To integrate

$$ht + kx + u(d_i u + a d_x u) = \phi(d_i u + a d_x u).$$

(4) To integrate

$$ht + kx + u \left(d_t u + \frac{x}{t} d_x u \right) = \phi \left(d_t u + \frac{x}{t} d_x u \right).$$

(5) To integrate

$$ht + kx + ly + u \left(d_t u + \frac{x}{t} d_x u + ad_y u \right) = \phi \left(d_t u + \frac{x}{t} d_x u + ad_y u \right).$$

II. The chief object in Chap. VIII. was to shew how to deprive a particular class of linear equations of coefficients which were not constant, in order to reduce them to linear equations with constant coefficients. These being always integrable, we were enabled to ascend from their integrals to those of the proposed equations. But in Chap. VII. we were often conducted for the value of u to a result of one of the forms following:

$$(\phi d_t)^n . u = 0, \quad \text{or} \quad (\phi d_t)^n t (\phi d_t)^{1-n} u = 0.$$

It is desirable to exhibit the direct dependence of u on v ; v being an auxiliary quantity defined by the equation

$$(\phi d_t) v = 0 \dots \dots \dots (1).$$

By actual integration we have

$$s_t^n 0 = A_1 + A_2 t + A_3 t^2 + \dots + A_n t^{n-1};$$

that is, $s_t^n 0$ will represent a series in integer powers of t from 0 to $n-1$, the coefficients of which are all *arbitrary*. We shall therefore take $s_t^n 0$ to represent any series of the above form when all its coefficients are arbitrary. It is to be distinctly understood that the series represented by $s_t^n 0$ is *definite* in form and *indefinite* in the coefficients of its terms. This will give rise to a peculiarity in its use,—it will enable us with proper explanations to use the sign of *equality* in

some cases where the strict logic of the steps would require the use of a sign of *equivalence* only.

Since $s_i 0$ is constant, by multiplying equation (1) by it we have

$$(\phi d_i) \cdot v s_i 0 = 0 \dots \dots \dots (2).$$

Also, since $s_i^2 0$ contains t to the first power only, if we operate with (ϕd_i) on $v s_i^2 0$, by formula (1) Art. 39 we have

$$\begin{aligned} (\phi d_i) \cdot v s_i^2 0 &= s_i^2 0 \cdot (\phi d_i) v + s_i 0 \cdot (\phi' d_i) v \\ &= (\phi' d_i) \cdot v s_i 0, \text{ by (1);} \end{aligned}$$

$$\begin{aligned} \therefore (\phi d_i)^2 \cdot v s_i^2 0 &= (\phi' d_i) (\phi d_i) \cdot v s_i 0 \\ &= 0, \text{ by (2).} \end{aligned}$$

We see then that the formula

$$(\phi d_i)^n \cdot v s_i^n 0 = 0 \dots \dots \dots (3)$$

holds good for $n=1$, and $n=2$. We see also that it holds good in these cases because v satisfies equation (1); and that in consequence of this we may write for v any quantity whatever (or the sum of any such quantities) which satisfies equation (1). The form (3) will therefore hold good for $n=1, 2$, if we write for v any quantity such as $(\phi' d_i) v$, or $(\phi'' d_i) v$, ... because each of these when written for v in equation (1) will satisfy it as well as v .

We will now suppose that equation (3) is true for all integer values of n from 1 to $(n-1)$; and we shall shew that in that case it is true for the next value of n .

By the formula of Art. 36 we have

$$(\phi d_i) \cdot v s_i^n 0 = s_i^n 0 \cdot (\phi d_i) v + s_i^{n-1} 0 \cdot (\phi' d_i) v + \frac{1}{1 \cdot 2} s_i^{n-2} 0 \cdot (\phi'' d_i) v + \dots$$

$$= s_i^{n-1} 0 \cdot (\phi' d_i) v + \frac{1}{1 \cdot 2} s_i^{n-2} 0 \cdot (\phi'' d_i) v + \dots$$

But each term of the right hand will vanish, by hypothesis, if it be operated on with $(\phi d_i)^{n-1}$;

$$\therefore (\phi d_i)^n \cdot v s_i^n 0 = 0 \dots \dots \dots (3).$$

This equation is therefore universally true for positive integer values of n , both for v and for any quantity instead of v which will satisfy equation (1).

Now if equation (1) be a linear equation of the m^{th} order, we know that the complete value of v will consist of the sum of m independent partial values, each of which will satisfy equation (1); and consequently the left-hand member of equation (3) will be made up of the sum of m parts, each of which independently satisfies it.

Now a glance at the process by which equation (3) has been proved will convince us that it is not necessary that the arbitrary constants in $s_i^n 0$ should have the same values in each of the independent parts of v . We therefore interpret the symbol $v s_i^n 0$ as meaning that *each part of v* is to be multiplied by $s_i^n 0$; and that $s_i^n 0$ is to be considered in each of the parts to be independently arbitrary.

From this explanation it will be seen that if $(\phi d_i) \cdot v = 0$ be of the m^{th} order, v involves m independent partial integrals; and, as $s_i^n 0$ involves n arbitrary constants which are different for each part of v , the expression $v s_i^n 0$ will involve mn arbitrary constants.

If, therefore, we have such an equation as the following,

$$(\phi d_i)^n \cdot u = 0 \dots \dots \dots (4),$$

by comparing it with equation (3) we have the following as

its integral,

$$u = v s_i^n 0 \dots\dots\dots (5),$$

and this is the *complete* integral of equation (4), because it contains mn arbitrary constants, and equation (4) is of the degree mn .

We are now able also to bring u in the equation,

$$(\phi d_i)^n t (\phi d_i)^{1-n} u = 0 \dots\dots\dots (6),$$

into connexion with v ; for on comparing this equation with equation (4), we have here $t (\phi d_i)^{1-n} u$ for u there; and consequently by equation (5) we have

$$t (\phi d_i)^{1-n} u = v s_i^n 0 \dots\dots\dots (7).$$

Now in dividing this equation by t we must remember that $s_i^n 0$ is a series in integral powers of t from t^0 to t^{n-1} ; and therefore, if we divide it by t , the quotient will consist of two parts, one of which contains powers of t from t^0 to t^{n-2} , and which may therefore be represented by $s_i^{n-1} 0$; and the other will be represented by $t^{-1} s_i 0$;

$$\therefore (\phi d_i)^{1-n} u = v s_i^{n-1} 0 + \frac{v}{t} s_i 0.$$

Operate on both sides of this equation with $(\phi d_i)^{n-1}$, {see equation (3)},

$$\therefore u = (\phi d_i)^{n-1} \cdot \frac{v}{t} s_i 0 \dots\dots\dots (8).$$

This is the integral of equation (6).

REMARK. $s_i 0$ will (as before explained) be different for every part of v ; hence the value of u just found will contain m arbitrary constants if $(\phi d_i)v$ be of the m^{th} order.

It may also be remarked, that we are at liberty to write D_i for d_i ; in which case

$$\phi(D_i)v = 0$$

will denote a linear partial differential equation of the m^{th} order, with constant or quasi-constant coefficients, and of any number of variables. Also, arbitrary constants will then be replaced in v by arbitrary functions of the quasi-constants.

$$\text{III. To integrate } t d_i^2 u + \left(a + \frac{1}{2}\right) d_i u = \frac{1}{4} b^2 u.$$

Let $t = \tau^2$, and change the independent variable from t to τ ;

$$\therefore \tau(d_\tau^2 - b^2)u + a \cdot 2d_\tau u = 0.$$

Operate on this with $(d_\tau^2 - b^2)^{-a}$, (see Ex. 9, Chap. VII.);

$$\begin{aligned} \therefore \tau(d_\tau^2 - b^2)^{1-a}u &= (d_\tau^2 - b^2)^{-a}0 \\ &= (d_\tau - b)^{-a}0 + (d_\tau + b)^{-a}0 \dots (\text{Art. 16}). \end{aligned}$$

As the two parts of u are independent, we may take them separately and add the results together for the complete value of u ;

$$\begin{aligned} \therefore \tau(d_\tau - b)^{1-a}(d_\tau + b)^{1-a}u &= (d_\tau - b)^{-a}0; \\ \therefore \tau \cdot \epsilon^{b\tau} d_\tau^{1-a} \epsilon^{-b\tau} (d_\tau + b)^{1-a}u &= \epsilon^{b\tau} d_\tau^{-a} \epsilon^{-b\tau} 0, \\ &= \epsilon^{b\tau} s_\tau^a 0. \end{aligned}$$

If now we divide by τ , and consider that $s_\tau^a 0$ contains powers of τ from τ^0 to τ^{a-1} , and that all the coefficients are arbitrary, we shall see that although

$$\tau^{-1} s_\tau^a 0 \text{ is not } = s_\tau^{a-1} 0 + \tau^{-1} s_\tau 0,$$

the two members of this inequality are *equivalent* for the purpose before us.

$$\therefore \epsilon^{b\tau} d_\tau^{1-a} \epsilon^{-b\tau} (d_\tau + b)^{1-a} u = \epsilon^{b\tau} s_\tau^{a-1} 0 + \epsilon^{b\tau} \tau^{-1} s_\tau 0,$$

$$\therefore (d_\tau + b)^{1-a} u = \epsilon^{b\tau} d_\tau^{a-1} \tau^{-1} s_\tau 0.$$

The right-hand member, though not equal to $\epsilon^{b\tau} \tau^{-a} s_\tau 0$, is equivalent to it, because $s_\tau 0$ is arbitrary. By actual differentiation, and preserving the arbitrary character of $s_\tau 0$, we have

$$\epsilon^{-b\tau} d_\tau^{1-a} \epsilon^{b\tau} u = \epsilon^{b\tau} \tau^{-a} s_\tau 0;$$

$$\therefore u = A \epsilon^{-b\tau} d_\tau^{a-1} (\epsilon^{2b\tau} \tau^{-a}),$$

A being an arbitrary constant.

The other part of u will be obtained from this by writing $-b$ for b ;

$$\therefore u = A \epsilon^{-b\tau} d_\tau^{a-1} (\epsilon^{2b\tau} \tau^{-a}) + B \epsilon^{b\tau} d_\tau^{a-1} (\epsilon^{-2b\tau} \tau^{-a}).$$

After the differentiations have been performed we are to write $\sqrt{t}$ for τ .

$$\text{IV. To integrate } t d_t^2 u + \left(\frac{1}{2} - a\right) d_t u = \frac{1}{4} b^2 u.$$

Change the variable as in the preceding Article,

$$\therefore \tau (d_\tau^2 - b^2) u = 2a d_\tau u \dots \dots \dots (1).$$

Operate on this with $(d_\tau^2 - b^2)^a$,

$$\therefore \tau (d_\tau^2 - b^2)^{a+1} u = 0;$$

$$\therefore u = (d_\tau - b)^{-a-1} 0 + (d_\tau + b)^{-a-1} 0$$

$$= \epsilon^{b\tau} s_\tau^{a+1} \epsilon^{-b\tau} 0 + \epsilon^{-b\tau} s_\tau^{a+1} \epsilon^{b\tau} 0$$

$$= \epsilon^{b\tau} s_\tau^{a+1} 0 + \epsilon^{-b\tau} s_\tau^{a+1} 0.$$

The two parts of u are independent. Each part contains $a + 1$ arbitrary constants; and, as there ought only to be one in each part, the supernumeraries are to be determined by substitution in equation (1). Taking one of the two parts of

u and writing $\epsilon^{-b\tau} s_r^{a+1} 0$ for u in equation (1), and assuming

$$s_r^{a+1} 0 = B_0 + B_1\tau + B_2\tau^2 + \dots + B_a\tau^a,$$

we find the following to be the law of the coefficients,

$$B_{k+1} = \frac{2b}{k+1} \cdot \frac{a-k}{2a-k} \cdot B_k.$$

If in the other term of u we assume

$$s_r^{a+1} 0 = A_0 + A_1\tau + A_2\tau^2 + \dots + A_a\tau^a,$$

we have merely to write $-b$ for b in the above law;

$$\therefore A_{k+1} = -\frac{2b}{k+1} \cdot \frac{a-k}{2a-k} \cdot A_k;$$

$$\begin{aligned} \therefore u &= \epsilon^{b\tau}(A_0 + A_1\tau + A_2\tau^2 + \dots + A_a\tau^a) \\ &+ \epsilon^{-b\tau}(B_0 + B_1\tau + B_2\tau^2 + \dots + B_a\tau^a); \end{aligned}$$

and thus the integral required is completely determined. It only remains to write in this result $\sqrt{t}$ for τ .

V. In the preceding Chapters we have spoken of the equation $d_t d_x u + cu = 0$ as in general unintegrable in finite terms; and in Art. 27 we have shewn that its integral, if it admit of one in finite terms, cannot contain arbitrary functions. We shall proceed to shew how a complete *first* integral of it can always be obtained; and also that with a certain restriction the *final* integral can be obtained also in finite terms.

For, since

$$d_t d_x u + cu = 0 \dots\dots\dots (1),$$

multiply by t and differentiate with d_t , having first commutated d_t, d_x ;

$$\therefore d_t d_x \cdot t d_t u + c t d_t u + cu = 0,$$

$$\text{or } (d_t d_x + c) \cdot t d_t u + cu = 0.$$

Similarly we obtain

$$(d_t d_x + c) \cdot x d_x u + cu = 0;$$

$$\therefore (d_t d_x + c) (t d_t u - x d_x u) = 0,$$

which being compared with equation (1) gives, by integrating with $(d_t d_x + c)^{-1}$,

$$t d_t u - x d_x u = ku \dots \dots \dots (2).$$

This integral has been obtained in a general manner, and is therefore entitled to be taken as a first integral of the proposed equation. It contains no arbitrary function but only an arbitrary constant.

(1) Let us now eliminate d_x between (1) and (2);

$$\therefore t d_t^2 u + (1 - k) d_t u + c x u = 0 \dots \dots \dots (3).$$

Here we see the proposed equation, which is of two independent variables, reduced to an equivalent equation of one independent variable, for x is here a quasi-constant. Unfortunately this equation is of a form that does not admit of general integration for all possible values of k , k being arbitrary. If, however, we refer to Arts. III. and IV. we shall find it there proved that it does admit of integration in finite terms when k differs from $\frac{1}{2}$ by a positive or negative integer; and that the simplest or root-form of the integral is that which corresponds to $k = \frac{1}{2}$.

Now, by Art. III., if U be the integral for this value of k ,

$$U = A e^{-b r} s_r e^{2 b r} + B e^{b r} s_r e^{-2 b r};$$

and this is equivalent to

$$U \sqrt{x} = A e^{2 \sqrt{-c x t}} + B e^{-2 \sqrt{-c x t}} \dots \dots \dots (4),$$

since $b^2 = -4cx$, and $\tau = \sqrt{t}$. This is only one part of the integral, for in obtaining this equation we treated t as the leading independent variable. Had we taken x instead of t we should have found,

$$U\sqrt{t} = A'\epsilon^{2\sqrt{-cxt}} + B'\epsilon^{-2\sqrt{-cxt}}.$$

Consequently the complete integral, found by taking the sum of the partial values of u , will be

$$U = \left(\frac{A}{\sqrt{x}} + \frac{A'}{\sqrt{t}}\right)\epsilon^{2\sqrt{-cxt}} + \left(\frac{B}{\sqrt{x}} + \frac{B'}{\sqrt{t}}\right)\epsilon^{-2\sqrt{-cxt}} \dots\dots(5).$$

This is the complete finite root-integral of the proposed equation.

We call it the root-integral because the integrals for all other values of k which differ by integers from $\frac{1}{2}$ can be obtained from this by mere integration or differentiation with s_i or d_i .

For U in (4) satisfies the equation (found from (3) by writing $\frac{1}{2}$ for k),

$$td_i^2 U + \frac{1}{2} d_i U + cxU = 0;$$

operate on (3) with $d_i^{k-\frac{1}{2}}$;

$$\therefore td_i^2 \cdot d_i^{k-\frac{1}{2}} u + \frac{1}{2} d_i \cdot d_i^{k-\frac{1}{2}} u + cx \cdot d_i^{k-\frac{1}{2}} u = 0.$$

Comparing these two equations, we see that

$$d_i^{k-\frac{1}{2}} u = U;$$

$$\therefore u = s_i^{k-\frac{1}{2}} U.$$

Thus u can be derived from U exactly in the same way as in all linear equations with constant coefficients. The arbitrary constant k in this case points only to the fact that if U be an integral, then $d_i^n U$ or $s_i^n U$ will be integrals also.

We may therefore take as the root-integral that which is the simplest in form, which is that given in (5) and corresponds to $k = \frac{1}{2}$. From that we may at pleasure derive other forms corresponding to different values of k by integration or differentiation.

(2) We have called equation (2) a *first* integral of the proposed equation, but it is so in a peculiar sense only. It was obtained by first raising the proposed equation an order higher, and then lowering it two orders by integration with the compound symbol $(d_i d_x + c)^{-1}$. This method of obtaining an integral always furnishes a result which requires to be checked and limited by substitution in the proposed or some equivalent equation, since on account of having been obtained from an equation of too high an order it may produce a result which is too general.

(3) Though we cannot for all possible values of the arbitrary constant k obtain an integral of equation (3), the equation (2) admits of general integration, giving

$$ux^k = F(tx) \dots\dots\dots(6).$$

If therefore (2), the equation from which this is derived, be general, this is the form of the integral for all values of k . Nothing remains but to find the form of F for any proposed value of k . If u taken from this be substituted in (1), it produces an equation of the second order of the same form

as (3); so that the root-form of F can only be found when $k = \frac{1}{2}$. We learn however from (6), that the subject of F is the same for all values of k .

(4) We must now draw attention to the circumstance that the proposed equation is known to be satisfied by the following equation,

$$u = Ae^{ht-kx} + Be^{kx-ht} \dots\dots\dots(7),$$

h, k being arbitrary constants subject to the condition

$$hk = c.$$

Now this value of u does not satisfy (2); and consequently (2), which in one direction embraces too much, in another direction does not embrace enough. By raising the proposed equation to the third order we expected to embrace within our results more than belonged to us, and we now find that we have lost some results which ought to be included in a general integral. We shall recur to this subject in Art. XII.

VI. The equation discussed in the preceding Article is a particular case of the more general equation

$$\phi(d_t d_x) u = 0 \dots\dots\dots(1).$$

We shall shew that a first integral of this equation can be obtained.

By formula (2) of Art. 39 we have

$$\phi(d_t d_x) \cdot t d_t u = t \phi(d_t d_x) \cdot d_t u + \phi'(d_t d_x) \cdot d_t d_x u.$$

Now $d_t u$ if written for u in equation (1) will satisfy it,

$$\therefore \phi(d_t d_x) \cdot d_t u = 0;$$

$$\therefore \phi(d_t d_x) \cdot t d_t u = \phi'(d_t d_x) \cdot d_t d_x u.$$

Similarly,

$$\phi(d_t d_x) \cdot x d_x u = \phi'(d_t d_x) \cdot d_t d_x u;$$

$$\therefore \phi(d_t d_x) (t d_t u - x d_x u) = 0:$$

which being compared with equation (1) gives, integrating this with $(\phi \cdot d_t d_x)^{-1}$,

$$t d_t u - x d_x u = k u \dots\dots\dots (2).$$

This is therefore a first integral of the whole class of equations comprehended in equation (1).

The integral of the equation (2) is

$$u = t^k F(tx) \dots\dots\dots (3).$$

And this must be substituted in equation (1); and the resulting equation, if it can be integrated, will give the form of F and any restrictions on the values of k which may be necessary. It teaches us that the *subject* of F must be tx .

VII. There is another interesting equation, viz.

$$d_t d_x u + d_t d_y u + d_x d_y u = 0 \dots\dots\dots (1),$$

of which a first integral may be obtained in a similar way.

By commutation of the symbols of operation it will take the form

$$(d_x + d_y) \cdot d_t u + d_x d_y u = 0.$$

Multiply by t and differentiate with d_t ,

$$\therefore d_t (d_x + d_y) \cdot t d_t u + d_x d_y \cdot t d_t u + d_x d_y u = 0;$$

$$\therefore (d_t d_x + d_t d_y + d_x d_y) \cdot t d_t u + d_x d_y u = 0.$$

Similarly,

$$(d_t d_x + d_t d_y + d_x d_y) \cdot x d_x u + d_t d_y u = 0,$$

and

$$(d_t d_x + d_t d_y + d_x d_y) \cdot y d_y u + d_t d_x u = 0.$$

Adding the last three equations together and taking account of equation (1), we have

$$(d_t d_x + d_t d_y + d_x d_y) (t d_t u + x d_x u + y d_y u) = 0,$$

and integrating this with $(d_t d_x + d_t d_y + d_x d_y)^{-1}$, we have

$$t d_t u + x d_x u + y d_y u = k u \dots\dots\dots (2).$$

This is a first integral of equation (1); and being integrated it gives

$$u = t^k f\left(\frac{x}{t}, \frac{y}{t}\right) \dots\dots\dots (3)$$

as the integral of equation (1). But the form of f will have to be determined by the substitution of this value of u in equation (1). It will depend on the integration of an equation of *two* independent variables.

VIII. As a companion Article to VI. we may shew how to obtain a first integral of another important equation,

$$\phi(D_t \Delta_t) u = 0 \dots\dots\dots (1),$$

in which we suppose $D_t x = a$, and $\Delta_t x = b$; and $\therefore \xi = x - at$ and $\xi' = x - bt$.

By formula (3) of Art. 39 we have (since $\Delta_t \xi = b - a$, and $D_t \xi' = a - b$)

$$\begin{aligned} \phi(D_t \Delta_t) \cdot \xi \Delta_t u &= \xi \cdot \phi(D_t \Delta_t) \Delta_t u + (b - a) D_t \Delta_t u \\ &= (b - a) D_t \Delta_t u. \end{aligned}$$

Similarly,

$$\phi(D_t \Delta_t) \cdot \xi' D_t u = (a-b) D_t \Delta_t u;$$

$$\therefore \phi(D_t \Delta_t) \cdot (\xi \Delta_t u + \xi' D_t u) = 0;$$

$$\therefore \xi \Delta_t u + \xi' D_t u = ku \dots\dots\dots(2).$$

This is the first integral of equation (1). Its integral is

$$u = \xi^{\frac{k}{b-a}} F(\xi \xi'); \text{ or } \xi'^{\frac{k}{a-b}} f(\xi \xi') \dots\dots\dots(3).$$

These two forms of u are mutually equivalent. The form of the arbitrary function F or f , and any necessary restrictions on the values of k , will have to be ascertained by substitution in the proposed equation (1).

IX. If in the equation

$$d_t^2 u = d_x^2 u + cu \dots\dots\dots(1),$$

we change the independent variables by writing $t+x$ for t , and $t-x$ for x , it becomes

$$d_t d_x u = cu \dots\dots\dots(2),$$

the root-integral of which we know. We can therefore pass at once from the root-integral of this equation to the corresponding integral of equation (1) by writing $\frac{1}{2}(t+x)$ for t , and $\frac{1}{2}(t-x)$ for x . Consequently the root-integral of (1) is

$$Vu = A\epsilon^{\sqrt{(t^2-x^2)}c} + B\epsilon^{-\sqrt{(t^2-x^2)}c} \dots\dots\dots(3),$$

V being equal to either of the quantities

$$\sqrt{t+x}, \text{ and } \sqrt{t-x}.$$

Again: from this we may obtain the root-integral of the equation

$$d_t^2 u + d_x^2 u = cu \dots\dots\dots(4),$$

by writing $x\sqrt{-1}$ for x . Consequently the root-integral of (4) is

$$Vu = A\epsilon^{\sqrt{(t^2+x^2)}c} + B\epsilon^{-\sqrt{(t^2+x^2)}c} \dots\dots\dots(5),$$

V being equal to either of the quantities,

$$\sqrt{(t+x\sqrt{-1})}, \text{ and } \sqrt{(t-x\sqrt{-1})}.$$

REMARK. Both in (3) and (5) there are two values of u arising from the double values of V , and we may in each case take the sum of the two values of u for the complete integral.

X. To integrate the equation

$$d_t^2 u = d_x^2 u + d_y^2 u.$$

This may be written in the two forms

$$d_t^2 u = d_x^2 u + cu \dots\dots\dots(1),$$

where c stands for d_y^2 ;

$$d_t^2 u = d_y^2 u + cu \dots\dots\dots(2),$$

where c stands for d_x^2 .

The latter may be deduced from the former by interchanging x and y .

From Art. IX., writing d_y for $\sqrt{c}$ in (3) we obtain

$$V_1 u = A\epsilon^{\sqrt{t^2-x^2}d_y} + B\epsilon^{-\sqrt{t^2-x^2}d_y}.$$

But in the case before us A and B are functions of y , being constants only in reference to d_t and d_x .

$$\text{Now} \quad e^{\sqrt{t^2-x^2}d_y} F(y) = F(y + \sqrt{t^2-x^2}),$$

$$\text{and} \quad e^{-\sqrt{t^2-x^2}d_y} f(y) = f(y - \sqrt{t^2-x^2});$$

$$\therefore V_1 u = F(y + \sqrt{t^2-x^2}) + f(y - \sqrt{t^2-x^2}) \dots (3);$$

V_1 being equal to either of the two quantities,

$$\sqrt{t+x}, \text{ and } \sqrt{t-x}.$$

To obtain the value of u corresponding to the form (2) of the proposed equation, interchange x and y ,

$$\therefore V_2 u = F_1(x + \sqrt{t^2-y^2}) + f_1(x - \sqrt{t^2-y^2}) \dots (4),$$

V_2 being equal to either of the two quantities,

$$\sqrt{t+y}, \text{ and } \sqrt{t-y}.$$

The complete integral will be found by adding together the partial values of u .

Again; from the integral of

$$d_t^2 u = d_x^2 u + d_y^2 u,$$

we may deduce that of Laplace's equation,

$$d_t^2 u + d_x^2 u + d_y^2 u = 0,$$

by writing $x\sqrt{-1}$ for x , and $y\sqrt{-1}$ for y . The constituent parts of its integral may be written in the following forms;

$$V_1 u = F(\sqrt{t^2+x^2} + y\sqrt{-1}) + f(\sqrt{t^2+x^2} - y\sqrt{-1});$$

$$V_2 u = F_1(\sqrt{t^2+y^2} + x\sqrt{-1}) + f_1(\sqrt{t^2+y^2} - x\sqrt{-1}).$$

$$V_1 = (t + x\sqrt{-1})^{\frac{1}{2}}, \text{ or } (t - x\sqrt{-1})^{\frac{1}{2}};$$

$$V_2 = (t + y\sqrt{-1})^{\frac{1}{2}}, \text{ or } (t - y\sqrt{-1})^{\frac{1}{2}}.$$

The same remark respecting the complete value of u may be made in this Article as in the preceding one.

XI. If we have an equation of the form

$$d_t^2 u = d_x^2 u + d_y^2 u + d_z^2 u + \dots + cu \dots \dots \dots (1),$$

the number of independent variables being n , we can reduce it to another of $n-1$ independent variables. And this we can do with $n-1$ independent results.

For brevity use C to represent $d_y^2 + d_z^2 + \dots + c$, and write the equation in the form

$$d_t^2 u - d_x^2 u = Cu \dots \dots \dots (2).$$

If (as in Art. IX.) we write in this equation $t+x$ for t , and $t-x$ for x , it becomes

$$d_t d_x u = Cu \dots \dots \dots (3).$$

In the manner of Art. v. we find from this equation

$$td_t u - xd_x u = ku \dots \dots \dots (4).$$

By eliminating d_x between this and (3), we obtain

$$td_t^2 u + (1-k) d_t u = Cxu \dots \dots \dots (5),$$

in which x is a quasi-constant. Consequently this equation contains only $n-1$ independent variables; and it is equivalent to the proposed equation.

Thus it appears that by the above process we can render any one of the independent variables $x, y, z, \dots$ in equation (1) a quasi-constant; and consequently there are $n - 1$ ways of reducing their number from n to $n - 1$; and there are also $n - 1$ independent equations of the form of equation (4).

Again; if we confine ourselves to the obtaining of the root-integrals of equation (1) we are to assume $k = \frac{1}{2}$ in equation (5); and if we then write τ^2 for $4tx$ this equation will take the form

$$d_{\tau}^2 u = Cu,$$

or
$$d_{\tau}^2 u = d_y^2 u + d_x^2 u + \dots + cu;$$

which form is the same as that of the proposed equation deprived of one of its terms.

Thus by reversing the process we can pass from the form

$$d_i^2 u = cu, \text{ to } d_i^2 u = d_x^2 u + cu;$$

and from the last to

$$d_i^2 u = d_x^2 u + d_y^2 u + cu;$$

and so on to a similar equation of any number of independent variables.

XII. The following equation is interesting as throwing some light on the nature of the peculiar first integral obtained in some of the preceding Articles, and on which indeed the success obtained has been entirely dependent. The equation referred to is

$$d_i d_x u + d_y d_x u = 0 \dots\dots\dots(1).$$

By proceeding with this equation as in Art. v. we easily obtain the four following equations:

$$(d_t d_x + d_y d_z) \cdot t d_t u + d_y d_z u = 0 \dots \dots \dots (2),$$

$$(d_t d_x + d_y d_z) \cdot x d_x u + d_y d_z u = 0 \dots \dots \dots (3),$$

$$(d_t d_x + d_y d_z) \cdot y d_y u + d_t d_z u = 0 \dots \dots \dots (4),$$

$$(d_t d_x + d_y d_z) \cdot z d_z u + d_t d_x u = 0 \dots \dots \dots (5).$$

Taking account of equation (1) we deduce from these the following results:

$$(2) - (3) \text{ gives } (d_t d_x + d_y d_z) (t d_t u - x d_x u) = 0,$$

$$(4) - (5) \text{ gives } (d_t d_x + d_y d_z) (y d_y u - z d_z u) = 0,$$

$$(2) + (4) \text{ gives } (d_t d_x + d_y d_z) (t d_t u + y d_y u) = 0,$$

$$(2) + (5) \text{ gives } (d_t d_x + d_y d_z) (t d_t u + z d_z u) = 0,$$

$$(3) + (4) \text{ gives } (d_t d_x + d_y d_z) (x d_x u + y d_y u) = 0,$$

$$(3) + (5) \text{ gives } (d_t d_x + d_y d_z) (x d_x u + z d_z u) = 0.$$

It will be seen on examination that of the last six equations only *three* are independent. Integrating them with $(d_t d_x + d_y d_z)^{-1}$, we obtain the following six first integrals of which only three are independent:

$$t d_t u - x d_x u = k_1 u, \text{ and } y d_y u - z d_z u = k_2 u \dots \dots (6),$$

$$t d_t u + y d_y u = l_1 u, \text{ and } x d_x u + z d_z u = l_2 u \dots \dots (7),$$

$$t d_t u + z d_z u = m_1 u, \text{ and } x d_x u + y d_y u = m_2 u \dots \dots (8).$$

The integrals of these are (taking them two and two as they stand),

$$u = t^k y^k F(tx, yz);$$

$$u = t^l x^l f\left(\frac{y}{t}, \frac{z}{x}\right);$$

$$u = t^m x^m \phi\left(\frac{z}{t}, \frac{y}{x}\right).$$

The forms of F, f, ϕ are to be determined by the separate substitution of these values of u in equation (1).

In this way, for example, we obtain from the first of these the following finite integral of the proposed equation,

$$Vu = F(\sqrt{tx} + \sqrt{-yz}) + f(\sqrt{tx} - \sqrt{-yz}),$$

the quantity represented by V being any one of these four following:

$$\sqrt{ty}, \quad \sqrt{tz}, \quad \sqrt{xy}, \quad \sqrt{xz}.$$

The Theory of the Peculiar First Integrals.

(1) On turning to the three sets of first integrals (6), (7), (8), we perceive at once that no two of the sets can be satisfied with the same value of u . From this it follows that they are not to be used simultaneously. They point to three independent integrals of the proposed equation.

(2) On looking at the equations from which the *first* integrals were obtained we learn that the symbols

$$td_t - xd_x, \quad yd_y - zd_z, \quad xd_x + zd_z, \quad \&c.$$

are commutative with $d_t d_x + d_y d_z$ on u .

(3) To avoid an unnecessary complexity of symbols let us confine ourselves to the consideration of the peculiar first integral in connexion with the equation

$$d_t d_x u + cu = 0 \dots\dots\dots(9).$$

In this case $(d_t d_x + c)$ is commutative with $(td_t - xd_x)$ on u .

Let H represent any simple or compound symbol of operation which is thus commutative with $(d_t d_x + c)$ on u ,

$$\therefore (d_t d_x + c) H . u = 0.$$

Integrating this equation with $(d_t d_x + c)^{-1}$ we have the following first integral,

$$H . u = ku \dots\dots\dots(10).$$

If then we can find a form of H which will satisfy the one condition contained in its definition, viz. that it is commutative with $(d_t d_x + c)$ on u , the above first integral will hold good; and between (9) and (10) the final integral is to be obtained. Thus there will be as many first integrals as we can find independent forms for H ; and as many final integrals as can be formed out of them.

(4) That H may have several values seems to be rendered probable by the fact that in the case of the equation we are now considering we already know two, viz.

$$H = d_t + md_x, \text{ and } H = td_t - xd_x \dots\dots\dots(11),$$

for both of these are of the first order and commutative with $(d_t d_x + c)$ on u .

The latter has been already considered; and the former gives, by the elimination of d_x from $d_t d_x u + cu = 0$, the following:

$$d_t^2 u - kd_t u - mcu = 0,$$

the integral of which is of the form

$$u = e^{at} F(x) + e^{\beta t} f(x).$$

And as t and x enter similarly into the proposed equation, this may be written in the form

$$u = A e^{at - \alpha' x} + B e^{\beta t - \beta' x},$$

in which $\alpha x' = c$, and $\beta \beta' = c$. The two terms may be both included in the form

$$u = A e^{at - \alpha' x} + B e^{\alpha' x - at}.$$

Thus the existence of this final integral is accounted for by the fact that $d_t + m d_x$ is commutative with $(d_t d_x + c)$.

(5) We have found two different and apparently independent forms of H , and from them two corresponding apparently independent integrals of the proposed equation. If these integrals are really independent, and even if the two forms of H are really independent, great uncertainty would exist as to the generality of the integral of

$$d_t d_x u + cu = 0$$

found in Art. v. We shall therefore now shew that the form of H given by the equation

$$H = t d_t - x d_x \dots \dots \dots (12)$$

includes the other form of H which we have just discussed.

We have not considered it necessary to remark respecting the equation

$$d_t d_x u + cu = 0,$$

and the other equations derived from it and discussed in Arts. IX, X., that as the independent variables t , x enter

this equation only in the form dt, dx , we may write $t + c_1, x + c_2$ instead of t and x . This property is general for all independent variables which enter an equation only in the form of differentials. We ought therefore to have written in the integrals derived from those equations $t + c_1, x + c_2, y + c_3, \dots$ for $t, x, y, \dots$. We abstained from doing so for simplicity of symbolic expression.

Now if in (12) we write $at + l, bx + m$ for t and x , we have

$$H = \left(t + \frac{l}{a}\right) d_t - \left(x + \frac{m}{b}\right) d_x \dots \dots \dots (13),$$

and we observe that this form of H includes *both* the forms of H before given; and that H is precisely the same as if we had merely written

$$t + \frac{l}{a} \text{ for } t, \text{ and } x + \frac{m}{b} \text{ for } x.$$

We have therefore to make these changes in the integral of $d_t d_x u + cu = 0$ given in Art. v., and we shall then have the integral in a form that includes both forms of H . It is therefore not correct to say that H has *two* independent forms. The form discussed in section (4) of this Article is *necessarily* included in the form $td_t - xd_x$ by the general property that the general value of t is $t + c_1$, and that of x is $x + c_2$; in other words, the forms in which the independent variables enter into the proposed differential equations are such that we are at liberty arbitrarily to change the origins of the independent variables in the integrals.

(6) But can we reverse the process and shew that the integral

$$u = A e^{at - a'x} + B e^{a'x - at}$$

is included in the general integral found in Art. v. the general form of which is

$$u \sqrt{t} = A\epsilon^{\sqrt{-clx}} + B\epsilon^{-\sqrt{-clx}},$$

by simply changing the origins of t and x ? It can be done as follows.

For t and x write $t + \frac{l}{a}$ and $x + \frac{m}{a}$ in this integral; and for brevity let us confine ourselves to the first term of the integral, as the result for the other term can be deduced from it;

$$\therefore u \sqrt{t + \frac{l}{a}} = A\epsilon^{\left(t + \frac{l}{a}\right)^{\frac{1}{2}} \left(x + \frac{m}{a}\right)^{\frac{1}{2}} \sqrt{-c}};$$

$$\therefore u \sqrt{at + l} = A \sqrt{a} \cdot \epsilon^{\frac{\sqrt{-clm}}{a}} \cdot \epsilon^{\sqrt{-clm} \left(\frac{t}{2l} + \frac{x}{2m} + aW\right)};$$

wherein aW represents the remainder of the series obtained by the extraction of the root of

$$\left(1 + \frac{at}{l}\right) \left(1 + \frac{ax}{m}\right).$$

It is evident we may suppose the factors

$$\sqrt{a}, \quad \epsilon^{\frac{\sqrt{-clm}}{a}},$$

which are independent of t and x , absorbed into the arbitrary constant, for which we may therefore write $A \sqrt{l}$;

$$\therefore u \sqrt{at + l} = A \sqrt{l} \cdot \epsilon^{\sqrt{-clm} \left(\frac{t}{2l} + \frac{x}{2m} + aW\right)}.$$

Now as an extreme case let us suppose $a = 0$;

$$\therefore u = A\epsilon^{\sqrt{-clm} \cdot \left(\frac{t}{2l} + \frac{x}{2m}\right)}.$$

The generality of this form is not diminished by the supposition that $lm + 1 = 0$; and as we are at liberty to assume that l is negative and m positive, or conversely, that l is positive and m negative, we have the two following results corresponding to the two terms of the general integral,

$$u = A\epsilon^{(t\sqrt{\frac{m}{l}} - x\sqrt{\frac{l}{m}})\sqrt{\frac{c}{2}}} + B\epsilon^{(x\sqrt{\frac{l}{m}} - t\sqrt{\frac{m}{l}})\sqrt{\frac{c}{2}}}.$$

But it will be objected to this result that it has been obtained by assuming $t + \infty$ and $x + \infty$ for t and x ; and that these assumptions are not lawful. We allow the illegality of such assumptions; and our reply is that these are not the assumptions we have made. If the reader will look back he will see that we assumed $t = at + l$, and $x = ax + m$; and that by these we were directed to write $t + \frac{l}{a}$ for t , and $x + \frac{m}{a}$ for x in the integral. We have done this, and it has conducted us to the above result.

We feel ourselves therefore entitled to affirm that the general value of H is $td_t - xd_x$; and that the general finite integral of the equation $d_t d_x u + cu = 0$ is that given in Art. v., with the understanding that we are at liberty to arbitrarily change the origins of the independent variables in that integral.

SEPTEMBER, 1869.

*A CATALOGUE of EDUCATIONAL BOOKS,
with a Short Account of their
Character and Aim,*

Published by

MACMILLAN AND CO.

Bedford Street, Covent Garden, London.

CLASSICAL.

ÆSCHYLI EUMENIDES. The Greek Text, with English Notes and English Verse, Translation, and an Introduction. By BERNARD DRAKE, M.A., late Fellow of King's College, Cambridge. 8vo. 3s. 6d.

The Greek text adopted in this Edition is based upon that of Wellauer, which may be said, in general terms, to represent that of the best manuscripts. But in correcting the Text, and in the Notes, advantage has been taken of the suggestions of Hermann, Paley, Linwood, and other commentators. In the Translation, the simple character of the Æschylean dialogues has generally enabled the author to render them without any material deviation from the construction and idioms of the original Greek.

ARISTOTLE ON FALLACIES; OR, THE SOPHISTICAL
ELENCHI. With a Translation and Notes by EDWARD POSTE,
M.A., Fellow of Oriel College, Oxford. 8vo. 8s. 6d.

Besides the doctrine of Fallacies, Aristotle offers, either in this treatise or in other passages quoted in the commentary, various glances over the world of science and opinion, various suggestions or problems which are still agitated, and a vivid picture of the ancient system of dialectics, which it is hoped may be found both interesting and instructive.

Aristotle. — AN INTRODUCTION TO ARISTOTLE'S
RHETORIC. With Analysis, Notes, and Appendices. By E.
M. COPE, Senior Fellow and Tutor of Trinity College, Cam-
bridge. 8vo. 14s.

This work is introductory to an edition of the Greek Text of Aristotle's Rhetoric, which is in course of preparation. Its object is to render that treatise thoroughly intelligible. The author has aimed to illustrate, as preparatory to the detailed explanation of the work, the general bearings and relations of the Art of Rhetoric in itself, as well as the special mode of treating it adopted by Aristotle in his peculiar system. The evidence upon obscure or doubtful questions connected with the subject is examined; and the relations which Rhetoric bears, in Aristotle's view, to the kindred art of Logic are fully considered. A connected Analysis of the work is given, sometimes in the form of paraphrase; and a few important matters are separately discussed in Appendices. There is added, as a general Appendix, by way of specimen of the antagonistic system of Isocrates and others, a complete analysis of the treatise called Ῥητορικὴ πρὸς Ἀλέξανδρον, with a discussion of its authorship and of the probable results of its teaching.

CATULLI VERONENSIS LIBER. Edited by R. ELLIS, Fellow
of Trinity College, Oxford. 18mo. 3s. 6d.

An elaborate Preface deals with the arrangement of the verse, and discusses various readings. "Rarely," says the Saturday Review, "have we read a classic author with so reliable, acute, and safe a guide."

Cicero.—THE SECOND PHILIPPIC ORATION. With an Introduction and Notes, translated from the German of KARL HALM. Edited, with Corrections and Additions, by JOHN E. B. MAYOR, M.A., Fellow and Classical Lecturer of St. John's College, Cambridge. Third Edition, revised. Fcap. 8vo. 5s.

This volume opens with a List of Books useful to the Student of Cicero, including History, Chronology, Lexicons, and some account of various editions, mostly German, of the works of Cicero. The Introduction is based on Halm: where Halm gives a reference to a classic, the passage has been commonly printed at length; where the reference is to Halm's notes on other Ciceronian speeches, or to modern books, the additional matter has been incorporated: and the numerous Greek quotations have been rendered into English. The English editor has further illustrated the work by additions drawn, for the most part, (1) from the ancient authorities; (2) from his own private marginal references, and from collections; (3) from the notes of previous commentators. A copious 'argument' is also given.

DEMOSTHENES ON THE CROWN. The Greek Text with English Notes. By B. DRAKE, M.A., late Fellow of King's College, Cambridge. Third Edition, to which is prefixed **ÆSCHINES AGAINST CTESIPHON**, with English Notes. Fcap. 8vo. 5s.

An Introduction discusses the immediate causes of the two orations, and their general character. The Notes contain frequent references to the best authorities. Among the appendices at the end is a chronological table of the life and public career of Æschines and Demosthenes.

Hodgson.—MYTHOLOGY FOR LATIN VERSIFICATION. A brief Sketch of the Fables of the Ancients, prepared to be rendered into Latin Verse for Schools. By F. HODGSON, B.D., late Provost of Eton. New Edition, revised by F. C. HODGSON, M.A. 18mo. 3s.

The late Provost of Eton has here supplied a help to the composition of Latin Verse, combined with a brief introduction to Classical Mythology.

In this new edition a few mistakes have been rectified ; rules have been added to the Prosody ; and a more uniform system has been adopted with regard to the help afforded.

JUVENAL FOR SCHOOLS. With English Notes. By J. E. B. MAYOR, M.A. New and cheaper Edition. Crown 8vo.

[*In the Press.*]

Marshall.—A TABLE OF IRREGULAR GREEK VERBS, classified according to the arrangement of Curtius' Greek Grammar. By J. M. MARSHALL, M.A., Fellow and late Lecturer of Brasenose College, Oxford ; one of the Masters in Clifton College. 8vo. cloth. 1s.

The system of this table has been borrowed from the excellent Greek Grammar of Dr. Curtius.

Mayor, John E. B.—FIRST GREEK READER. Edited after KARL HALM, with Corrections and large Additions by JOHN E. B. MAYOR, M.A. Fellow and Classical Lecturer of St. John's College, Cambridge. Fcap. 8vo. 6s.

A selection of short passages, serving to illustrate especially the Greek Accidence. A good deal of syntax is incidentally taught, and Madvig and other books are cited, for the use of masters : but no learner is expected to know more of syntax than is contained in the Notes and Vocabulary. A preface "To the Reader," not only explains the aim and method of the volume, but also deals with classical instruction generally. The extracts are uniformly in the Attic dialect, and any Hellenistic forms occurring in the original classic authors, such as Ælian and Polybius, have been discarded in favour of the corresponding Attic expressions. This book may be used in connexion with Mayor's "Greek for Beginners."

Mayor, Joseph B.—GREEK FOR BEGINNERS. By the Rev. J. B. MAYOR, M.A. Fcap. 8vo. 4s. 6d.

The distinctive method of this book consists in building up a boy's knowledge of Greek upon the foundation of his knowledge of English and Latin, instead of trusting everything to the unassisted memory. The forms and constructions of Greek have been thoroughly compared with

those of Latin, and no Greek words have been used in the earlier part of the book except such as have connexions either in English or Latin. Each step leads naturally on to its successor, grammatical forms and rules are at once applied in a series of graduated exercises, accompanied by ample vocabularies. Thus the book serves as Grammar, Exercise book, and Vocabulary. Where possible, the Grammar has been simplified; the ordinary ten declensions are reduced to three, which correspond to the first three in Latin; and the system of stems is adopted. A general Vocabulary, and Index of Greek words, completes the work.

Plato.—THE REPUBLIC OF PLATO. Translated into English, with an Analysis and Notes, by J. LL. DAVIES, M.A., and D. J. VAUGHAN, M.A. Third Edition, with Vignette Portraits of Plato and Socrates, engraved by JEENS from an Antique Gem. 18mo. 4s. 6d.

An introductory notice supplies some account of the life of Plato, and the translation is preceded by an elaborate analysis. "The translators have," in the judgment of the Saturday Review, "produced a book which any reader, whether acquainted with the original or not, can peruse with pleasure as well as profit."

Plautus (Ramsay).—THE MOSTELLARIA OF PLAUTUS. With Notes Critical and Explanatory, Prolegomena, and Excursus. By WILLIAM RAMSAY, M.A., formerly Professor of Humanity in the University of Glasgow. Edited by Professor GEORGE G. RAMSAY, M.A., of the University of Glasgow. 8vo. 14s.

"The fruits of that exhaustive research and that ripe and well-digested scholarship which its author brought to bear upon everything that he undertook are visible throughout it. It is furnished with a complete apparatus of prolegomena, notes, and excursus; and for the use of veteran scholars it probably leaves nothing to be desired."—PALL MALL GAZETTE.

Roby.—A LATIN GRAMMAR for the Higher Classes in Grammar Schools. By H. J. ROBY, M.A., based on the Elementary Latin Grammar. [In the Press.]

Sallust.—CAII SALLUSTII CRISPI CATILINA ET JUGURTHA. For Use in Schools. With copious Notes. By C. MERIVALE, B.D. (In the present Edition the Notes have been carefully revised, and a few remarks and explanations added.) Second Edition. Fcap. 8vo. 4s. 6d.

The JUGURTHA and the CATILINA may be had separately, price 2s. 6d. each.

This edition of Sallust, prepared by the distinguished historian of Rome, contains an introduction, concerning the Life and works of Sallust, lists of the Consuls, and elaborate notes.

Tacitus.—THE HISTORY OF TACITUS TRANSLATED INTO ENGLISH. By A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Notes and a Map. 8vo. 10s. 6d.

The translators have endeavoured to adhere as closely to the original as was thought consistent with a proper observance of English idiom. At the same time, it has been their aim to reproduce the precise expressions of the author. The campaign of Civilis is elucidated in a note of some length, which is illustrated by a map, containing the names of places and of tribes occurring in the work. There is also a complete account of the Roman army as it was constituted in the time of Tacitus. This work is characterised by the Spectator as "a scholarly and faithful translation."

THE AGRICOLA AND GERMANIA OF TACITUS. A Revised Text, English Notes, and Maps. By ALFRED J. CHURCH, M.A., and W. J. BRODRIBB, M.A. Fcap. 8vo. 3s. 6d.

"We have endeavoured, with the aid of recent editions, thoroughly to elucidate the text, explaining the various difficulties, critical and grammatical, which occur to the student. We have consulted throughout, besides the older commentators, the editions of Ritter and Orelli, but we are under special obligations to the labours of the recent German editors, Wex, and Kritz." Two Indexes are appended, (1) of Proper Names, (2) of Words and Phrases explained.

THE AGRICOLA and GERMANIA may be had separately, price 2s. each.

Tacitus—continued.

THE AGRICOLA AND GERMANIA. Translated into English by A. J. CHURCH, M.A., and W. J. BRODRIBB, M.A. With Maps and Notes. Extra fcap. 8vo. 2s. 6d.

The translators have sought to produce such a version as may satisfy scholars who demand a faithful rendering of the original, and English readers who are offended by the baldness and frigidity which commonly disfigure translations. The treatises are accompanied by introductions, notes, maps, and a chronological summary. The Athenæum says of this work that it is "a version at once readable and exact, which may be perused with pleasure by all, and consulted with advantage by the classical student."

Thring.—A LATIN GRADUAL. A First Latin Construing Book for Beginners. By EDWARD THRING, M.A. Fcap. 8vo. 2s. 6d.

The Head Master of Uppingham has here sought to supply by easy steps a knowledge of grammar, combined with a good Vocabulary. Passages have been selected from the best Latin authors in prose and verse. These passages are gradually built up in their grammatical structure, and finally printed in full. A short practical manual of common mood constructions, with their English equivalents, forms a second part.

A MANUAL OF MOOD CONSTRUCTIONS. Fcap. 8vo. 1s. 6d.

Treats of the ordinary mood constructions, as found in the Latin, Greek, and English languages.

A CONSTRUING BOOK. Fcap. 8vo. 2s. 6d.

Thucydides.—THE SICILIAN EXPEDITION. Being Books VI. and VII. of Thucydides, with Notes. A New Edition, revised and enlarged, with a Map. By the Rev. PERCIVAL FROST, M.A., late Fellow of St. John's College, Cambridge. Fcap. 8vo. 5s.

This edition is mainly a grammatical one. Attention is called to the force of compound verbs, and the exact meaning of the various tenses employed.

Wright.—Works by T. WRIGHT, M.A., late Head Master of Sutton Coldfield School.

HELLENICA ; OR, A HISTORY OF GREECE IN GREEK, as related by Diodorus and Thucydides ; being a First Greek Reading Book, with explanatory Notes, Critical and Historical. Third Edition, with a Vocabulary. 12mo. 3s. 6d.

In the last twenty chapters of this volume, Thucydides sketches the rise and progress of the Athenian Empire in so clear a style and in such simple language, that the editor has doubts whether any easier or more instructive passages can be selected for the use of the pupil who is commencing Greek. This book includes a chronological table of the events recorded.

A HELP TO LATIN GRAMMAR ; or, The Form and Use of Words in Latin, with Progressive Exercises. Crown 8vo. 4s. 6d.

This book is not intended as a rival to any of the excellent Grammars now in use ; but as a help to enable the beginner to understand them.

THE SEVEN KINGS OF ROME. An Easy Narrative, abridged from the First Book of Livy by the omission of Difficult Passages ; being a First Latin Reading Book, with Grammatical Notes. With Vocabulary and Exercises. Second Edition. 5s.

This work is intended to supply the pupil with an easy construing book, which may at the same time be made the vehicle for instructing him in the rules of grammar and principles of composition. The notes profess to teach what is commonly taught in grammars. It is conceived that the pupil will learn the rules of construction of the language much more easily from separate examples, which are pointed out to him in the course of his reading, and which he may himself set down in his note-book after some scheme of his own, than from a heap of quotations amassed for him by others.

Or, separately,

SEVEN KINGS OF ROME. 3s.

VOCABULARY AND EXERCISES TO "THE SEVEN KINGS." 2s. 6d.

CLASSIC VERSIONS OF ENGLISH BOOKS, AND LATIN HYMNS.

THE following works are, as the heading indicates, classic renderings of English books. For scholars, and particularly for writers of Latin Verse, the series has a special value. The *Hymni Ecclesiæ* are here inserted, as partly falling under the same class.

Church and Kebbel.—HORÆ TENNYSONIANÆ. Curantibus A. J. CHURCH, A.M., T. E. KEBBEL, A.M. [*In the Press.*]

Latham.—SERTUM SHAKSPERIANUM, Subnexis aliquot aliunde excerptis floribus. Latine reddidit Rev. H. LATHAM, M.A. Extra fcap. 8vo. 5s.

Besides versions of Shakspeare this volume contains, among other pieces, Gray's "Elegy," Campbell's "Hohenlinden," Wolfe's "Burial of Sir John Moore," and selections from Cowper and George Herbert.

Lyttelton.—THE COMUS OF MILTON, rendered into Greek Verse. By LORD LYTTTELTON. Extra fcap. 8vo. 6s. 6d.

THE SAMSON AGONISTES OF MILTON, rendered into Greek Verse. By LORD LYTTTELTON. Extra fcap. 8vo. 6s. 6d.

Merivale.—KEATS' HYPERION, rendered into Latin Verse. By C. MERIVALE, B.D. Second Edit. Extra fcap. 8vo. 3s. 6d.

Hymni Ecclesiæ.—Edited by Rev. DR. NEWMAN. Extra fcap. 8vo. 7s. 6d.

Hymns of the Mediæval Church. The first Part contains selections from the Parisian Breviary; the second from those of Rome, Salisbury, and York.

Trench (Archbishop).—SACRED LATIN POETRY, chiefly Lyrical, selected and arranged for Use; with Notes and Introduction. Fcap. 8vo. 7s.

In this work the editor has selected hymns of a catholic religious sentiment that are common to Christendom, while rejecting those of a distinctively Romish character.

MATHEMATICS.

Airy.—Works by G. B. AIRY, Astronomer Royal :—

ELEMENTARY TREATISE ON PARTIAL DIFFERENTIAL EQUATIONS. Designed for the Use of Students in the Universities. With Diagrams. Crown 8vo. cloth. 5s. 6d.

It is hoped that the methods of solution here explained, and the instances exhibited, will be found sufficient for application to nearly all the important problems of Physical Science, which require for their complete investigation the aid of Partial Differential Equations.

ON THE ALGEBRAICAL AND NUMERICAL THEORY OF ERRORS OF OBSERVATIONS AND THE COMBINATION OF OBSERVATIONS. Crown 8vo. cloth. 6s. 6d.

In order to spare astronomers and observers in natural philosophy the confusion and loss of time which are produced by referring to the ordinary treatises embracing both branches of probabilities (the first relating to chances which can be altered only by the changes of entire units or integral multiples of units in the fundamental conditions of the problem ; the other concerning those chances which have respect to insensible gradations in the value of the element measured) the present tract has been drawn up. It relates only to errors of observation, and to the rules, derivable from the consideration of these errors, for the combination of the results of observations.

Airy, G. B.—continued.

UNDULATORY THEORY OF OPTICS. Designed for the Use of Students in the University. New Edition. Crown 8vo. cloth. 6s. 6d.

The undulatory theory of optics is presented to the reader as having the same claims to his attention as the theory of gravitation: namely, that it is certainly true, and that, by mathematical operations of general elegance, it leads to results of great interest. This theory explains with accuracy a vast variety of phenomena of the most complicated kind. The plan of this tract has been to include those phenomena only which admit of calculation, and the investigations are applied only to phenomena which actually have been observed.

ON SOUND AND ATMOSPHERIC VIBRATIONS. With the Mathematical Elements of Music. Designed for the Use of Students of the University. Crown 8vo. 9s.

This volume consists of sections, which again are divided into numbered articles, on the following topics: General recognition of the air as the medium which conveys sound; Properties of the air on which the formation and transmission of sound depend; Theory of undulations as applied to sound, &c.; Investigation of the motion of a wave of air through the atmosphere; Transmission of waves of soniferous vibrations through different gases, solids, and fluids; Experiments on the velocity of sound, &c.; On musical sounds, and the manner of producing them; On the elements of musical harmony and melody, and of simple musical composition; On instrumental music; On the human organs of speech and hearing.

Bayma.—THE ELEMENTS OF MOLECULAR MECHANICS. By JOSEPH BAYMA, S.J., Professor of Philosophy, Stonyhurst College. Demy 8vo. cloth. 10s. 6d.

Of the twelve Books into which the present treatise is divided, the first and second give the demonstration of the principles which bear directly on the constitution and the properties of matter. The next three books contain a series of theorems and of problems on the laws of motion of elementary substances. In the sixth and seventh, the mechanical constitution of mole-

cules is investigated and determined: and by it the general properties of bodies are explained. The eighth book treats of luminiferous æther. The ninth explains some special properties of bodies. The tenth and eleventh contain a radical and lengthy investigation of chemical principles and relations, which may lead to practical results of high importance. The twelfth and last book treats of molecular masses, distances, and powers.

Beasley.—AN ELEMENTARY TREATISE ON PLANE TRIGONOMETRY. With Examples. By R. D. BEASLEY, M.A., Head Master of Grantham Grammar School. Second Edition, revised and enlarged. Crown 8vo. cloth. 3s. 6d.

This treatise is specially intended for use in schools. The choice of matter has been chiefly guided by the requirements of the three days' examination at Cambridge. About four hundred examples have been added to this edition, mainly collected from the examination papers of the last ten years.

Boole.—Works by G. BOOLE, D.C.L., F.R.S., Professor of Mathematics in the Queen's University, Ireland.

A TREATISE ON DIFFERENTIAL EQUATIONS. New and Revised Edition. Edited by I. TODHUNTER. Crown 8vo. cloth. 14s.

Professor Boole has endeavoured in this treatise to convey as complete an account of the present state of knowledge on the subject of Differential Equations, as was consistent with the idea of a work intended, primarily, for elementary instruction. The earlier sections of each chapter contain that kind of matter which has usually been thought suitable for the beginner, while the latter ones are devoted either to an account of recent discovery, or the discussion of such deeper questions of principle as are likely to present themselves to the reflective student in connexion with the methods and processes of his previous course.

A TREATISE ON DIFFERENTIAL EQUATIONS. Supplementary Volume. Edited by I. TODHUNTER. Crown 8vo. cloth. 8s. 6d.

This volume contains all that Professor Boole wrote for the purpose of enlarging his treatise on Differential Equations.

Boole, G., D.C.L.—continued.

THE CALCULUS OF FINITE DIFFERENCES. Crown 8vo.
cloth. 10s. 6d.

In this exposition of the Calculus of Finite Differences, particular attention has been paid to the connexion of its methods with those of the Differential Calculus—a connexion which in some instances involves far more than a merely formal analogy. The work is in some measure designed as a sequel to Professor Boole's Treatise on Differential Equations.

**CAMBRIDGE SENATE-HOUSE PROBLEMS AND RIDERS,
WITH SOLUTIONS:—**

1848-1851.—PROBLEMS. By FERRERS and JACKSON. 8vo.
cloth. 15s. 6d.

1848-1851.—RIDERS. By JAMESON. 8vo. cloth. 7s. 6d.

1854.—PROBLEMS AND RIDERS. By WALTON and
MACKENZIE. 8vo. cloth. 10s. 6d.

1857.—PROBLEMS AND RIDERS. By CAMPION and
WALTON. 8vo. cloth. 8s. 6d.

1860.—PROBLEMS AND RIDERS. By WATSON and ROUTH.
Crown 8vo. cloth. 7s. 6d.

1864.—PROBLEMS AND RIDERS. By WALTON and WIL-
KINSON. 8vo. cloth. 10s. 6d.

These volumes will be found of great value to Teachers and Students, as indicating the style and range of mathematical study in the University of Cambridge.

**CAMBRIDGE COURSE OF ELEMENTARY NATURAL
PHILOSOPHY,** for the Degree of B.A. Originally compiled by
J. C. SNOWBALL, M.A., late Fellow of St. John's College.
Fifth Edition, revised and enlarged, and adapted for the Middle-
Class Examinations by THOMAS LUND, B.D., Late Fellow and
Lecturer of St. John's College, Editor of Wood's Algebra, &c.
Crown 8vo. cloth. 5s.

This work will be found adapted to the wants, not only of University Students, but also of many others who require a short course of Mechanics

and Hydrostatics, and especially of the candidates at our Middle Class Examinations. At the end of each chapter a series of easy questions is added for the exercise of the student.

CAMBRIDGE AND DUBLIN MATHEMATICAL JOURNAL.

The Complete Work, in Nine Vols. 8vo. cloth, 7l. 4s.

Only a few copies remain on hand. Among Contributors to this work will be found Sir W. Thomson, Stokes, Adams, Boole, Sir W. R. Hamilton, De Morgan, Cayley, Sylvester, Jellett, and other distinguished mathematicians.

Candler.—HELP TO ARITHMETIC. Designed for the use of Schools. By H. CANDLER, M.A. Mathematical Master of Uppingham School. Extra fcap. 8vo. 2s. 6d.

This work is intended as a companion to any text book that may be in use.

Cheyne.—AN ELEMENTARY TREATISE ON THE PLANETARY THEORY. With a Collection of Problems. By C. H. H. CHEYNE, B.A. Crown 8vo. cloth. 6s. 6d.

In this volume, an attempt has been made to produce a treatise on the Planetary theory, which, being elementary in character, should be so far complete, as to contain all that is usually required by students in the University of Cambridge.

THE EARTH'S MOTION OF ROTATION. By C. H. H. CHEYNE, M.A. Crown 8vo. 3s. 6d.

The first part of this work consists of an application of the method of the variation of elements to the general problem of rotation. In the second part the general rotation formulæ are applied to the particular case of the earth.

Childe.—THE SINGULAR PROPERTIES OF THE ELLIPSOID AND ASSOCIATED SURFACES OF THE NTH DEGREE. By the Rev. G. F. CHILDE, M.A., Author of "Ray Surfaces," "Related Caustics," &c. 8vo. 10s. 6d.

The object of this volume is to develop peculiarities in the Ellipsoid; and, further, to establish analogous properties in the unlimited congeneric series of which this remarkable surface is a constituent.

Christie.—A COLLECTION OF ELEMENTARY TEST-QUESTIONS IN PURE AND MIXED MATHEMATICS; with Answers and Appendices on Synthetic Division, and on the Solution of Numerical Equations by Horner's Method. By JAMES R. CHRISTIE, F.R.S., late First Mathematical Master at the Royal Military Academy, Woolwich. Crown 8vo. cloth. 8s. 6d.

The series of Mathematical exercises here offered to the public is collected from those which the author has, from time to time, proposed for solution by his pupils during a long career at the Royal Military Academy. A student who finds that he is able to solve the larger portion of these exercises, may consider that he is thoroughly well grounded in the elementary principles of pure and mixed Mathematics.

Dalton.—ARITHMETICAL EXAMPLES. Progressively arranged, with Exercises and Examination Papers. By the Rev. T. DALTON, M.A., Assistant Master of Eton College. 18mo. cloth. 2s. 6d.

Answers to the Examples are appended.

Day.—PROPERTIES OF CONIC SECTIONS PROVED GEOMETRICALLY. PART I., THE ELLIPSE, with Problems. By the Rev. H. G. DAY, M.A., Head Master of Sedburgh Grammar School. Crown 8vo. 3s. 6d.

The object of this book is the introduction of a treatment of Conic Sections which should be simple and natural, and lead by an easy transition to the analytical methods, without departing from the strict geometry of Euclid.

Dodgson.—AN ELEMENTARY TREATISE ON DETERMINANTS, with their Application to Simultaneous Linear Equations and Algebraical Geometry. By CHARLES L. DODGSON, M.A., Student and Mathematical Lecturer of Christ Church, Oxford. Small 4to. cloth. 10s. 6d.

The object of the author is to present the subject as a continuous chain of argument, separated from all accessories of explanation or illustration. All such explanation and illustration as seemed necessary for a beginner, are introduced either in the form of foot-notes, or, where that would have occupied too much room, of Appendices.

Drew.—GEOMETRICAL TREATISE ON CONIC SECTIONS. By W. H. DREW, M.A., St. John's College, Cambridge. Third Edition. Crown 8vo. cloth. 4s. 6d.

In this work the subject of Conic Sections has been placed before the student in such a form that, it is hoped, after mastering the elements of Euclid, he may find it an easy and interesting continuation of his geometrical studies. With a view, also, of rendering the work a complete manual of what is required at the Universities, there have either been embodied into the text or inserted among the examples, every book-work question, problem, and rider, which has been proposed in the Cambridge examinations up to the present time.

SOLUTIONS TO THE PROBLEMS IN DREW'S CONIC SECTIONS. Crown 8vo. cloth. 4s. 6d.

Ferrers.—AN ELEMENTARY TREATISE ON TRILINEAR CO-ORDINATES, the Method of Reciprocal Polars, and the Theory of Projectors. By the Rev. N. M. FERRERS, M.A., Fellow and Tutor of Gonville and Caius College, Cambridge. Second Edition. Crown 8vo. 6s. 6d.

The object of the author in writing on this subject has mainly been to place it on a basis altogether independent of the ordinary Cartesian system, instead of regarding it as only a special form of Abridged Notation. A short chapter on Determinants has been introduced.

Frost.—THE FIRST THREE SECTIONS OF NEWTON'S PRINCIPIA. With Notes and Illustrations. Also a collection of Problems, principally intended as Examples of Newton's Methods. By PERCIVAL FROST, M.A., late Fellow of St. John's College, Mathematical Lecturer of King's College, Cambridge. Second Edition. 8vo. cloth. 10s. 6d.

The author's principal intention is to explain difficulties which may be encountered by the student on first reading the Principia, and to illustrate the advantages of a careful study of the methods employed by Newton, by showing the extent to which they may be applied in the solution of problems; he has also endeavoured to give assistance to the student who is engaged in the study of the higher branches of mathematics, by representing in a geometrical form several of the processes employed in the Differential and Integral Calculus, and in the analytical investigations of Dynamics.

Frost and Wolstenholme.—A TREATISE ON SOLID GEOMETRY. By PERCIVAL FROST, M.A., and the Rev. J. WOLSTENHOLME, M.A., Fellow and Assistant Tutor of Christ's College. 8vo. cloth. 18s.

The authors have endeavoured to present before students as comprehensive a view of the subject as possible. Intending to make the subject accessible, at least in the earlier portion, to all classes of students, they have endeavoured to explain completely all the processes which are most useful in dealing with ordinary theorems and problems, thus directing the student to the selection of methods which are best adapted to the exigencies of each problem. In the more difficult portions of the subject, they have considered themselves to be addressing a higher class of students; and they have there tried to lay a good foundation on which to build, if any reader should wish to pursue the science beyond the limits to which the work extends.

Godfray.—A TREATISE ON ASTRONOMY, for the Use of Colleges and Schools. By HUGH GODFRAY, M.A., Mathematical Lecturer at Pembroke College, Cambridge. 8vo. cloth. 12s. 6d.

This book embraces all those branches of Astronomy which have, from time to time, been recommended by the Cambridge Board of Mathematical

Godfray, Hugh—*continued.*

Studies: but by far the larger and easier portion, adapted to the first three days of the Examination for Honours, may be read by the more advanced pupils in many of our schools. The author's aim has been to convey clear and distinct ideas of the celestial phenomena.

AN ELEMENTARY TREATISE ON THE LUNAR THEORY,
with a Brief Sketch of the Problem up to the time of Newton.
By HUGH GODFRAY, M.A. Second Edition, revised. Crown
8vo. cloth. 5s. 6d.

These pages will, it is hoped, form an introduction to more recondite works. Difficulties have been discussed at considerable length. The selection of the method followed with regard to analytical solutions, which is the same as that of Airy, Herschel, &c. was made on account of its simplicity; it is, moreover, the method which has obtained in the University of Cambridge.

Hemming.—AN ELEMENTARY TREATISE ON THE
DIFFERENTIAL AND INTEGRAL CALCULUS, for the
Use of Colleges and Schools. By G. W. HEMMING, M.A.,
Fellow of St. John's College, Cambridge. Second Edition, with
Corrections and Additions. 8vo. cloth. 9s.

Jones and Cheyne.—ALGEBRAICAL EXERCISES. Pro-
gressively arranged. By the Rev. C. A. JONES, M.A., and C. H.
CHEYNE, M.A., Mathematical Masters of Westminster School.
New Edition. 18mo. cloth. 2s. 6d.

This little book is intended to meet a difficulty which is probably felt more or less by all engaged in teaching Algebra to beginners. It is, that while new ideas are being acquired, old ones are forgotten. In the belief that constant practice is the only remedy for this, the present series of miscellaneous exercises has been prepared. Their peculiarity consists in this, that though miscellaneous they are yet progressive, and may be used by

the pupil almost from the commencement of his studies. They are not intended to supersede the systematically arranged examples to be found in ordinary treatises on Algebra, but rather to supplement them.

The book being intended chiefly for Schools and Junior Students, the higher parts of Algebra have not been included.

Kitchener.—A GEOMETRICAL NOTE-BOOK, containing Easy Problems in Geometrical Drawing preparatory to the Study of Geometry. For the Use of Schools. By F. E. KITCHENER, M.A., Mathematical Master at Rugby. 4to. 2s.

It is the object of this book to make some way in overcoming the difficulties of Geometrical conception, before the mind is called to the attack of Geometrical theorems. A few simple methods of construction are given; and space is left on each page, in order that the learner may draw in the figures.

Morgan.—A COLLECTION OF PROBLEMS AND EXAMPLES IN MATHEMATICS. With Answers. By H. A. MORGAN, M.A., Sadlerian and Mathematical Lecturer of Jesus College, Cambridge. Crown 8vo. cloth. 6s. 6d.

This book contains a number of problems, chiefly elementary, in the Mathematical subjects usually read at Cambridge. They have been selected from the papers set during late years at Jesus College. Very few of them are to be met with in other collections, and by far the larger number are due to some of the most distinguished Mathematicians in the University.

Parkinson.—Works by S. PARKINSON, B.D., Fellow and Prælector of St. John's College, Cambridge.

AN ELEMENTARY TREATISE ON MECHANICS. For the Use of the Junior Classes at the University and the Higher Classes in Schools. With a Collection of Examples. Fourth edition, revised. Crown 8vo. cloth. 9s. 6d.

In preparing a fourth edition of this work the author has kept the same object in view as he had in the former editions—namely, to include in it such portions of Theoretical Mechanics as can be conveniently investigated

Parkinson, S.—continued.

without the use of the Differential Calculus, and so render it suitable as a manual for the junior classes in the University and the higher classes in Schools. With one or two short exceptions, the student is not presumed to require a knowledge of any branches of Mathematics beyond the elements of Algebra, Geometry, and Trigonometry. Several additional propositions have been incorporated in the work for the purpose of rendering it more complete; and the collection of Examples and Problems has been largely increased.

A TREATISE ON OPTICS. Second Edition, revised. Crown 8vo. cloth. 10s. 6d.

A collection of examples and problems has been appended to this work, which are sufficiently numerous and varied in character to afford useful exercise for the student. For the greater part of them, recourse has been had to the Examination Papers set in the University and the several Colleges during the last twenty years.

Phear.—ELEMENTARY HYDROSTATICS. With Numerous Examples. By J. B. PHEAR, M.A., Fellow and late Assistant Tutor of Clare College, Cambridge. Fourth Edition. Crown 8vo. cloth. 5s. 6d.

This edition has been carefully revised throughout, and many new illustrations and examples added, which it is hoped will increase its usefulness to students at the Universities and in Schools. In accordance with suggestions from many engaged in tuition, answers to all the Examples have been given at the end of the book.

Pratt.—A TREATISE ON ATTRACTIONS, LAPLACE'S FUNCTIONS, AND THE FIGURE OF THE EARTH. By JOHN H. PRATT, M.A., Archdeacon of Calcutta, Author of "The Mathematical Principles of Mechanical Philosophy." Third Edition. Crown 8vo. cloth. 6s. 6d.

The author's chief design in this treatise is to give an answer to the question, "Has the Earth acquired its present form from being originally in a fluid state?" This Edition is a complete revision of the former ones.

Puckle.—AN ELEMENTARY TREATISE ON CONIC SECTIONS AND ALGEBRAIC GEOMETRY. With Numerous Examples and Hints for their Solution; especially designed for the Use of Beginners. By G. H. PUCKLE, M.A., Head Master of Windermere College. Third Edition, revised and enlarged. Crown 8vo. cloth. 7s. 6d.

This work has been completely rewritten, and a large amount of new matter has been added to suit the requirements of the present time. The author has written with special reference to those difficulties and misapprehensions which are found most common to beginners. The treatise is complete as far as regards Cartesian Co-ordinates.

Rawlinson.—ELEMENTARY STATICS, by the Rev. GEORGE RAWLINSON, M.A. Edited by the Rev. EDWARD STURGES, M.A., of Emmanuel College, Cambridge, and late Professor of the Applied Sciences, Elphinstone College, Bombay. Crown 8vo. cloth. 4s. 6d.

Published under the authority of Her Majesty's Secretary of State for India, for use in the Government Schools and Colleges in India.

Reynolds.—MODERN METHODS IN ELEMENTARY GEOMETRY. By E. M. REYNOLDS, M.A., Mathematical Master in Clifton College. Crown 8vo. 3s. 6d.

Geometry has received extensive developments in modern times, but in England there has been no corresponding improvement in elementary teaching.

Routh.—AN ELEMENTARY TREATISE ON THE DYNAMICS OF THE SYSTEM OF RIGID BODIES. With Numerous Examples. By EDWARD JOHN ROUTH, M.A., late Fellow and Assistant Tutor of St. Peter's College, Cambridge; Examiner in the University of London. Second Edition, enlarged. Crown 8vo. cloth. 14s.

In this edition the author has made several additions to each chapter. He has tried, even at the risk of some little repetition, to make each

chapter, as far as possible, complete in itself, so that all that relates to any one part of the subject may be found in the same place. This arrangement will enable every student to select his own order in which to read the subject. The Examples which will be found at the end of each chapter have been chiefly selected from the Examination Papers which have been set in the University and the Colleges in the last few years.

Smith (Barnard).—Works by BARNARD SMITH, M.A.,
Rector of Glaston, Rutlandshire, late Fellow and Senior Bursar
of St. Peter's College, Cambridge.

ARITHMETIC AND ALGEBRA, in their Principles and Application; with numerous systematically arranged Examples taken from the Cambridge Examination Papers, with especial reference to the Ordinary Examination for the B.A. Degree. Tenth Edition. Crown 8vo. cloth. 10s. 6d.

This manual is now extensively used in Schools and Colleges, both in England and in the Colonies. It has also been found of great service for students preparing for the Middle Class and Civil and Military Service Examinations, from the care that has been taken to elucidate the principles of all the rules. The present edition has been carefully revised. "To all those whose minds are sufficiently developed to comprehend the simplest mathematical reasoning, and who have not yet thoroughly mastered the principles of Arithmetic and Algebra, it is calculated to be of great advantage."—ATHENÆUM.

Of this work, also, one of the highest possible authorities, the late Dean Peacock, writes: "Mr. Smith's work is a most useful publication. The rules are stated with great clearness. The examples are well selected, and worked out with just sufficient detail, without being encumbered by too minute explanations; and there prevails throughout it that just proportion of theory and practice, which is the crowning excellence of an elementary work."

Smith, Barnard—continued.

ARITHMETIC FOR SCHOOLS. New Edition. Crown 8vo. cloth. 4s. 6d.

Adapted from the author's work on "Arithmetic and Algebra," by the omission of the algebraic portion, and by the introduction of new exercises. The reason of each arithmetical process is fully exhibited. The system of Decimal Coinage is explained; and answers to the exercises are appended at the end. This Arithmetic is characterised as "admirably adapted for instruction, combining just sufficient theory with a large and well-selected collection of exercises for practice."—JOURNAL OF EDUCATION.

COMPANION TO ARITHMETIC FOR SCHOOLS.

[Preparing.

A KEY TO THE ARITHMETIC FOR SCHOOLS. Seventh Edition. Crown 8vo. cloth. 8s. 6d.

EXERCISES IN ARITHMETIC. With Answers. Crown 8vo. limp cloth. 2s. 6d.

Or sold separately, Part I. 1s. ; Part II. 1s. ; Answers, 6d.

These Exercises have been published in order to give the pupil examples in every rule of Arithmetic. The greater number have been carefully compiled from the latest University and School Examination Papers.

SCHOOL CLASS-BOOK OF ARITHMETIC. 18mo. cloth. 3s.

Or sold separately, Parts I. and II. 10d. each ; Part III. 1s.

This manual, published at the request of many schoolmasters, and chiefly intended for National and Elementary Schools, has been prepared on the same plan as that adopted in the author's School Arithmetic, which is in extensive circulation in England and abroad. The Metrical Tables have been introduced, from the conviction on the part of the author, that the knowledge of such tables, and the mode of applying them, will be of great use to the rising generation.

KEYS TO SCHOOL CLASS-BOOK OF ARITHMETIC. Complete in one volume, 18mo. cloth, 6s. 6d. ; or Parts I. II. and III. 2s. 6d. each.

Smith, Barnard—*continued.*

SHILLING BOOK OF ARITHMETIC FOR NATIONAL AND
ELEMENTARY SCHOOLS. 18mo. cloth. Or separately,
Part I. 2d.; Part II. 3d.; Part III. 7d. Answers, 6d.

THE SAME, with Answers complete. 18mo. cloth. 1s. 6d.

This Shilling Book of Arithmetic has been prepared for the use of National and other schools at the urgent request of numerous masters of schools both at home and abroad. The Explanations of the Rules, and the Examples will, it is hoped, be found suited to the most elementary classes.

KEY TO SHILLING BOOK OF ARITHMETIC. 18mo. cloth.
4s. 6d.

EXAMINATION PAPERS IN ARITHMETIC. 18mo. cloth.
1s. 6d. The same, with Answers, 18mo. 1s. 9d.

The object of these Examination Papers is to test students both in the theory and practice of Arithmetic. It is hoped that the method adopted will lead students to deduce results from general principles rather than to apply stated rules. The author believes that the practice of giving examples under particular rules makes the working of Arithmetic quite mechanical, and tends to throw all but very clever boys off their balance when a general paper on the subject is put before them.

KEY TO EXAMINATION PAPERS IN ARITHMETIC.
18mo. cloth. 4s. 6d.

Smith (J. H.).—Works by J. H. SMITH, M.A., Gonville and
Caius College, Cambridge.

A TREATISE ON ELEMENTARY STATICS. Second Edition.
8vo. 5s. 6d.

This treatise is intended to give a simple explanation of the part of Statics required in the previous examination and the second examination for ordinary degrees in the University of Cambridge. The examples have been

Smith, J. H.—continued.

selected from papers set in University Examinations. In this edition some explanations of the elementary definitions have been added; and a new and simple proof of the first part of the "Parallelogram of Forces," a new introduction to the Principle of Moments, and a short account of Friction have been inserted.

A TREATISE ON ELEMENTARY HYDROSTATICS. 8vo. 4s. 6d.

An explanation of the subject of Elementary Hydrostatics as required in the general examination for ordinary degrees at Cambridge. The Examples given at the end of the book are taken from Papers set in the University Examinations.

A TREATISE ON ELEMENTARY TRIGONOMETRY. 8vo. 5s.

This work treats of the elementary parts of Trigonometry, as required of the Candidates for Honours in the previous examination.

A TREATISE ON ELEMENTARY ALGEBRA. For the Use of Colleges and Schools. Crown 8vo. cloth. 6s. 6d.

The object of this treatise is to explain those parts of Algebra which precede the Binomial Theorem. The Author has carefully abstained from making extracts from works on the subject in common use. He has written at considerable length on the subject of Factors and Fractions, because those branches of the subject present great difficulties to beginners. The rule has been never to evade a difficulty. The Examples are progressive, easy, and to a great extent original. Some have been selected from the Senate House Papers of the last three years, and a few from old English, French, or German works.

Snowball.—THE ELEMENTS OF PLANE AND SPHERICAL TRIGONOMETRY; with the Construction and Use of Tables of Logarithms. By J. C. SNOWBALL, M.A. Tenth Edition. Crown 8vo. cloth. 7s. 6d.

In preparing the present edition for the press, the text has been subjected to a careful revision; the proofs of some of the more impor-

tant propositions have been rendered more strict and general; and a considerable addition of more than two hundred examples, taken principally from the questions set of late years in the public examinations of the University and of individual Colleges, has been made to the collection of Examples and Problems for practice.

Tait and Steele.—DYNAMICS OF A PARTICLE. With numerous Examples. By Professor TAIT and Mr. STEELE. New Edition. Crown 8vo. cloth. 10s. 6d.

In this treatise will be found all the ordinary propositions, connected with the Dynamics of Particles, which can be conveniently deduced without the use of D'Alembert's Principle. Throughout the book will be found a number of illustrative examples introduced in the text, and for the most part completely worked out; others with occasional solutions or hints to assist the student are appended to each chapter. For by far the greater portion of these, the Cambridge Senate-House and College Examination Papers have been applied to.

Taylor.—GEOMETRICAL CONICS; including Anharmonic Ratio and Projection, with numerous Examples. By C. TAYLOR, B.A., Scholar of St. John's College, Cambridge. Crown 8vo. cloth. 7s. 6d.

This work contains elementary proofs of the principal properties of Conic Sections, together with chapters on Projection and Anharmonic Ratio.

Tebay.—ELEMENTARY MENSURATION FOR SCHOOLS. With numerous Examples. By SEPTIMUS TEBAY, B.A., Head Master of Queen Elizabeth's Grammar School, Rivington. Extra fcap. 8vo. 3s. 6d.

The object of the present work is to enable boys to acquire a moderate knowledge of Mensuration in a reasonable time. All difficult and useless matter has been avoided. The examples for the most part are easy, and the rules are concise.

Todhunter.—Works by I. TODHUNTER, M.A., F.R.S.,
of St. John's College, Cambridge.

THE ELEMENTS OF EUCLID. For the Use of Colleges and
Schools. New Edition. 18mo. cloth. 3s. 6d.

As the elements of Euclid are usually placed in the hands of young students, it is important to exhibit the work in such a form as will assist them in overcoming the difficulties which they experience on their first introduction to processes of continuous argument. No method appears to be so useful as that of breaking up the demonstrations into their constituent parts; a plan strongly recommended by Professor De Morgan. In the present Edition each distinct assertion in the argument begins a new line: and at the ends of the lines are placed the necessary references to the preceding principles on which the assertions depend. The longer propositions are distributed into subordinate parts, which are distinguished by breaks at the beginning of the lines. Notes, appendix, and a collection of exercises are added.

MENSURATION FOR BEGINNERS. With Numerous Examples.
18mo. cloth. 2s. 6d.

The subjects included in the present work are those which have usually found a place in Elementary Treatises on Mensuration. The mode of treatment has been determined by the fact that the work is intended for the use of beginners. Accordingly it is divided into short independent chapters, which are followed by appropriate examples. A knowledge of the elements of Arithmetic is all that is assumed; and in connection with most of the Rules of Mensuration it has been found practicable to give such explanations and illustrations as will supply the place of formal mathematical demonstrations, which would have been unsuitable to the character of the work.

ALGEBRA FOR BEGINNERS. With numerous Examples. New
Edition. 18mo. cloth. 2s. 6d.

Great pains have been taken to render this work intelligible to young students, by the use of simple language and by copious explanations. In

Todhunter, I.—continued.

determining the subjects to be included and the space to be assigned to each, the Author has been guided by the papers given at the various examinations in elementary Algebra which are now carried on in this country. The book may be said to consist of three parts. The first part contains the elementary operations in integral and fractional expressions; the second the solution of equations and problems; the third treats of various subjects which are introduced but rarely into examination papers, and are more briefly discussed. Provision has at the same time been made for the introduction of easy equations and problems at an early stage—for those who prefer such a course.

KEY TO ALGEBRA FOR BEGINNERS. Crown 8vo. cloth.
6s. 6d.

TRIGONOMETRY FOR BEGINNERS. With numerous Examples.
New Edition. 18mo. cloth. 2s. 6d.

Intended to serve as an introduction to the larger treatise on Plane Trigonometry, published by the Author. The same plan has been adopted as in the Algebra for Beginners: the subject is discussed in short chapters, and a collection of examples is attached to each chapter. The first fourteen chapters present the geometrical part of Plane Trigonometry; and contain all that is necessary for practical purposes. The range of matter included is such as seems required by the various examinations in elementary Trigonometry which are now carried on in the country. Answers are appended at the end.

MECHANICS FOR BEGINNERS. With numerous Examples.
18mo. cloth. 4s. 6d.

Intended as a companion to the two preceding books. The work forms an elementary treatise on demonstrative mechanics. It may be true that this part of mixed mathematics has been sometimes made too abstract and speculative; but it can hardly be doubted that a knowledge of the elements at least of the theory of the subject is extremely valuable even for those who are mainly concerned with practical results. The Author has accordingly endeavoured to provide a suitable introduction to the study of applied

Todhunter, I.—continued.

as well as of theoretical mechanics. The work consists of two parts, namely, Statics and Dynamics. It will be found to contain all that is usually comprised in elementary treatises on Mechanics, together with some additions.

ALGEBRA. For the Use of Colleges and Schools. Fourth Edition. Crown 8vo. cloth. 7s. 6d.

This work contains all the propositions which are usually included in elementary treatises on Algebra, and a large number of Examples for Exercise. The author has sought to render the work easily intelligible to students, without impairing the accuracy of the demonstrations, or contracting the limits of the subject. The Examples, about Sixteen hundred and fifty in number, have been selected with a view to illustrate every part of the subject. Each chapter is complete in itself; and the work will be found peculiarly adapted to the wants of students who are without the aid of a teacher. The Answers to the examples, with hints for the solution of some in which assistance may be needed, are given at the end of the book.

AN ELEMENTARY TREATISE ON THE THEORY OF EQUATIONS. Second Edition, revised. Crown 8vo. cloth. 7s. 6d.

This treatise contains all the propositions which are usually included in elementary treatises on the theory of Equations, together with Examples for exercise. These have been selected from the College and University Examination Papers, and the results have been given when it appeared necessary. In order to exhibit a comprehensive view of the subject, the treatise includes investigations which are not found in all the preceding elementary treatises, and also some investigations which are not to be found in any of them. For the second edition the work has been revised and some additions have been made, the most important being an account of the researches of Professor Sylvester respecting Newton's Rule.

PLANE TRIGONOMETRY. For Schools and Colleges. Third Edition. Crown 8vo. cloth. 5s.

The design of this work has been to render the subject intelligible to beginners, and at the same time to afford the student the opportunity of

Todhunter, I.—continued.

obtaining all the information which he will require on this branch of Mathematics. Each chapter is followed by a set of Examples: those which are entitled Miscellaneous Examples, together with a few in some of the other sets, may be advantageously reserved by the student for exercise after he has made some progress in the subject. In the Second Edition the hints for the solution of the Examples have been considerably increased.

A TREATISE ON SPHERICAL TRIGONOMETRY. Second Edition, enlarged. Crown 8vo. cloth. 4s. 6d.

The present work is constructed on the same plan as the treatise on Plane Trigonometry, to which it is intended as a sequel. In the account of Napier's Rules of Circular Parts, an explanation has been given of a method of proof devised by Napier, which seems to have been overlooked by most modern writers on the subject. Considerable labour has been bestowed on the text in order to render it comprehensive and accurate, and the Examples (selected chiefly from College Examination Papers) have all been carefully verified.

PLANE CO-ORDINATE GEOMETRY, as applied to the Straight Line and the Conic Sections. With numerous Examples. Fourth Edition, revised and enlarged. Crown 8vo. cloth. 7s. 6d.

The Author has here endeavoured to exhibit the subject in a simple manner for the benefit of beginners, and at the same time to include in one volume all that students usually require. In addition, therefore, to the propositions which have always appeared in such treatises, he has introduced the methods of abridged notation, which are of more recent origin; these methods, which are of a less elementary character than the rest of the work, are placed in separate chapters, and may be omitted by the student at first.

A TREATISE ON THE DIFFERENTIAL CALCULUS. With numerous Examples. Fourth Edition. Crown 8vo. cloth. 10s. 6d.

The Author has endeavoured in the present work to exhibit a comprehensive view of the Differential Calculus on the method of limits. In the more elementary portions he has entered into considerable detail in the

Todhunter, I.—continued.

explanations, with the hope that a reader who is without the assistance of a tutor may be enabled to acquire a competent acquaintance with the subject. The method adopted is that of Differential Coefficients. To the different chapters are appended examples sufficiently numerous to render another book unnecessary; these examples being mostly selected from College Examination Papers.

A TREATISE ON THE INTEGRAL CALCULUS AND ITS APPLICATIONS. With numerous Examples. Third Edition, revised and enlarged. Crown 8vo. cloth. 10s. 6d.

This is designed as a work at once elementary and complete, adapted for the use of beginners, and sufficient for the wants of advanced students. In the selection of the propositions, and in the mode of establishing them, it has been sought to exhibit the principles clearly, and to illustrate all their most important results. The process of summation has been repeatedly brought forward, with the view of securing the attention of the student to the notions which form the true foundation of the Calculus itself, as well as of its most valuable applications. Every attempt has been made to explain those difficulties which usually perplex beginners, especially with reference to the limits of integrations. A new method has been adopted in regard to the transformation of multiple integrals. The last chapter deals with the Calculus of Variations. A large collection of exercises, selected from College Examination Papers, has been appended to the several chapters.

EXAMPLES OF ANALYTICAL GEOMETRY OF THREE DIMENSIONS. Second Edition, revised. Crown 8vo. cloth. 4s.

A TREATISE ON ANALYTICAL STATICS. With numerous Examples. Third Edition, revised and enlarged. Crown 8vo. cloth. 10s. 6d.

In this work on statics (treating of the laws of the equilibrium of bodies) will be found all the propositions which usually appear in treatises on Theoretical Statics. To the different chapters examples are appended, which have been principally selected from University Examination Papers.

In the Third Edition many additions have been made, in order to illustrate the application of the principles of the subject to the solution of problems.

Wilson, J. M. — **ELEMENTARY GEOMETRY.** Angles, Parallels, Triangles, and Equivalent Figures, the Circle and Proportion. By J. M. WILSON, M.A., Fellow of St. John's College, Cambridge, and Mathematical Master in Rugby School. Extra fcap. 8vo. 3s. 6d.

"It is an actual substitute for the first two books of Euclid, in which many of his propositions are drawn out from the conception of straightness, parallelism, angles, with wonderful ease and simplicity, and the methods employed have the great merit of suggesting a ready application to the solution of fresh problems."—GUARDIAN.

ELEMENTARY GEOMETRY. PART II. (separately). The Circle and Proportion. By J. M. WILSON, M.A. Extra fcap. 8vo. 2s. 6d.

Wilson (W. P.) — **A TREATISE ON DYNAMICS.** By W. P. WILSON, M.A., Fellow of St. John's College, Cambridge, and Professor of Mathematics in Queen's College, Belfast. 8vo. 9s. 6d.

Wolstenholme. — **A BOOK OF MATHEMATICAL PROBLEMS,** on Subjects included in the Cambridge Course. By JOSEPH WOLSTENHOLME, Fellow of Christ's College, sometime Fellow of St. John's College, and lately Lecturer in Mathematics at Christ's College. Crown 8vo. cloth. 8s. 6d.

CONTENTS:—*Geometry (Euclid)—Algebra—Plane Trigonometry—Geometrical Conic Sections—Analytical Conic Sections—Theory of Equations—Differential Calculus—Integral Calculus—Solid Geometry—Statics—Elementary Dynamics—Newton—Dynamics of a Point—Dynamics of a Rigid Body—Hydrostatics—Geometrical Optics—Spherical Trigonometry and Plane Astronomy.*

SCIENCE..

THE importance of Science as an element of sound education is now generally acknowledged; and accordingly it is obtaining a prominent place in the ordinary course of school instruction. It is the intention of the Publishers to produce a complete series of Scientific Manuals, affording full and accurate elementary information, conveyed in clear and lucid English. The authors are well known as among the foremost men of their several departments; and their names form a ready guarantee for the high character of the books. Subjoined is a list of those manuals that have already appeared, with a short account of each. Others are in active preparation; and the whole will constitute a standard series specially adapted to the requirements of beginners, whether for private study or for school instruction.

ASTRONOMY, by the Astronomer Royal.

POPULAR ASTRONOMY. With Illustrations. By G. B. AIRY, Astronomer Royal. Sixth and cheaper Edition. 18mo. cloth. 4s. 6d.

This work consists of six lectures, which are intended "to explain to intelligent persons the principles on which the instruments of an Observatory are constructed (omitting all details, so far as they are merely sub-

sidary), and the principles on which the observations made with these instruments are treated for deduction of the distances and weights of the bodies of the Solar System, and of a few stars, omitting all minutiae of formulae, and all troublesome details of calculation." The speciality of this volume is the direct reference of every step to the Observatory, and the full description of the methods and instruments of observation.

ASTRONOMY.

MR. LOCKYER'S ELEMENTARY LESSONS IN ASTRONOMY. With Coloured Diagram of the Spectra of the Sun, Stars, and Nebulae, and numerous Illustrations. By J. NORMAN LOCKYER, F.R.A.S. Third Thousand. 18mo. 5s. 6d.

The author has here aimed to give a connected view of the whole subject, and to supply facts, and ideas founded on the facts, to serve as a basis for subsequent study and discussion. The chapters treat of the Stars and Nebulae; the Sun; the Solar System; Apparent Movements of the Heavenly Bodies; the Measurement of Time; Light; the Telescope and Spectroscope; Apparent Places of the Heavenly Bodies; the Real Distances and Dimensions; Universal Gravitation. The most recent astronomical discoveries are incorporated. Mr. Lockyer's work supplements that of the Astronomer Royal mentioned in the previous article.

PHYSIOLOGY.

PROFESSOR HUXLEY'S LESSONS IN ELEMENTARY PHYSIOLOGY. With numerous Illustrations. By T. H. HUXLEY, F.R.S. Professor of Natural History in the Royal School of Mines. Eighth Thousand. 18mo. cloth. 4s. 6d.

This book describes and explains, in a series of graduated lessons, the principles of Human Physiology; or the Structure and Functions of the Human Body. The first lesson supplies a general view of the subject. This is followed by sections on the Vascular or Veinous System, and the Circulation; the Blood and the Lymph; Respiration; Sources of Loss and of Gain to the Blood; the Function of Alimentation; Motion and Locomotion; Sensations and Sensory Organs; the Organ of Sight; the Coalescence of Sensations with one another and with other States of Consciousness; the Nervous System and Innervation; Histology, or the Minute Structure of the Tissues. A Table of Anatomical and Physiological Constants is appended. The lessons are fully illustrated by numerous engravings. The manual is primarily intended to serve as a text-book for teachers and learners in boys' and girls' schools.

QUESTIONS ON HUXLEY'S PHYSIOLOGY FOR SCHOOLS.

By T. ALCOCK, M.D. 18mo. 1s. 6d.

These Questions were drawn up as aids to the instruction of a class of young people in Physiology.

BOTANY.

PROFESSOR OLIVER'S LESSONS IN ELEMENTARY BOTANY. With nearly Two Hundred Illustrations, By DANIEL OLIVER, F.R.S., F.L.S. Sixth Thousand. 18mo. cloth, 4s. 6d.

This book is designed to teach the Elements of Botany on Professor Henslow's plan of selected Types and by the use of Schedules. The earlier chapters, embracing the elements of Structural and Physiological Botany, introduce us to the methodical study of the Ordinal Types. The concluding chapters are entitled, "How to dry Plants" and "How to describe Plants." A valuable Glossary is appended to the volume. In the preparation of this work free use has been made of the manuscript materials of the late Professor Henslow.

CHEMISTRY.

PROFESSOR ROSCOE'S LESSONS IN ELEMENTARY CHEMISTRY, INORGANIC AND ORGANIC. By HENRY E. ROSCOE, F.R.S., Professor of Chemistry in Owens College, Manchester. With numerous Illustrations and Chromo-Litho. of the Solar Spectrum, and of the Alkalies and Alkaline Earths, *New Edition.* Thirteenth Thousand. 18mo. cloth. 4s. 6d.

It has been the endeavour of the author to arrange the most important facts and principles of Modern Chemistry in a plain but concise and scientific form, suited to the present requirements of elementary instruction. For the purpose of facilitating the attainment of exactitude in the knowledge of the subject, a series of exercises and questions upon the lessons have been added. The metric system of weights and measures, and the centigrade thermometric scale, are used throughout the work. The new Edition, besides new wood-cuts, contains many additions and improvements, and includes the most important of the latest discoveries.

MISCELLANEOUS.

ATLAS OF EUROPE. GLOBE EDITION. Uniform in size with Macmillan's Globe Series, containing 45 Coloured Maps, on a uniform scale and projection : with Plans of London and Paris, and a copious Index. Strongly bound in half-morocco, with flexible back, 9s.

This Atlas includes all the countries of Europe in a series of 48 Maps drawn on the same scale, with an Alphabetical Index to the situation of more than ten thousand places ; and the relation of the various maps and countries to each other is defined in a general Key-map. The identity of scale in all the maps facilitates the comparison of extent and distance, and conveys a just impression of the magnitude of different countries. The size suffices to show the provincial divisions, the railways and main roads, the principal rivers and mountain ranges. "This atlas," writes the British Quarterly, "will be an invaluable boon for the school, the desk, or the traveller's portmanteau."

Bates & Lockyer.—A CLASS-BOOK OF GEOGRAPHY.

Adapted to the recent Programme of the Royal Geographical Society. By H. W. BATES, Assistant Secretary to the Royal Geographical Society, and J. N. LOCKYER, F.R.A.S.

[*In the Press.*]

CAMEOS FROM ENGLISH HISTORY. From Rollo to Edward

II. By the Author of "The Heir of Redclyffe." Extra fcap. 8vo. Second Edition. 5s.

The endeavour has not been to chronicle facts, but to put together a series of pictures of persons and events, so as to arrest the attention, and give some individuality and distinctness to the recollection, by gathering together details at the most memorable moments. The "Cameos" are intended as a book for young people just beyond the elementary histories of England,

and able to enter in some degree into the real spirit of events, and to be struck with characters and scenes presented in some relief. "Instead of dry details," says the Nonconformist, "we have living pictures, faithful, vivid, and striking."

HISTORICAL SELECTIONS. Readings from the best Authorities on English and European History. Selected and Arranged by E. M. SEWELL and C. M. YONGE. Crown 8vo. 6s.

When young children have acquired the outlines of History from abridgments and catechisms, and it becomes desirable to give a more enlarged view of the subject, in order to render it really useful and interesting, a difficulty often arises as to the choice of books. Two courses are open, either to take a general and consequently dry history of facts, such as Russel's *Modern Europe*, or to choose some work treating of a particular period or subject, such as the works of Macaulay and Froude. The former course usually renders history uninteresting; the latter is unsatisfactory, because it is not sufficiently comprehensive. To remedy this difficulty selections, continuous and chronological, have, in the present volume, been taken from the larger works of Freeman, Milman, Palgrave, and others, which may serve as distinct landmarks of historical reading. "We know of scarcely anything," says the Guardian, of this volume, "which is so likely to raise to a higher level the average standard of English education."

Hole.—A GENEALOGICAL STEMMA OF THE KINGS OF ENGLAND AND FRANCE. By the Rev. C. HOLE. On Sheet. 1s.

The different families are printed in distinguishing colours, thus facilitating reference.

A BRIEF BIOGRAPHICAL DICTIONARY. Compiled and Arranged by CHARLES HOLE, M.A., Trinity College, Cambridge. Second Edition, 18mo. neatly and strongly bound in cloth. 4s. 6d.

The most comprehensive Biographical Dictionary in the world, containing more than 18,000 names of persons of all countries, with dates of birth and death, and what they were distinguished for. Reference is also made to Biographies of the persons mentioned. Extreme care has been bestowed on the verification of the dates, and thus numerous errors, current in previous works, have been corrected.

"An invaluable addition to our manuals of reference, and, from its moderate price, cannot fail to become as popular as it is useful."—TIMES.

Jephson.—SHAKESPEARE'S TEMPEST. With Glossarial and Explanatory Notes. By the Rev. J. M. JEPHSON. 18mo. 1s. 6d.

It is important to find some substitute for classical study, and it is believed that such a substitute may be found in the Plays of Shakespeare. Each sentence of Shakespeare becomes, like a sentence in Thucydides or Cicero, a lesson in the origin and derivation of words, and in the fundamental rules of grammatical construction. On this principle the present edition of the "Tempest" has been prepared. The text is taken from the "Cambridge Shakespeare."

Oppen.—FRENCH READER. For the Use of Colleges and Schools. Containing a graduated Selection from modern Authors in Prose and Verse; and copious Notes, chiefly Etymological. By EDWARD A. OPPEN. Fcap. 8vo. cloth. 4s. 6d.

This is a Selection from the best modern authors of France. Its distinctive feature consists in its etymological notes, connecting French with the classical and modern languages, including the Celtic. This subject has hitherto been little discussed even by the best-educated teachers.

A SHILLING BOOK OF GOLDEN DEEDS. A Reading Book for Schools and General Readers. By the Author of "The Heir of Redclyffe." 18mo. cloth.

A record of some of the good and great deeds of all time, abridged from the larger work of the same author in the Golden Treasury Series.

Vaughan, C. M. — A SHILLING BOOK OF WORDS FROM THE POETS. By C. M. VAUGHAN. 18mo. cloth.

It has been felt of late years that the children of our parochial schools, and those classes of our countrymen which they commonly represent, are capable of being interested, and therefore benefited also, by something higher in the scale of poetical composition than those brief and somewhat puerile fragments to which their knowledge was formerly restricted. An attempt has here been made to supply the want by forming a selection at once various and unambitious; healthy in tone, just in sentiment, elevating in thought, and beautiful in expression.

Thring.—Works by EDWARD THRING, M.A., Head Master of Uppingham.

THE ELEMENTS OF GRAMMAR TAUGHT IN ENGLISH,
with Questions. Fourth Edition. 18mo. 2s.

This little work is chiefly intended for teachers and learners. It took its rise from questionings in National Schools, and the whole of the first part is merely the writing out in order the answers to questions which have been used already with success. A chapter on Learning Language is especially addressed to teachers.

THE CHILD'S GRAMMAR. Being the Substance of "The Elements of Grammar taught in English," adapted for the Use of Junior Classes. A New Edition. 18mo. 1s.

SCHOOL SONGS. A Collection of Songs for Schools. With the Music arranged for four Voices. Edited by the Rev. E. THRING and H. RICCIUS. Folio. 7s. 6d.

There is a tendency in schools to stereotype the forms of life. Any genial solvent is valuable. Games do much; but games do not penetrate to domestic life, and are much limited by age. Music supplies the want. The collection includes the "Agnus Dei," Tennyson's "Light Brigade," Macaulay's "Ivry," &c. among other pieces.

Trench, Archbishop.—HOUSEHOLD BOOK OF ENGLISH POETRY. Selected and Arranged, with Notes, by R. C. TRENCH, D.D., Archbishop of Dublin. Extra fcap. 8vo. 5s. 6d.

This volume is called a "Household Book," by this name implying that it is a book for all—that there is nothing in it to prevent it from being confidently placed in the hands of every member of the household. Specimens of all classes of poetry are given, including selections from living authors. The Editor has aimed to produce a book "which the emigrant, finding room for little not absolutely necessary, might yet find room for it in his trunk, and the traveller in his knapsack, and that on some narrow shelves where there are few books this might be one."

"The Archbishop has conferred in this delightful volume an important gift on the whole English-speaking population of the world."—PALL MALL GAZETTE.

DIVINITY

Wright, W. Aldis (Co-Editor of the "Cambridge Shakespeare").
 —THE BIBLE WORD-BOOK. A Glossary of Old English
 Bible Words. By J. EASTWOOD, M.A., of St. John's College, and
 W. ALDIS WRIGHT, M.A., Trinity College, Cambridge. 18mo.
 5s. 6d.

It is the object of this Glossary to explain and illustrate all such words, phrases, and constructions, in the Authorized Version of the Old and New Testaments and the Apocrypha, and in the Book of Common Prayer, as are either obsolete or archaic. Full explanations are supplied, and these illustrated by numerous citations from the elder writers. An index of editions quoted is appended. Apart from its immediate subject, this work serves to illustrate a well-marked period in the history of the English language. It is thus of distinct philological value.

Hardwick.—A HISTORY OF THE CHRISTIAN CHURCH.
 Middle Age. From Gregory the Great to the Excommunication
 of Luther. By ARCHDEACON HARDWICK. Edited by FRANCIS
 PROCTER, M.A. With Four Maps constructed for this work by
 A. KEITH JOHNSTON. Second Edition. Crown 8vo. 10s. 6d.

The ground-plan of this treatise coincides in many points with one adopted at the close of the last century in the colossal work of Schröckh, and since that time by others of his thoughtful countrymen; but in arranging the materials a very different course has frequently been pursued. With regard to the opinions of the author, he is willing to avow distinctly that he has construed history with the specific prepossessions of an Englishman and a member of the English Church. The reader is constantly referred to the authorities, both original and critical, on which the statements are founded.

Hardwick, Archdeacon—continued.

- A HISTORY OF THE CHRISTIAN CHURCH DURING THE REFORMATION.** By ARCHDEACON HARDWICK. Revised by FRANCIS PROCTER, M.A. Second Edition. Crown 8vo. 10s. 6d.

This volume is intended as a sequel and companion to the "History of the Christian Church during the Middle Age." The author's earnest wish has been to give the reader a trustworthy version of those stirring incidents which mark the Reformation period, without relinquishing his former claim to characterise peculiar systems, persons, and events according to the shades and colours they assume, when contemplated from an English point of view, and by a member of the Church of England.

Maclear.—Works by the Rev. G. F. MACLEAR, B.D., Head Master of King's College School, and Preacher at the Temple Church.

- A CLASS-BOOK OF OLD TESTAMENT HISTORY.** Fourth Edition, with Four Maps. 18mo. cloth. 4s. 6d.

This volume forms a Class-book of Old Testament History from the earliest times to those of Ezra and Nehemiah. In its preparation the most recent authorities have been consulted, and wherever it has appeared useful, Notes have been subjoined illustrative of the Text, and, for the sake of more advanced students, references added to larger works. The Index has been so arranged as to form a concise dictionary of the persons and places mentioned in the course of the narrative; while the maps, which have been prepared with considerable care at Stanford's Geographical Establishment, will, it is hoped, materially add to the value and usefulness of the Book.

- A. CLASS-BOOK OF NEW TESTAMENT HISTORY,** including the Connexion of the Old and New Testament. With Four Maps. Third Edition. 18mo. cloth. 5s. 6d.

A sequel to the author's Class-book of Old Testament History, continuing the narrative from the point at which it there ends, and carrying it on to the close of St. Paul's second imprisonment at Rome. In its preparation, as in that of the former volume, the most recent and trustworthy authorities have been consulted, notes subjoined, and references to larger works added. It is thus hoped that it may prove at once an useful class-book and a convenient companion to the study of the Greek Testament.

Maclear, Rev. G. F.—*continued.*

A SHILLING BOOK OF OLD TESTAMENT HISTORY, for National and Elementary Schools. With Map. 18mo. cloth.

A SHILLING BOOK OF NEW TESTAMENT HISTORY, for National and Elementary Schools. With Map. 18mo. cloth.

These works have been carefully abridged from the author's larger manuals.

CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND. Second Edition. 18mo. cloth. 2s. 6d.

This may be regarded as a sequel to the Class-books of Old and New Testament History. Like them, it is furnished with notes and references to larger works, and it is hoped that it may be found, especially in the higher forms of our Public Schools, to supply a suitable manual of Instruction in the chief doctrines of the English Church, and a useful help in the preparation of Candidates for Confirmation.

A FIRST CLASS-BOOK OF THE CATECHISM OF THE CHURCH OF ENGLAND, with Scripture Proofs, for Junior Classes and Schools. 18mo. 6d.

THE ORDER OF CONFIRMATION. A Sequel to the Class Book of the Catechism. For the use of Candidates for Confirmation. With Prayers and Collects. 18mo. 3d.

Procter.—A HISTORY OF THE BOOK OF COMMON PRAYER, with a Rationale of its Offices. By FRANCIS PROCTER, M.A. Eighth Edition, revised and enlarged. Crown 8vo. 10s. 6d.

In the course of the last twenty years the whole question of Liturgical knowledge has been reopened with great learning and accurate research, and it is mainly with the view of epitomizing extensive publications, and correcting the errors and misconceptions which had obtained currency, that the present volume has been put together.

Procter and Maclear.—AN ELEMENTARY INTRODUCTION TO THE BOOK OF COMMON PRAYER.

Third Edition, re-arranged and supplemented by an Explanation of the Morning and Evening Prayer and the Litany. By the Rev. F. PROCTER and the Rev. G. F. MACLEAR. Third Edition. 18mo. 2s. 6d.

As in the other Class-books of the series, notes have also been subjoined, and references given to larger works, and it is hoped that the volume will be found adapted for use in the higher forms of our Public Schools, and a suitable manual for those preparing for the Oxford and Cambridge local examinations. This new Edition has been considerably altered, and several important additions have been made. Besides a re-arrangement of the work generally, the Historical Portion has been supplemented by an Explanation of the Morning and Evening Prayer and of the Litany.

PSALMS OF DAVID CHRONOLOGICALLY ARRANGED.

BY FOUR FRIENDS. An Amended Version, with Historical Introduction and Explanatory Notes. Crown 8vo. 10s. 6d.

To restore the Psalter as far as possible to the order in which the Psalms were written,—to give the division of each Psalm into strophes, of each strophe into the lines which composed it,—to amend the errors of translation, is the object of the present Edition. Professor Ewald's works, especially that on the Psalms, have been extensively consulted.

This book has been used with satisfaction by masters for private work in higher classes in schools.

Ramsay.—THE CATECHISER'S MANUAL; or, the Church

Catechism illustrated and explained, for the use of Clergymen, Schoolmasters, and Teachers. By the Rev. ARTHUR RAMSAY, M.A. Second Edition. 18mo. 1s. 6d.

A clear explanation of the Catechism, by way of question and answer.

Simpson.—AN EPITOME OF THE HISTORY OF THE CHRISTIAN CHURCH.

By WILLIAM SIMPSON, M.A. Fourth Edition. Fcap. 8vo. 3s. 6d.

A compendious summary of Church History.

Swainson.—A HANDBOOK to BUTLER'S ANALOGY. By C. A. SWAINSON, D.D., Norrisian Professor of Divinity at Cambridge. Crown 8vo. 1s. 6d.

This manual is designed to serve as a handbook or road-book to the Student in reading the Analogy, to give the Student a sketch or outline map of the country on which he is entering, and to point out to him matters of interest as he passes along.

Westcott.—A GENERAL SURVEY OF THE HISTORY OF THE CANON OF THE NEW TESTAMENT DURING THE FIRST FOUR CENTURIES. By BROOKE FOSS WESTCOTT, B.D., Canon of Peterborough. Second Edition, revised. Crown 8vo. 10s. 6d.

The Author has endeavoured to connect the history of the New Testament Canon with the growth and consolidation of the Church, and to point out the relation existing between the amount of evidence for the authenticity of its component parts, and the whole mass of Christian literature. Such a method of inquiry will convey both the truest notion of the connexion of the written Word with the living Body of Christ, and the surest conviction of its divine authority.

Of this work the Saturday Review writes: "Theological students, and not they only, but the general public, owe a deep debt of gratitude to Mr. Westcott for bringing this subject fairly before them in this candid and comprehensive essay. . . . As a theological work it is at once perfectly fair and impartial, and imbued with a thoroughly religious spirit; and as a manual it exhibits, in a lucid form and in a narrow compass, the results of extensive research and accurate thought. We cordially recommend it."

INTRODUCTION TO THE STUDY OF THE FOUR GOSPELS. By BROOKE FOSS WESTCOTT, B.D. Third Edition. Crown 8vo. 10s. 6d.

This book is intended to be an Introduction to the Study of the Gospels. The author has made it a point carefully to study the researches of the great writers, and consciously to neglect none. There is an elaborate discussion appended "On the Primitive Doctrine of Inspiration."

Westcott, Canon—continued.

A GENERAL VIEW OF THE HISTORY OF THE ENGLISH BIBLE. By BROOKE FOSS WESTCOTT, B.D. Crown 8vo. 1cs. 6d.

"The first trustworthy account we have had of that unique and marvellous monument of the piety of our ancestors."—DAILY NEWS.

THE BIBLE IN THE CHURCH. A Popular Account of the Collection and Reception of the Holy Scriptures in the Christian Churches. Second Edition. By BROOKE FOSS WESTCOTT, B.D. 18mo. cloth, 4s. 6d.

The present book is an attempt to answer a request, which has been made from time to time, to place in a simple form, for the use of general readers, the substance of the author's "History of the Canon of the New Testament." An elaborate and comprehensive Introduction is followed by chapters on the Bible of the Apostolic Age; on the Growth of the New Testament; the Apostolic Fathers; the Age of the Apologists; the First Christian Bible; the Bible Proscribed and Restored; the Age of Jerome and Augustine; the Bible of the Middle Ages in the West and in the East, and in the Sixteenth Century. Two appendices on the History of the Old Testament Canon before the Christian Era, and on the Contents of the most ancient MSS. of the Christian Bible, complete the volume.

THE GOSPEL OF THE RESURRECTION. Thoughts on its Relation to Reason and History. By BROOKE FOSS WESTCOTT, B.D. New Edition. Fcap. 8vo. 4s. 6d.

This Essay is an endeavour to consider some of the elementary truths of Christianity as a miraculous Revelation from the side of History and Reason. If the arguments which are here adduced are valid, they will go far to prove that the Resurrection, with all that it includes, is the key to the history of man, and the complement of reason.

Wilson.—AN ENGLISH, HEBREW, AND CHALDEE LEXICON AND CONCORDANCE, to the more Correct Understanding of the English translation of the Old Testament, by reference to the Original Hebrew. By WILLIAM WILSON, D.D., Canon of Winchester, late Fellow of Queen's College, Oxford. Second Edition, carefully Revised. 4to. cloth. 25s.

The aim of this work is, that it should be useful to clergymen and all persons engaged in the study of the Bible, even when they do not possess a knowledge of Hebrew; while able Hebrew scholars have borne testimony to the help that they themselves have found in it.

BOOKS ON EDUCATION.

Arnold.—A FRENCH ETON; OR, MIDDLE CLASS EDUCATION AND THE STATE. By MATTHEW ARNOLD. Fcap. 8vo. cloth. 2s. 6d.

"A very interesting dissertation on the system of secondary instruction in France, and on the advisability of copying the system in England."
SATURDAY REVIEW.

SCHOOLS AND UNIVERSITIES ON THE CONTINENT. 8vo. 10s. 6d.

The Author was in 1865 charged by the Schools Inquiry Commission with the task of investigating the system of education for the middle and upper classes in France, Italy, Germany, and Switzerland. In the discharge of this task he was on the continent nearly seven months, and during that time he visited the four countries named, and made a careful study of the matters to which the Commissioners had directed his attention. The present volume contains the report which he made to them. It is here adapted to the use of the general reader.

ESSAYS ON A LIBERAL EDUCATION. Edited by the Rev. F. W. FARRAR, M.A., F.R.S., Assistant Master at Harrow, late Fellow of Trinity College, Cambridge, and Hon. Fellow of King's College, London. Second Edition. 8vo. cloth. 10s. 6d.

CONTENTS:—*History of Classical Education, by Charles S. Parker, M.A.; Theory of Classical Education, by Henry Sedgwick, M.A.;*

Liberal Education in Universities, by John Seeley, M.A. ; *Teaching by means of Grammar*, by E. E. Bowen, M.A. ; *Greek and Latin Verse-Composition*, by the Rev. F. W. Farrar ; *Natural Science in Schools*, by J. M. Wilson, M.A., F.G.S. ; *The Teaching of English*, by J. W. Hales, M.A. ; *Education of the Reasoning Faculties*, by W. Johnson, M.A. ; *The present Social Results of Classical Education*, by Lord Houghton.

The Authors have sought to hasten the expansion and improvement of liberal education by showing in what light some of the most interesting questions of Educational Reform are viewed by men who have had opportunities for forming a judgment respecting them, and several of whom have been for some time engaged in the work of education at our Universities and Schools.

Farrar.—ON SOME DEFECTS IN PUBLIC SCHOOL EDUCATION. A Lecture delivered at the Royal Institution. With Notes and Appendices. Crown 8vo. 1s.

Jex-Blake.—A VISIT TO SOME AMERICAN SCHOOLS AND COLLEGES. By SOPHIA JEX-BLAKE. Crown 8vo. cloth. 6s.

"In the following pages I have endeavoured to give a simple and accurate account of what I saw during a series of visits to some of the Schools and Colleges in the United States. . . . I wish simply to give other teachers an opportunity of seeing through my eyes what they cannot perhaps see for themselves, and to this end I have recorded just such particulars as I should myself care to know."—AUTHOR'S PREFACE.

"Miss Blake gives a living picture of the Schools and Colleges themselves, in which that education is carried on."—PALL MALL GAZETTE.

Thring.—EDUCATION AND SCHOOL. By the Rev. EDWARD THRING, M.A., Head Master of Uppingham. Second Edition. Crown 8vo. cloth. 5s. 6d.

Youmans.—**MODERN CULTURE :** its True Aims and Requirements. A Series of Addresses and Arguments on the Claims of Scientific Education. Edited by EDWARD L. YOUMANS, M.D. Crown 8vo. 8s. 6d.

CONTENTS :—*Professor Tyndall on the Study of Physics ; Dr. Daubeny on the Study of Chemistry ; Professor Hensley on the Study of Botany ; Professor Huxley on the Study of Zoology ; Dr. J. Paget on the Study of Physiology ; Dr. Whewell on the Educational History of Science ; Dr. Faraday on the Education of the Judgment ; Dr. Hodgson on the Study of Economic Science ; Mr. Herbert Spencer on Political Education ; Professor Masson on College Education and Self Education ; Dr. Youmans on the Scientific Study of Human Nature. An Appendix contains extracts from distinguished authors, and from the Scientific Evidence given before the Public Schools Commission.*

